

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
3 January 2003 (03.01.2003)

PCT

(10) International Publication Number
WO 03/000851 A2

- (51) International Patent Classification⁷: **C12N**
- (21) International Application Number: **PCT/US02/19735**
- (22) International Filing Date: **20 June 2002 (20.06.2002)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:
60/300,501 22 June 2001 (22.06.2001) **US**
60/385,632 4 June 2002 (04.06.2002) **US**
- (71) Applicant (*for all designated States except US*): **THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; 3700 Market Street, Suite 300, Philadelphia, PA 19104-3147 (US).**
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): **GAO, Guangping [CN/US]; 408 Yorkshire Road, Rosemont, PA 19010 (US). WILSON, James, M. [US/US]; 1350 N. Avignon Drive, Gladwyne, PA 19035 (US).**
- (74) Agents: **KODROFF, Cathy, A. et al.; Howson and Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US).**
- (81) Designated States (*national*): **AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.**
- (84) Designated States (*regional*): **ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).**
- Published:**
- *without international search report and to be republished upon receipt of that report*
 - *with sequence listing part of description published separately in electronic form and available upon request from the International Bureau*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: **METHOD FOR RAPID SCREENING OF BACTERIAL TRANSFORMANTS AND NOVEL SIMIAN ADENOVIRUS PROTEINS**

(57) Abstract: Chimpanzee serotype C68 proteins, peptides, and polypeptide are provided. Also provided are novel adenoviruses derived from these proteins, as well as compositions containing these proteins and methods of using same for immunization and therapy. Further, a rapid method for screening recombinant transformants using a visually detectable method is described.

WO 03/000851 A2

METHOD FOR RAPID SCREENING OF BACTERIAL TRANSFORMANTS AND NOVEL SIMIAN ADENOVIRUS PROTEINS

BACKGROUND OF THE INVENTION

5 Recombinant adenoviruses have been described for gene therapy and vaccine uses.

 Adenoviruses have a characteristic morphology with an icosahedral capsid consisting of three major proteins, hexon (II), penton base (III) and a knobbed fibre (IV), along with a number of other minor proteins, VI, VIII, IX, IIIa and IVa2 [W.C. Russell, *J. Gen Virol.*, 81:2573-2604 (Nov 2000)]. The virus genome is a linear, double-stranded DNA
10 with a terminal protein attached covalently to the 5' termini, which have inverted terminal repeats (ITRs). The virus DNA is intimately associated with the highly basic protein VII and a small peptide termed *mu*. Another protein, V, is packaged with this DNA-protein complex and provides a structural link to the capsid via protein VI. The virus also contains a virus-
15 encoded protease, which is necessary for processing of some of the structural proteins to produce mature infectious virus.

 There continues to be a need for recombinant viral vectors and improved methods for making these vectors.

20 SUMMARY OF THE INVENTION

 In one aspect, the invention provides a method for rapid screening of bacterial transformants. The method involves engineering a recombinant shuttle vector comprising a nucleic acid cassette containing a transgene and a nucleic acid sequence encoding prokaryotic green fluorescent protein (GFP) operably linked to regulatory sequences which permit its
25 expression in a host cell. Thereafter, host cells are transfected with the shuttle vector and screened for expression of GFP. The absence of green color (i.e., white) is indicative of a cell carrying the recombinant virus. Expression of GFP is readily detected by the green color when activated by fluorescent light, and indicates the presence of parent virus (i.e., absence of recombinant).

In another aspect, the invention provides capsid proteins of C68, isolated from other C68 proteins, and characterized by the amino acids provided herein.

In still another embodiment, the invention provides adenoviral vectors and non-viral targeting proteins derived from the C68 capsid proteins, termed herein C68-derived
5 constructs.

Yet other advantages of the present invention will be readily apparent from the following detailed description of the invention.

Brief Description of the Drawings

10 Fig. 1 summarizes the genetic organization of the chimpanzee adenovirus C68 genome. In Fig. 1A the genome of the C68 chimpanzee adenovirus is schematically represented by the box at the top. The inverted terminal repeats are shaded black and the early regions are shaded gray. The arrowheads above the box indicate the direction of expression of the early genes. The line below the box represents the division of the genome
15 into 100 map units. The arrows below the line represent the five late gene regions and the proteins encoded in each region. The numbers below the box or arrows indicate the start (promoter or initiation codon) and end (canonical PolyA signal) for each region. * represents the E2A late promoter. Fig. 1B illustrates the PstI clones; Fig. 1C illustrates the BamHI clones. Fig. 1D illustrates the HindIII clones. For parts 1B-1D, the unshaded regions indicate
20 that a fragment was cloned into a plasmid vector, while the shaded regions indicate that the restriction fragment was not cloned. For each section the fragment name, alphabetical with A being the largest fragment, and the fragment size are listed above the box and the fragment end points are listed below the box.

Fig. 2 provides a sequence alignment of the C68 hexon protein [aa 131 to 441 of SEQ
25 ID NO:16] with Ad4 [SEQ ID NO:34], Ad16 [SEQ ID NO:35], Ad3 [SEQ ID NO:36], Ad7 [SEQ ID NO:37], and Ad2 [SEQ ID NO:38]. The deduced amino acid sequences of highly similar human adenovirus hexons were compared with the C68 chimpanzee adenovirus using CLUSTAL X. Serotypes and subgroups are indicated on the left margin, followed by the residue number. The numbering refers to the amino acid position with respect to the start of
30 translation. Amino acids are shaded with respect to C68 to highlight sequence similarities

(gray) and identities (black). The seven hypervariable regions within loop domains DE1 and FG1 are labeled along the bottom and correspond to the following Ad2 sequences in the alignment: HVR1, 137-188; HVR2, 194-204; HVR3, 222-229; HVR4, 258-271; HVR5, 278-294; HVR6, 316-327; and HVR7, 433-465 of SEQ ID NO:16. The GenBank accession numbers for the sequences shown are as follow: AAD03657 (Ad4), S37216 (Ad16), S39298 (Ad3), AAD03663 (Ad7), and NP040525 (Ad2).

Fig. 3 provides an alignment of the amino acid sequences of the fiber knob domains of chimpanzee C68 (Pan-9) [amino acids 247 to 425 of SEQ ID NO: 27] and the human adenovirus serotypes 2 [SEQ ID NO: 39] and 5[SEQ ID NO:40].

Fig. 4 provides an alignment of the amino acid sequences of the L1 and a portion of the L2 loops of the capsid hexon on the human adenovirus serotype 5 [SEQ ID NO:41] and chimpanzee C68 (Pan-9) [amino acids 125 to 443 of SEQ ID NO:16] adenovirus sequences. The intervening conserved region is part of the pedestal domain conserved between adenovirus serotypes.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel adenovirus capsid proteins derived from the unique sequences of chimpanzee adenovirus C68. The capsid proteins of the invention are useful for a variety of purposes, including non-viral targeted delivery to cells and for creating recombinant viral vectors. These proteins and viral vectors are useful for delivery of heterologous molecules to target cells.

The invention further provides a novel method for rapid screening of bacterial transformants obtained during production of the novel adenoviral capsids of the invention, and during production of a variety of other viral or non-viral constructs. In this method, at least the shuttle vector is engineered to contain a marker gene, e.g., green fluorescent protein (GFP), gene under the control of a suitable promoter. The transformed cells are screened for expression of marker. In the case of GFP, white colonies are recombinants while green colonies are residual parental plasmid.

I. Novel Adenovirus Capsid Proteins

In one aspect, the invention provides unique C68 adenoviral capsid proteins, including the C68 hexon region, the C68 penton region, and the C68 fiber region, and fragments thereof. Suitably, these capsid proteins can be substantially pure, i.e., are free of other proteins. Preferably, these proteins are at least 10% homogeneous, more preferably 60% homogeneous, and most preferably 95% homogeneous.

In addition, the invention provides unique C68-derived capsid proteins. As used herein, a C68-derived capsid protein includes any C68 capsid protein or a fragment thereof including, without limitation, a polypeptide, peptide or a consecutive sequence of at least 8 amino acid residues unique to a C68 capsid protein and which is free of other proteins. A C68-derived capsid protein also includes a capsid protein that contains a C68 capsid protein or fragment thereof as defined above, including, without limitation, a chimeric capsid protein, a fusion protein, an artificial capsid protein, a synthetic capsid protein, and a recombinant capsid proteins, without limitation to means of generating these proteins. Suitably, these C68-derived capsid proteins contain one or more C68 regions or fragments thereof (e.g., a hexon) in combination with capsid regions or fragments thereof of different adenoviral serotypes, or of non-adenoviral sources, as described herein. These C68-derived capsid proteins may be used in non-viral targeting of useful molecules to cells, or for production of viral vectors, as described herein.

A "modification of a capsid protein associated with altered tropism" as used herein includes an altered capsid protein, i.e, a penton, hexon or fiber protein region, or fragment thereof, such as the knob domain of the fiber region, or a polynucleotide encoding same, such that specificity is altered.

In one embodiment, the amino acid sequences of the C68 penton protein are provided

in SEQ ID NO:12: MMRRAYPEGPPPSYESVMQQAMAAAAMQPPLEAPYVPPRYLAPT
EGRNSIRYSELAPLYDTTRLYLVDNKSADIASLNYQNDHSNFLTTVVQNNDFTPTEAS
TQTINFDESRWGGQLKTIMHTNMPNVNEFMYSNKFARVMVSRKTPNGVTVTEDYDG
SQDELKYEWFELPEGNFVMTMTIDLNNAIIDNYLAVGRQNGVLES DIGVKFDTRN
FRLGWDPVTELVMPGVYTNEAFHPDIVLLPGCGVDFTESRLSNLLGIRKRQPFQEGFQ
IMYEDLEGGNIPALLDVDAYEKSKEAATAAVATASTEVRGDNFASAAVAAAEA
AETESKIVIQPEKDSKNRSYNVLPDKINTAYRSWYLAYNYGDPEKGVRSWTLTSD

VTCGVEQVYWSLPDMMQDPVTFRSTRQVSNPVVGAE LLPVYSKSFNEQAVYSQQLR
 AFTSLTHVFNRFENQILVRPPAPTITTVSENPALTDHGTLPLRSSIRGVQRVTVD
 ARRRTCPIVYKALGIVAPRVLSSRTF.

Suitably, this penton protein, or unique fragments thereof, may be utilized for a
 5 variety of purposes. Examples of suitable fragments include the C68 penton having N-
 terminal and/or C-terminal truncations of about 50, 100, 150, or 200 amino acids, based upon
 the amino acid numbering provided above and in SEQ ID NO:12. Other suitable fragments
 include shorter internal, C-terminal, or N-terminal fragments. Further, the penton protein
 may be modified for a variety of purposes known to those of skill in the art.

10 The sequences of the C68 hexon are provided in SEQ ID NO:16:

MATPSMLPQWAYMHIAGQDASEYLSPLVQFARATDTYFSLGNK
 FRNPTVAPTHDVTDRSQRLTLRFVPVDREDNTYSYKVRTLAVGDNRVLDMASTYFD
 15 IRGVLDRGPSFKPYSGTAYNSLAPKGAPNTCQWTYKADGETATEKTYTYGNAPVQGIN
 ITKDGILGTDTDQPIYADKTYQPEPQVGDAEWHDTGTDEKYGGRALKPDTKMKPC
 20 YGSFAKPTNKEGGQANVKTGTGTTKEYDIDMAFFDNRSAAAAGLAPEIVLYTENVDL
 TPDTHIVYKAGTDDSSSSINLGQQAMPNRPNIIGFRDNFIGLMYYNSTGNMGVLAGQA
 25 SQLNAVVDLQDRNTELSYQLLLDSLGRTRYFSMWNQAVDSYDPDVRIENHGVDEL
 PNYCFPLDAVGRTDTYQGIKANGTDQTTWTKDDSVNDANEIGKGNPFAMEINIQANLW
 RNFLYANVALYLPDSYKYTPANVTLPNTNTYDYMNGRVVAPSLVDSYINIGARWSLD
 30 PMDNVNPFNHHRNAGLRYRSMLLGNRGYVPFHIQVPQKFFAIKSLLLPGSYTYEWNF
 RKDVNMILQSSLGNDLRTDGASISFTSINLYATFFPMAHNTASTLEAMLRNDTNDQSF
 NDYLSAANMLYPIPANATNPISIPSRNWA AFRGWSFTRLKTETPSLGSFGDPYFVY
 35 SGSIPYLDGTFYLNHTFKKVSITFDSSVSWPGNDRLLTPNEFEIKRTVDGEGYNVAQC
 NMTKDWFLVQMLAHYNIGYQGFYVPEGYKDRMYSFFRNFPMSRQVVDEVNYKDYQAV
 40 TLAYQHNNSGFVGYLAPTMRQGQPYXPYPLIGKSAVTSVTQKKFLCDRVMWRIPF
 SSNFMMSGALTDLGQNMLYANSAHALDMNFEVDPMDESTLLYVVFVDFVVRVHQPHR
 GVIEAVYXRTPFSAGNATT.

45 Suitably, this hexon protein, or unique fragments thereof, may be utilized for a variety
 of purposes. Examples of suitable fragments include the C68 hexon having N-terminal
 and/or C-terminal truncations of about 50, 100, 150, 200, 300, 400, or 500 amino acids, based
 upon the amino acid numbering provided above and in SEQ ID NO:16. Other suitable

fragments include shorter internal, C-terminal, or N-terminal fragments. For example, one suitable fragment the loop region (domain) of the hexon protein, designated DE1 and FG1, or a hypervariable region thereof. Such fragments include the regions spanning amino acid residues about 125 to 443; about 138 to 441, or smaller fragments, such as those spanning
 5 about residue 138 to residue 163; about 170 to about 176; about 195 to about 203; about 233 to about 246; about 253 to about 264; about 287 to about 297; and about 404 to about 430 of C68, with reference to SEQ ID NO:16. Other suitable fragments may be readily identified by one of skill in the art. Further, the hexon protein may be modified for a variety of purposes known to those of skill in the art.

10 In one example, it may be desirable to generate an adenovirus having an altered hexon protein utilizing the C68 hexon protein sequences of the invention. One suitable method for altering hexon proteins is described in US Patent 5,922,315, which is incorporated by reference. In this method, at least one loop region of the adenovirus hexon is changed with at least one loop region of another adenovirus serotype. Thus, at least one loop region of such
 15 an altered adenovirus hexon protein is a C68 hexon loop region. In one embodiment, a loop region of the C68 hexon protein is replaced by a loop region from another adenovirus serotype. In another embodiment, the loop region of the C68 hexon is used to replace a loop region from another adenovirus serotype. Suitable adenovirus serotypes may be readily selected from among human and non-human serotypes, as described herein. Where non-
 20 human adenoviruses are selected, the serotypes are preferably selected from non-human primates. However, the selection of a suitable serotype is not a limitation of the present invention. Still other uses for the C68 hexon protein sequences of the invention will be readily apparent to those of skill in the art.

The sequences of the C68 fiber protein are: SEQ ID NO:27:

25 MSKKRVRVDDDFDPVYPYDADNAPTVPFINPPFVSSDGFQEKPL
 GVLSRLRLADPVTTKNGEITLKLGEVDLDSSGKLISNTATKAAAPLSFSNNTISLNMD
 PFYTKDGKLSLQVSPPLNILRTSILNTLALGFGSGLGLRGSALAVQLVSPLTFD TDGN
 IKLTLDRLHVTTGDAIESNISWAKGLKFEDGAIATNIGNGLEFGSSSTETGVDDAY
 30 PIQVKLGSGLSFDSTGAIMAGNKEDDKLTLTWTPDPSPNCQILAENDAKLTLCLTKCG
 SQILATVSVLVVSGNLNPITGTVSSAQVFLRFDANGVLLTEHSTLKKYWG YRQGDSI
 DGTPTYNAVGFMPNLKAYPKSQSSTTKNNIVGQVYMNGDVSKPMLLTITLNGTDDSNS
 TYSMSFSYTWNGSYVGATFGANSYTF SYIAQE.

Suitably, this fiber protein, or unique fragments thereof, may be utilized for a variety of purposes. One suitable fragment is the fiber knob, which spans about amino acids 247 to 425 of SEQ ID NO: 27. Examples of other suitable fragments include the C68 fiber having N-terminal and/or C-terminal truncations of about 50, 100, 150, or 200 amino acids, based upon the amino acid numbering provided above and in SEQ ID NO:27. Still other suitable fragments include internal fragments. Further, the fiber protein may be modified using a variety of techniques known to those of skill in the art.

The amino acid sequences of other useful gene products of C68 are provided in SEQ ID Nos. 1 – 11, 13 – 15, 17 – 26, and 28 -38 of the attached sequence listing. More particularly, these sequences are as follows.

Regions		Ad C68 – CDS, With ref to SEQ ID NO:33.	Ad C68 SEQ ID NO:
E1a	11kDa	578... 649, 1236 ... 1469	1
	28.2 kDa	578 ... 1142, 1236 ... 1444	2
	24.8 kDa	578... 1049, 1236 ... 1444	3
E1b	20.5kDa	1603 ... 2163	4
	54.7 kDa	1908 ... 3404	5
	18.5 kDa	1908 ... 2200, 3188 ... 3404	6
	10.1 kDa	1908 ... 2170, 3306 ... 3324	7
IX	Hexon-associated protein – pIX	3489 ... 3917	8
IVa2	Maturation protein – pIVa2	Complement (3976 ... 5309, 5588 ... 5600)	9
L1	21.9kDa	7858 ... 8460	10
	42.9 kDa	10825 ... 12000	11
L2	Penton – pIII	13888 ... 15492	12
	Major core protein – pVIII	15493 ... 16098	13
	Minor Core Protein – pV	16120 ... 17190	14
L3	Hexon-associated protein – pVI	17442 ... 18215	15
	Hexon – pII	18322 ... 21123	16
E2a	DNA-Binding Protein Endopeptidase	Complement (21835 ... 23376)	17
L4	Virion morphogenesis-associated protein 24.3 kDa	Complement (25529 ... 25862, 26032 ... 26366)	18
	Hexon-associated protein - pVIII	26446 ... 27129	19

Regions		Ad C68 – CDS, With ref to SEQ ID NO:33.	SEQ ID NO:
E3	11.6 kDa	27130 ... 27450	20
	16 kDa	(27404 ... 27477, 27666 ... 28032)	21
	19.3 kDa	28014 ... 28544	22
	22.3	28572 ... 29186	23
	9.9 kDa	30722 ... 30997	24
	15.6 kDa	31003 ... 31434	25
	14.7 kDa	31427 ... 31834	26
L5	Fiber - pIV	32137 ... 33414	27
E4	ORF7-like protein	Complement (33521 ... >33772)	28
	Orf 6 – 33 kDa	Complement (33769 ... 34674)	29
	Orf4 – 13.2 kDa	Complement (34580 ... 34945)	30
	Orf 3 – 12. 8 kDa	Complement (34955 ... 35308)	31
	Orf 2 – 14.2 kDa	Complement (35305 ... 35694)	32

Thus, the invention provides unique C68 proteins, peptides and fragments thereof, which are produced recombinantly or by other methods. Suitably, such fragments are at least 8 amino acids in length. However, fragments of other desired lengths are readily utilized. In addition, the invention encompasses such modifications as may be introduced to enhance yield and/or expression of a C68 protein or fragment, construction of a fusion molecule in which all or a fragment of the C68 protein or fragment is fused (either directly or via a linker) with a fusion partner to enhance. Other suitable modifications include, without limitation, truncation of a coding region (e.g., a protein or enzyme) to eliminate a pre- or pro-protein ordinarily cleaved to produce the mature protein or enzyme and/or mutation of a coding region to provide a secretable gene product. Still other modifications will be readily apparent to one of skill in the art. The invention further encompasses proteins having at least about 95% to 99% identity to the C68 proteins provided herein.

The term “substantial homology” or “substantial similarity,” when referring to a protein or fragment thereof, indicates that, when optimally aligned with appropriate amino acid insertions or deletions with another protein, there is nucleotide sequence identity in at least about 95 to 99% of the aligned sequences .

The term “percent sequence identity” or “identical” in the context of proteins or fragments thereof refers to the amino acids in the two sequences that are the same when

aligned for maximum correspondence. The length of sequence identity comparison may be over the full length of a protein, enzyme, polypeptide, peptide, or other fragment of at least about 200 to 500 amino acids, is desired. However, identity among smaller fragments, e.g. of at least about 8 amino acids, usually at least about 20 to 24 amino acids, at least about 28 to 32 amino acids, at least about 50 or more amino acids, may also be desired.

Identity is readily determined by one of skill in the art by resort to algorithms and computer programs known by those of skill in the art. As described herein, alignments are performed using any of a variety of publicly or commercially available Multiple Sequence Alignment Programs, such as "Clustal W", accessible through Web Servers on the internet.

Alternatively, Vector NTI utilities are also used. There are also a number of algorithms known in the art that can be used to measure amino acid sequence identity, including those contained in the programs described above. Generally, these programs are used at default settings, although one of skill in the art can alter these settings as needed. Alternatively, one of skill in the art can utilize another algorithm or computer program that provides at least the level of identity or alignment as that provided by the referenced algorithms and programs.

As described herein, the C68-derived capsid proteins of the invention are particularly well suited for use in applications in which the neutralizing antibodies diminish the effectiveness of other Ad serotype based targeting proteins and vectors, as well as other viral vectors. The C68-derived constructs of the invention are particularly advantageous in readministration for repeat gene therapy or for boosting immune response (vaccine titers).

Also provided by the present invention are artificial adenoviral capsid proteins, which involve modifications and chimeric capsids constructed using the C68 adenoviral capsid proteins of the invention. Such artificial capsid proteins can be constructed using the amino acid sequences of the chimp C68 Ad hexon of the invention. Because the hexon protein is the determinant for serotype of an adenovirus, such artificial hexon proteins would result in adenoviruses having artificial serotypes. Other artificial capsid proteins can also be constructed using the chimp Ad penton sequences and/or fiber sequences of the invention and/or fragments thereof.

In one embodiment, a chimeric C68 capsid is constructed using C68 hexon and C68 fiber and a penton from another adenovirus. Alternatively, a chimeric C68 capsid comprises

a C68 hexon and a fiber and penton from one or more different adenoviruses. Another chimeric adenovirus capsid comprises the C68 fiber and a penton and a hexon from one or more different different adenovirus serotypes. Yet another chimeric adenovirus capsid comprises the C68 penton and a fiber and hexon from one or more different adenovirus serotypes. Suitably, for such chimeric and artificial capsids constructed from C68 proteins, the non-C68 adenovirus components may be readily selected from other adenovirus serotypes.

Under certain circumstances, it may be desirable to use one or more of the C68-derived capsid proteins or a fragment thereof to generate an antibody. The term "an antibody," as used herein, refers to an immunoglobulin molecule which is able to specifically bind to an epitope. The antibodies in the present invention exist in a variety of forms including, for example, high affinity polyclonal antibodies, monoclonal antibodies, synthetic antibodies, chimeric antibodies, recombinant antibodies and humanized antibodies. Such antibodies originate from immunoglobulin classes IgG, IgM, IgA, IgD and IgE. Such antibodies may be generated using any of a number of methods known in the art. Suitable antibodies may be generated by well-known conventional techniques, e.g. Kohler and Milstein and the many known modifications thereof. Similarly desirable high titer antibodies are generated by applying known recombinant techniques to the monoclonal or polyclonal antibodies developed to these antigens [see, e.g., PCT Patent Application No. PCT/GB85/00392; British Patent Application Publication No. GB2188638A; Amit *et al.*, 1986 *Science*, 233:747-753; Queen *et al.*, 1989 *Proc. Nat'l. Acad. Sci. USA*, 86:10029-10033; PCT Patent Application No. PCT/WO9007861; and Riechmann *et al.*, *Nature*, 332:323-327 (1988); Huse *et al.*, 1988a *Science*, 246:1275-1281]. Alternatively, antibodies can be produced by manipulating the complementarity determining regions of animal or human antibodies to the antigen of this invention. See, e.g., E. Mark and Padlin, "Humanization of Monoclonal Antibodies", Chapter 4, The Handbook of Experimental Pharmacology, Vol. 113, The Pharmacology of Monoclonal Antibodies, Springer-Verlag (June, 1994); Harlow *et al.*, 1999, Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, NY; Harlow *et al.*, 1989, Antibodies: A Laboratory Manual, Cold Spring Harbor, New York;

Houston *et al.*, 1988, *Proc. Natl. Acad. Sci. USA* 85:5879-5883; and Bird *et al.*, 1988, *Science* 242:423-426.

Alternatively, one or more of the C68 capsid proteins of the invention are assembled as multi-antigenic complexes [see, e.g., European Patent Application 0339695, published November 2, 1989] and employed to elicit high titer antibodies. Further provided by the present invention are anti-idiotypic antibodies (Ab2) and anti-anti-idiotypic antibodies (Ab3). See, e.g., M. Wettendorff *et al.*, "Modulation of anti-tumor immunity by anti-idiotypic antibodies." In *Idiotypic Network and Diseases*, ed. by J. Cerny and J. Hiernaux, 1990 *J. Am. Soc. Microbiol.*, Washington DC: pp. 203-229]. These anti-idiotypic and anti-anti-idiotypic antibodies are produced using techniques well known to those of skill in the art. These antibodies may be used for a variety of purposes, including diagnostic and clinical methods and kits.

Under certain circumstances, it may be desirable to introduce a detectable label or a tag onto a C68 antibody or other construct of the invention. As used herein, a detectable label is a molecule which is capable, alone or upon interaction with another molecule, of providing a detectable signal. Most desirably, the label is detectable visually, e.g. by fluorescence, for ready use in immunohistochemical analyses or immunofluorescent microscopy. For example, suitable labels include fluorescein isothiocyanate (FITC), phycoerythrin (PE), allophycocyanin (APC), coriphosphine-O (CPO) or tandem dyes, PE-cyanin-5 (PC5), and PE-Texas Red (ECD). All of these fluorescent dyes are commercially available, and their uses known to the art. Other useful labels include a colloidal gold label. Still other useful labels include radioactive compounds or elements. Additionally, labels include a variety of enzyme systems that operate to reveal a colorimetric signal in an assay, e.g., glucose oxidase (which uses glucose as a substrate) releases peroxide as a product which in the presence of peroxidase and a hydrogen donor such as tetramethyl benzidine (TMB) produces an oxidized TMB that is seen as a blue color. Other examples include horseradish peroxidase (HRP) or alkaline phosphatase (AP), and hexokinase in conjunction with glucose-6-phosphate dehydrogenase which reacts with ATP, glucose, and NAD⁺ to yield, among other products, NADH that is detected as increased absorbance at 340 nm wavelength. Other label systems that are utilized in the methods of this invention are detectable by other means, e.g., colored

latex microparticles [Bangs Laboratories, Indiana] in which a dye is embedded are used in place of enzymes to form conjugates with the target sequences provide a visual signal indicative of the presence of the resulting complex in applicable assays.

Methods for coupling or associating the label with a desired molecule are similarly conventional and known to those of skill in the art. Known methods of label attachment are described [see, for example, Handbook of Fluorescent probes and Research Chemicals, 6th Ed., R. P. M. Haugland, Molecular Probes, Inc., Eugene, OR, 1996; Pierce Catalog and Handbook, Life Science and Analytical Research Products, Pierce Chemical Company, Rockford, IL, 1994/1995]. Thus, selection of the label and coupling methods do not limit this invention.

The C68-derived proteins, peptides, and fragments described herein can be produced by any suitable means, including chemical synthesis, or other synthetic means, or by recombinant production and conventional genetic engineering methodologies. For example, peptides can be synthesized by the well known solid phase peptide synthesis methods (Merrifield, *J. Am. Chem. Soc.*, **85**:2149 (1962); Stewart and Young, *Solid Phase Peptide Synthesis* (Freeman, San Francisco, 1969) pp. 27-62). These and other suitable production methods are within the knowledge of those of skill in the art and are not a limitation of the present invention.

Alternatively, suitable methods for recombinant production can be used. Selection of suitable expression systems, including expression vectors and host cells for protein expression and/or viral packaging is within the ability of one of skill in the art and is not a limitation of the present invention. See, e.g., Sambrook et al, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (Cold Spring Harbor, NY).

Nucleic acid sequences for the C68 genome, which is 36521 bp in length, may be obtained using information available in US Patent 6,083,716 and from the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 (Pan-9). This sequences is also available from GenBank. Other chimpanzee adenovirus sequences are available from the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209, and other sources. Desirable chimpanzee strains Pan 5 [ATCC VR-591], Pan 6 [ATCC VR-592], and Pan 7 [ATCC VR-593]. Another particularly

desirable chimpanzee adenovirus strain is chimpanzee adenovirus strain Bertha or C1 [ATCC Accession No. VR-20]. The sequence of the C1 serotype, and the location of the adenovirus genes E1a, E1b, E2a, E2b, E3, E4, L1, L2, L3, L4 and L5 are provided in US Patent 6,083,716, which is incorporated by reference herein. Optionally, non-chimpanzee simian
5 adenoviral sequences may be used. Such non-chimpanzee adenovirus include those obtained from baboon adenovirus strains [e.g., ATCC VR-275], adenovirus strains isolated from rhesus monkeys [e.g., ATCC VR-209, ATCC VR-275, ATCC VR-353, ATCC VR-355], and adenovirus strains isolated from African green monkeys [e.g., ATCC VR-541; ATCC VR-941; ATCC VR-942; ATCC VR-943]. Alternatively, one may readily select from among the
10 at least 51 different human serotypes, including, without limitation, human adenovirus serotypes 1, 2, 3, 4, 5, 12, 35, 37, and 40, and other, non-human primate adenovirus serotypes. Further, the sequences of these and other suitable serotypes are available from a variety of databases including, e.g., PubMed and GenBank [see, for example, US Patent No. 5,240,846]. Selection of an appropriate adenovirus is not a limitation of the present
15 invention.

The invention further provides molecules useful for production of the C68 and C68-derived proteins of the invention, including such molecules which carry polynucleotides including DNA sequences. Thus, the invention further encompasses the nucleic acid sequences encoding the C68-derived constructs of the invention, and molecules and host cells
20 useful in expression thereof, including suitable DNA molecules and vectors, which can be any suitable genetic element as defined herein. Preferably, these vectors are DNA-based (e.g., plasmids) or viral vectors.

In one embodiment, the C68-derived capsid proteins and other C68 adenovirus proteins described herein are used for non-viral, protein-based delivery of genes, proteins, and
25 other desirable diagnostic, therapeutic and immunogenic molecules. A desired molecule for delivery to a target cell may be associated with a C68-derived capsid protein or other protein by any suitable means, including, e.g., covalent or non-covalent binding. For example, the C68 penton protein may be readily utilized for such a purpose by production of a fusion protein using the C68 penton sequences of SEQ ID NO:12 in a manner analogous to that
30 described in Medina-Kauwe LK, et al, *Gene Ther.* 2001 May; 8(10):795-803 and Medina-

Kauwe LK, et al, *Gene Ther.* 2001 Dec; 8(23): 1753-1761. Alternatively, the amino acid sequences of C68 protein IX may be utilized for targeting vectors by associating the protein IX with a ligand that binds to a cell surface receptor, as described in US Patent Appln 20010047081. Suitable ligands include a CD40 antigen, an RGD-containing or polylysine-containing sequence, and the like. Still other C68 proteins may be used for these and similar purposes.

Further, the C68 adenovirus proteins of the invention are particularly well suited for use in producing viral vectors in C68-derived capsids. Suitably, these adenoviruses are pseudotyped such that a nucleic acid molecule carrying adenovirus ITRs from a non-C68 serotype and a minigene are packaged in a C68-derived adenoviral capsid of the invention. Alternatively, adenoviruses may be generated which contain at least the 5' ITRs or the 3' ITRs from C68, in a C68-derived capsid protein. The adenoviral vectors described herein may contain adenoviral sequences derived from one, more than one adenoviral strain. In yet another alternative, other C68 elements described herein may be utilized in production of recombinant vectors, or other desirable constructs.

The C68 proteins of the invention are useful for a variety of purposes, including construction of recombinant viruses. The C68-derived capsid proteins of the invention are useful in producing hybrid vectors, including, hybrid C68-adeno-associated viruses, Epstein-Barr virus, and retroviruses [Caplen *et al*, *Gene Ther.* 6: 454-459 (1999); Tan *et al*, *J Virol.*, 73:7582-7589 (1999)]. Such viruses include C68-derived capsids which encapsidated vectors with adeno-associated virus (AAV) ITRs [Lieber *et al*, *J Virol*, 73:9314-9324 (1999), Recchia *et al*, *Proc Natl Acad Sci USA*, 96:2615-2620 (1999); or lentivirus ITRs (Zheng *et al*, *Nat Biotech*, 18:176-180 (2000), using Maloney leukemia virus long terminal repeats).

In a particularly desirable embodiment, the C68-derived capsid proteins, and optionally, the other C68 sequences described herein, are used to produce recombinant adenoviruses and pseudotyped adenoviruses. However, it will be readily understood that the C68-derived capsid proteins and other novel C68 sequences can be utilized for a variety of purposes, including production of other types of viral vectors (such as, e.g., hybrid vectors) carrying the therapeutic and immunogenic transgenes described below. Additionally, it will be readily understood that viral vectors carrying the unique C68 proteins and other sequences

of the invention can be utilized for targeting and/or delivery of other types of molecules, including proteins, chemical molecules and other moieties useful for diagnostic, therapeutic and/or immunization purposes.

5 II. Recombinant Adenoviral Vectors

The compositions of this invention include vectors that deliver a heterologous molecule to cells, either for therapeutic or vaccine purposes. As used herein, a vector may include any genetic element including, without limitation, a cosmid, episome, plasmid, or a virus. In a particularly preferred embodiment, these vectors are viral vectors having capsid
10 proteins derived from the C68 proteins of the invention. Alternatively, these vectors may contain other C68 sequences of the invention. These viral vectors suitably contain a minigene. By "minigene" is meant the combination of a selected heterologous gene and the other regulatory elements necessary to drive translation, transcription and/or expression of the gene product in a host cell.

15 Typically, an adenoviral vector is designed such that the minigene is flanked on its 5' end and/or its 3' end by adenoviral sequences which include, at a minimum, the cis-elements necessary for replication and virion encapsidation. Thus, in one embodiment, the vector contains adenoviral sequences encompassing at least the 5' end of the adenoviral genome, i.e., the 5' inverted terminal repeat sequences (which functions as origins of replication) and
20 the native 5' packaging enhancer domains (that contain sequences necessary for packaging linear Ad genomes and enhancer elements for the E1 promoter). The vector is also provided with the cis-acting 3' ITRs. Suitably, the minigene is located between the 5' adenoviral elements and the 3' adenoviral elements. An adenoviral vector of the invention may also contain additional adenoviral sequences. For example, the minigene may be located in the
25 site of such as the site of a functional E1 deletion or functional E3 deletion, among others that may be selected. Alternatively, the minigene may be inserted into an existing gene region to disrupt the function of that region, if desired.

The term "functionally deleted" or "functional deletion" means that a sufficient amount of the gene region is removed or otherwise damaged, e.g., by mutation or

modification, so that the gene region is no longer capable of producing functional products of gene expression. If desired, the entire gene region may be removed.

Suitably, these adenoviral vectors of the invention contain one or more adenoviral elements derived from C68. In one embodiment, the vectors contain adenoviral ITRs from an adenoviral serotype which differs from C68. Alternatively, C68 ITRs may be utilized in a viral vector of the invention in which the capsid is not naturally occurring, but contains one or more C68 proteins, or fragments thereof. The selection of the serotype of the ITRs and the serotype of any other adenoviral sequences present in vector is not a limitation of the present invention. A variety of adenovirus strains are described herein.

The viral sequences, helper viruses, if needed, and recombinant viral particles, and other vector components and sequences employed in the construction of the vectors described herein are obtained as described above. See, e.g., US Patent No. 5,240,846. The DNA sequences of the adenovirus sequences are employed to construct vectors and cell lines useful in the preparation of such vectors. See, e.g., US Patent No. 6,083,716.

Modifications of the nucleic acid sequences forming the vectors of this invention, including sequence deletions, insertions, and other mutations may be generated using standard molecular biological techniques and are within the scope of this invention.

A. The "Minigene"

The methods employed for the selection of the transgene, the cloning and construction of the "minigene" and its insertion into the viral vector are within the skill in the art given the teachings provided herein.

1. The transgene

The transgene is a nucleic acid sequence, heterologous to the vector sequences flanking the transgene, which encodes a polypeptide, protein, or other product, of interest. The nucleic acid coding sequence is operatively linked to regulatory components in a manner which permits transgene transcription, translation, and/or expression in a host cell.

The composition of the transgene sequence will depend upon the use to which the resulting vector will be put. For example, one type of transgene sequence includes a reporter sequence, which upon expression produces a detectable signal. Such reporter sequences include, without limitation, DNA sequences encoding β -lactamase, β -galactosidase

(LacZ), alkaline phosphatase, thymidine kinase, green fluorescent protein (GFP), chloramphenicol acetyltransferase (CAT), luciferase, membrane bound proteins including, for example, CD2, CD4, CD8, the influenza hemagglutinin protein, and others well known in the art, to which high affinity antibodies directed thereto exist or can be produced by

5 conventional means, and fusion proteins comprising a membrane bound protein appropriately fused to an antigen tag domain from, among others, hemagglutinin or Myc. These coding sequences, when associated with regulatory elements which drive their expression, provide signals detectable by conventional means, including enzymatic, radiographic, colorimetric, fluorescence or other spectrographic assays, fluorescent activating cell sorting assays and
10 immunological assays, including enzyme linked immunosorbent assay (ELISA), radioimmunoassay (RIA) and immunohistochemistry. For example, where the marker sequence is the LacZ gene, the presence of the vector carrying the signal is detected by assays for beta-galactosidase activity. Where the transgene is GFP or luciferase, the vector carrying the signal may be measured visually by color or light production in a luminometer.

15 However, desirably, the transgene is a non-marker sequence encoding a product which is useful in biology and medicine, such as proteins, peptides, RNA, enzymes, or catalytic RNAs. Desirable RNA molecules include tRNA, dsRNA, ribosomal RNA, catalytic RNAs, and antisense RNAs. One example of a useful RNA sequence is a sequence which extinguishes expression of a targeted nucleic acid sequence in the treated animal.

20 The transgene may be used for treatment, e.g., of genetic deficiencies, as a cancer therapeutic or vaccine, for induction of an immune response, and/or for prophylactic vaccine purposes. As used herein, induction of an immune response refers to the ability of a molecule (e.g., a gene product) to induce a T cell and/or a humoral immune response to the molecule. The invention further includes using multiple transgenes, e.g., to correct or
25 ameliorate a condition caused by a multi-subunit protein. In certain situations, a different transgene may be used to encode each subunit of a protein, or to encode different peptides or proteins. This is desirable when the size of the DNA encoding the protein subunit is large, e.g., for an immunoglobulin, the platelet-derived growth factor, or a dystrophin protein. In order for the cell to produce the multi-subunit protein, a cell is infected with the recombinant
30 virus containing each of the different subunits. Alternatively, different subunits of a protein

may be encoded by the same transgene. In this case, a single transgene includes the DNA encoding each of the subunits, with the DNA for each subunit separated by an internal ribozyme entry site (IRES). This is desirable when the size of the DNA encoding each of the subunits is small, e.g., the total size of the DNA encoding the subunits and the IRES is less than five kilobases. As an alternative to an IRES, the DNA may be separated by sequences encoding a 2A peptide, which self-cleaves in a post-translational event. See, e.g., M.L. Donnelly, *et al*, *J. Gen. Virol.*, 78(Pt 1):13-21 (Jan 1997); Furler, S., *et al*, *Gene Ther.*, 8(11):864-873 (June 2001); Klump H., *et al.*, *Gene Ther.*, 8(10):811-817 (May 2001). This 2A peptide is significantly smaller than an IRES, making it well suited for use when space is a limiting factor. However, the selected transgene may encode any biologically active product or other product, e.g., a product desirable for study.

Suitable transgenes may be readily selected by one of skill in the art. The selection of the transgene is not considered to be a limitation of this invention.

2. Regulatory Elements

In addition to the major elements identified above for the minigene, the vector also includes conventional control elements necessary which are operably linked to the transgene in a manner that permits its transcription, translation and/or expression in a cell transfected with the plasmid vector or infected with the virus produced by the invention. As used herein, "operably linked" sequences include both expression control sequences that are contiguous with the gene of interest and expression control sequences that act in *trans* or at a distance to control the gene of interest.

Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation (polyA) signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance secretion of the encoded product. A great number of expression control sequences, including promoters which are native, constitutive, inducible and/or tissue-specific, are known in the art and may be utilized.

Examples of constitutive promoters include, without limitation, the retroviral Rous sarcoma virus (RSV) LTR promoter (optionally with the RSV enhancer), the cytomegalovirus (CMV) promoter (optionally with the CMV enhancer) [see, e.g., Boshart *et al*, *Cell*, **41**:521-530 (1985)], the SV40 promoter, the dihydrofolate reductase promoter, the β -actin promoter, the phosphoglycerol kinase (PGK) promoter, and the EF1 α promoter [Invitrogen].

Inducible promoters allow regulation of gene expression and can be regulated by exogenously supplied compounds, environmental factors such as temperature, or the presence of a specific physiological state, e.g., acute phase, a particular differentiation state of the cell, or in replicating cells only. Inducible promoters and inducible systems are available from a variety of commercial sources, including, without limitation, Invitrogen, Clontech and Ariad. Many other systems have been described and can be readily selected by one of skill in the art. For example, inducible promoters include the zinc-inducible sheep metallothionine (MT) promoter and the dexamethasone (Dex)-inducible mouse mammary tumor virus (MMTV) promoter. Other inducible systems include the T7 polymerase promoter system [WO 98/10088]; the ecdysone insect promoter [No *et al*, *Proc. Natl. Acad. Sci. USA*, **93**:3346-3351 (1996)], the tetracycline-repressible system [Gossen *et al*, *Proc. Natl. Acad. Sci. USA*, **89**:5547-5551 (1992)], the tetracycline-inducible system [Gossen *et al*, *Science*, **268**:1766-1769 (1995), see also Harvey *et al*, *Curr. Opin. Chem. Biol.*, **2**:512-518 (1998)]. Other systems include the FK506 dimer, VP16 or p65 using castradiol, diphenol murislerone, the RU486-inducible system [Wang *et al*, *Nat. Biotech.*, **15**:239-243 (1997) and Wang *et al*, *Gene Ther.*, **4**:432-441 (1997)] and the rapamycin-inducible system [Magari *et al*, *J. Clin. Invest.*, **100**:2865-2872 (1997)]. The effectiveness of some inducible promoters increases over time. In such cases one can enhance the effectiveness of such systems by inserting multiple repressors in tandem, e.g., TetR linked to a TetR by an IRES. Alternatively, one can wait at least 3 days before screening for the desired function. Once can enhance expression of desired proteins by known means to enhance the effectiveness of this system. For example, using the Woodchuck Hepatitis Virus Posttranscriptional Regulatory Element (WPRE).

In another embodiment, the native promoter for the transgene will be used. The native promoter may be preferred when it is desired that expression of the transgene should mimic the native expression. The native promoter may be used when expression of the transgene must be regulated temporally or developmentally, or in a tissue-specific manner, or in response to specific transcriptional stimuli. In a further embodiment, other native expression control elements, such as enhancer elements, polyadenylation sites or Kozak consensus sequences may also be used to mimic the native expression.

Another embodiment of the transgene includes a transgene operably linked to a tissue-specific promoter. For instance, if expression in skeletal muscle is desired, a promoter active in muscle should be used. These include the promoters from genes encoding skeletal β -actin, myosin light chain 2A, dystrophin, muscle creatine kinase, as well as synthetic muscle promoters with activities higher than naturally occurring promoters (see Li *et al.*, *Nat. Biotech.*, 17:241-245 (1999)). Examples of promoters that are tissue-specific are known for liver (albumin, Miyatake *et al.*, *J. Virol.*, 71:5124-32 (1997); hepatitis B virus core promoter, Sandig *et al.*, *Gene Ther.*, 3:1002-9 (1996); alpha-fetoprotein (AFP), Arbuthnot *et al.*, *Hum. Gene Ther.*, 7:1503-14 (1996)), bone osteocalcin (Stein *et al.*, *Mol. Biol. Rep.*, 24:185-96 (1997)); bone sialoprotein (Chen *et al.*, *J. Bone Miner. Res.*, 11:654-64 (1996)), lymphocytes (CD2, Hansal *et al.*, *J. Immunol.*, 161:1063-8 (1998); immunoglobulin heavy chain; T cell receptor chain), neuronal such as neuron-specific enolase (NSE) promoter (Andersen *et al.*, *Cell. Mol. Neurobiol.*, 13:503-15 (1993)), neurofilament light-chain gene (Piccioli *et al.*, *Proc. Natl. Acad. Sci. USA*, 88:5611-5 (1991)), and the neuron-specific vgf gene (Piccioli *et al.*, *Neuron*, 15:373-84 (1995)), among others.

Optionally, vectors carrying transgenes encoding therapeutically useful or immunogenic products may also include selectable markers or reporter genes may include sequences encoding geneticin, hygromycin or purimycin resistance, among others. Such selectable reporters or marker genes (preferably located outside the viral genome to be packaged into a viral particle) can be used to signal the presence of the plasmids in bacterial cells, such as ampicillin resistance. Other components of the vector may include an origin of replication. Selection of these and other promoters and vector elements are conventional and many such sequences are available [see, e.g., Sambrook *et al.*, and references cited therein].

These vectors are generated using the techniques and sequences provided herein, in conjunction with techniques known to those of skill in the art. Such techniques include conventional cloning techniques of cDNA such as those described in texts [Sambrook et al, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, NY], use of overlapping oligonucleotide sequences of the adenovirus genomes, polymerase chain reaction, and any suitable method which provides the desired nucleotide sequence.

III. Production of the Recombinant Viral Particle

In one embodiment, the chimpanzee adenoviral plasmids (or other vectors) are used to produce recombinant adenoviral particles. In one embodiment, the recombinant adenoviruses are functionally deleted in the E1a or E1b genes, and optionally bearing other mutations, e.g., temperature-sensitive mutations or deletions in other genes. In other embodiments, it is desirable to retain an intact E1a and/or E1b region in the recombinant adenoviruses. Such an intact E1 region may be located in its native location in the adenoviral genome or placed in the site of a deletion in the native adenoviral genome (e.g., in the E3 region).

In the construction of useful chimpanzee adenovirus vectors for delivery of a gene to the human (or other mammalian) cell, a range of adenovirus nucleic acid sequences can be employed in the vectors. For example, all or a portion of the adenovirus delayed early gene E3 may be eliminated from the C68 adenovirus sequence which forms a part of the recombinant virus. The function of adenovirus E3 is believed to be irrelevant to the function and production of the recombinant virus particle. Adenovirus vectors may also be constructed having a deletion of at least the ORF6 region of the E4 gene, and more desirably because of the redundancy in the function of this region, the entire E4 region. Still another vector of this invention contains a deletion in the delayed early gene E2a. Deletions may also be made in any of the late genes L1 through L5 of the chimpanzee adenovirus genome. Similarly, deletions in the intermediate genes IX and IVa₂ may be useful for some purposes. Other deletions may be made in the other structural or non-structural adenovirus genes. The above discussed deletions may be used individually, i.e., an adenovirus sequence for use in the present invention may contain deletions in only a single region. Alternatively, deletions

of entire genes or portions thereof effective to destroy their biological activity may be used in any combination. For example, in one exemplary vector, the adenovirus sequence may have deletions of the E1 genes and the E4 gene, or of the E1, E2a and E3 genes, or of the E1 and E3 genes, or of E1, E2a and E4 genes, with or without deletion of E3, and so on. As
5 discussed above, such deletions may be used in combination with other mutations, such as temperature-sensitive mutations, to achieve a desired result.

An adenoviral vector lacking any essential adenoviral sequences (e.g., E1a, E1b, E2a, E2b, E4 ORF6, L1, L2, L3, L4 and L5) may be cultured in the presence of the missing adenoviral gene products which are required for viral infectivity and propagation of an
10 adenoviral particle. These helper functions may be provided by culturing the adenoviral vector in the presence of one or more helper constructs (e.g., a plasmid or virus) or a packaging host cell. See, for example, the techniques described for preparation of a "minimal" human Ad vector in International Patent Application WO96/13597, published May 9, 1996, and incorporated herein by reference.

15 1. Helper Viruses

Thus, depending upon the chimpanzee adenovirus gene content of the viral vectors employed to carry the minigene, a helper adenovirus or non-replicating virus fragment may be necessary to provide sufficient chimpanzee adenovirus gene sequences necessary to produce an infective recombinant viral particle containing the minigene. Useful
20 helper viruses contain selected adenovirus gene sequences not present in the adenovirus vector construct and/or not expressed by the packaging cell line in which the vector is transfected. In one embodiment, the helper virus is replication-defective and contains a variety of adenovirus genes in addition to the sequences described above. Such a helper virus is desirably used in combination with an E1-expressing cell line.

25 Helper viruses may also be formed into poly-cation conjugates as described in Wu *et al*, *J. Biol. Chem.*, 264:16985-16987 (1989); K. J. Fisher and J. M. Wilson, *Biochem. J.*, 299:49 (April 1, 1994). Helper virus may optionally contain a second reporter minigene. A number of such reporter genes are known to the art. The presence of a reporter gene on the helper virus which is different from the transgene on the adenovirus
30 vector allows both the Ad vector and the helper virus to be independently monitored. This

second reporter is used to enable separation between the resulting recombinant virus and the helper virus upon purification.

2. Complementation Cell Lines

To generate recombinant chimpanzee adenoviruses (Ad) deleted in any
5 of the genes described above, the function of the deleted gene region, if essential to the replication and infectivity of the virus, must be supplied to the recombinant virus by a helper virus or cell line, i.e., a complementation or packaging cell line. In many circumstances, a cell line expressing the human E1 can be used to transcomplement the chimp Ad vector. This is particularly advantageous because, due to the diversity between the chimp Ad sequences of
10 the invention and the human AdE1 sequences found in currently available packaging cells, the use of the current human E1-containing cells prevents the generation of replication-competent adenoviruses during the replication and production process. However, in certain circumstances, it will be desirable to utilize a cell line which expresses the E1 gene products can be utilized for production of an E1-deleted chimpanzee adenovirus. Such cell lines have
15 been described. See, e.g., US Patent 6,083,716.

If desired, one may utilize the sequences provided herein to generate a packaging cell or cell line that expresses, at a minimum, the adenovirus E1 gene under the transcriptional control of a promoter for expression in a selected parent cell line. Inducible or constitutive promoters may be employed for this purpose. Examples of such promoters are
20 described in detail elsewhere in this specification. A parent cell is selected for the generation of a novel cell line expressing any desired Ad gene. Without limitation, such a parent cell line may be HeLa [ATCC Accession No. CCL 2], A549 [ATCC Accession No. CCL 185], KB [CCL 17], Detroit [e.g., Detroit 510, CCL 72] and WI-38 [CCL 75] cells, among others. These cell lines are all available from the American Type Culture Collection, 10801
25 University Boulevard, Manassas, Virginia 20110-2209. Other suitable parent cell lines may be obtained from other sources.

Such E1-expressing cell lines are useful in the generation of recombinant chimpanzee adenovirus E1 deleted vectors. Additionally, or alternatively, the invention provides cell lines that express one or more chimpanzee adenoviral gene products,
30 e.g., E1a, E1b, E2a, and/or E4 ORF6, can be constructed using essentially the same

procedures for use in the generation of recombinant chimpanzee viral vectors. Such cell lines can be utilized to transcomplement adenovirus vectors deleted in the essential genes that encode those products, or to provide helper functions necessary for packaging of a helper-dependent virus (e.g., adeno-associated virus). The preparation of a host cell according to this invention involves techniques such as assembly of selected DNA sequences. This assembly may be accomplished utilizing conventional techniques. Such techniques include cDNA and genomic cloning, which are well known and are described in Sambrook et al., cited above, use of overlapping oligonucleotide sequences of the adenovirus genomes, combined with polymerase chain reaction, synthetic methods, and any other suitable methods which provide the desired nucleotide sequence.

In still another alternative, the essential adenoviral gene products are provided in *trans* by the adenoviral vector and/or helper virus. In such an instance, a suitable host cell can be selected from any biological organism, including prokaryotic (e.g., bacterial) cells, and eukaryotic cells, including, insect cells, yeast cells and mammalian cells.

Particularly desirable host cells are selected from among any mammalian species, including, without limitation, cells such as A549, WEHI, 3T3, 10T1/2, 293 cells (which express functional adenoviral E1), Saos, C2C12, L cells, HT1080, HepG2 and primary fibroblast, hepatocyte and myoblast cells derived from mammals including human, monkey, mouse, rat, rabbit, and hamster. The selection of the mammalian species providing the cells is not a limitation of this invention; nor is the type of mammalian cell, i.e., fibroblast, hepatocyte, tumor cell, etc.

3. Assembly of Viral Particle and Transfection of a Cell Line

Generally, when delivering the vector comprising the minigene by transfection, the vector is delivered in an amount from about 5 μ g to about 100 μ g DNA, and preferably about 10 to about 50 μ g DNA to about 1×10^4 cells to about 1×10^{13} cells, and preferably about 10^5 cells. However, the relative amounts of vector DNA to host cells may be adjusted, taking into consideration such factors as the selected vector, the delivery method and the host cells selected.

The vector may be any vector known in the art or disclosed above, including naked DNA, a plasmid, phage, transposon, cosmids, viruses, etc. Introduction into

the host cell of the vector may be achieved by any means known in the art or as disclosed above, including transfection, and infection. One or more of the adenoviral genes may be stably integrated into the genome of the host cell, stably expressed as episomes, or expressed transiently. The gene products may all be expressed transiently, on an episome or stably integrated, or some of the gene products may be expressed stably while others are expressed transiently. Furthermore, the promoters for each of the adenoviral genes may be selected independently from a constitutive promoter, an inducible promoter or a native adenoviral promoter. The promoters may be regulated by a specific physiological state of the organism or cell (i.e., by the differentiation state or in replicating or quiescent cells) or by exogenously-added factors, for example.

Introduction of the molecules (as plasmids or viruses) into the host cell may also be accomplished using techniques known to the skilled artisan and as discussed throughout the specification. In preferred embodiment, standard transfection techniques are used, e.g., CaPO₄ transfection or electroporation.

Assembly of the selected DNA sequences of the adenovirus (as well as the transgene and other vector elements into various intermediate plasmids, and the use of the plasmids and vectors to produce a recombinant viral particle are all achieved using conventional techniques. Such techniques include conventional cloning techniques of cDNA such as those described in texts [Sambrook et al, cited above], use of overlapping oligonucleotide sequences of the adenovirus genomes, polymerase chain reaction, and any suitable method which provides the desired nucleotide sequence. Standard transfection and co-transfection techniques are employed, e.g., CaPO₄ precipitation techniques. Other conventional methods employed include homologous recombination of the viral genomes, plaquing of viruses in agar overlay, methods of measuring signal generation, and the like.

For example, following the construction and assembly of the desired minigene-containing viral vector, the vector is transfected *in vitro* in the presence of a helper virus into the packaging cell line. Homologous recombination occurs between the helper and the vector sequences, which permits the adenovirus-transgene sequences in the vector to be replicated and packaged into virion capsids, resulting in the recombinant viral vector

particles. The current method for producing such virus particles is transfection-based. However, the invention is not limited to such methods.

The resulting recombinant chimpanzee adenoviruses are useful in transferring a selected transgene to a selected cell. In *in vivo* experiments with the recombinant virus grown in the packaging cell lines, the E1-deleted recombinant chimpanzee adenoviral vectors of the invention demonstrate utility in transferring a transgene to a non-chimpanzee, preferably a human, cell.

IV. Use of Non-Viral C68 Proteins and C68-derived Adenoviruses

The recombinant adenovirus vectors of the invention are useful for gene transfer to a human or non-chimpanzee veterinary patient *in vitro*, *ex vivo*, and *in vivo*. In addition, a variety of C68 proteins described herein are useful in non-viral targeting of transgenes, proteins, chemical molecules, and other moieties or molecules to cells. Suitable methods of delivery and dosing regimens are readily determined based upon the targeted molecule and targeting protein. Examples of suitable genes and sources of proteins for protein-mediated delivery are provided in the sections below relating to viral delivery of therapeutic and immunogenic molecules. While the discussion below focuses on viral vectors, it will be appreciated that the C68-derived proteins of the invention may be formulated as described herein for the C68-derived viral vectors and the same routes of administration and regimens may be utilized.

The recombinant adenovirus vectors described herein can be used as expression vectors for the production of the products encoded by the heterologous genes *in vitro*. For example, the recombinant adenoviruses containing a gene inserted into the location of an E1 deletion may be transfected into an E1-expressing cell line as described above. Alternatively, replication-competent adenoviruses may be used in another selected cell line. The transfected cells are then cultured in the conventional manner, allowing the recombinant adenovirus to express the gene product from the promoter. The gene product may then be recovered from the culture medium by known conventional methods of protein isolation and recovery from culture.

A C68-derived vector or C68-derived protein of the invention provides an efficient gene transfer vehicle that can deliver a selected transgene or other molecule to a selected host cell *in vivo* or *ex vivo* even where the organism has neutralizing antibodies to one or more AAV serotypes. In one embodiment, the rAAV and the cells are mixed *ex vivo*; the infected cells are cultured using conventional methodologies; and the transduced cells are re-infused into the patient. These compositions are particularly well suited to gene delivery for therapeutic purposes and for immunization, including inducing protective immunity.

More commonly, the C68-derived vectors and C68-derived proteins of the invention will be utilized for delivery of therapeutic or immunogenic molecules, as described below. It will be readily understood for both applications, that the C68-derived constructs of the invention are useful for use in regimens involving single administrations, as well as in regimens involving repeat delivery of adenoviral vectors or non-viral targeted delivery, or repeat delivery of the transgene or other molecule to the cells.

Such regimens typically involve delivery of a series of viral vectors in which the viral capsids are alternated. The viral capsids may be changed for each subsequent administration, or after a pre-selected number of administrations of a particular serotype capsid (e.g., one, two, three, four or more). For example, a regimen may involve delivery of a rAd with a C68-derived capsid and delivery with a rAd with another human or non-human primate adenovirus serotype. Optionally, these regimens may involve administration of rAd with capsids of other non-human primate adenoviruses, human adenoviruses, or artificial serotypes such as are described herein. Alternatively, the regimens involve administration of C68-derived proteins for non-viral targeting with repeat administrations of C68-derived proteins, or with other protein-based delivery systems. Each phase of these regimens can involve administration of a series of injections (or other delivery routes) with a single C68-derived construct followed by a series with another Ad serotype construct. Alternatively, the C68-derived vectors and proteins of the invention may be utilized in regimens involving other non-adenoviral-mediated delivery systems, including other viral systems, non-viral delivery systems, protein, peptides, and other biologically active molecules.

The following sections will focus on exemplary molecules which may be delivered via the adenoviral vectors of the invention.

A. Ad-Mediated Delivery of Therapeutic Molecules

In one embodiment, the above-described C68-derived constructs are administered to humans according to published methods for gene therapy. A C68-derived construct bearing a transgene can be administered to a patient, preferably suspended in a biologically compatible solution or pharmaceutically acceptable delivery vehicle. A suitable vehicle includes sterile saline. Other aqueous and non-aqueous isotonic sterile injection solutions and aqueous and non-aqueous sterile suspensions known to be pharmaceutically acceptable carriers and well known to those of skill in the art may be employed for this purpose.

The C68-derived adenoviral vectors are administered in sufficient amounts to transduce the target cells and to provide sufficient levels of gene transfer and expression to provide a therapeutic benefit without undue adverse or with medically acceptable physiological effects, which can be determined by those skilled in the medical arts. Conventional and pharmaceutically acceptable routes of administration include, but are not limited to, direct delivery to the retina and other intraocular delivery methods, direct delivery to the liver, intranasal, intravenous, intramuscular, intratracheal, subcutaneous, intradermal, rectal, oral and other parenteral routes of administration. Routes of administration may be combined, if desired, or adjusted depending upon the transgene or the condition. The route of administration primarily will depend on the nature of the condition being treated.

Dosages of the viral vector will depend primarily on factors such as the condition being treated, the age, weight and health of the patient, and may thus vary among patients. For example, a therapeutically effective adult human or veterinary dosage of the viral vector is generally in the range of from about 100 μ L to about 100 mL of a carrier containing concentrations of from about 1×10^6 to about 1×10^{15} particles, about 1×10^{11} to 1×10^{13} particles, or about 1×10^9 to 1×10^{12} particles virus. Dosages will range depending upon the size of the animal and the route of administration. For example, a suitable human or veterinary dosage (for about an 80 kg animal) for intramuscular injection is in the range of about 1×10^9 to about 5×10^{12} particles per mL, for a single site. Optionally, multiple sites of administration may be delivered. In another example, a suitable human or veterinary dosage may be in the range of about 1×10^{11} to about 1×10^{15} particles for an oral formulation.

When C68 proteins of the invention are utilized for targeted delivery, suitable dosage ranges, a therapeutically effective adult human or veterinary dosage of the construct is generally in the range of from about 100 μ L to about 100 mL of a carrier containing concentrations of from about 0.01 μ g to about 100 mg protein, about 0.1 μ g to about 10 mg, about 1 μ g to about 1 mg protein. Dosages will range depending upon the size of the animal and the route of administration. Routes of administration may be readily selected from any suitable route including, without limitation, the routes described above.

One of skill in the art may adjust these doses, depending the route of administration, and the therapeutic or vaccinal application for which the C68-derived construct is employed. The levels of expression of the transgene, or for an immunogen, the level of circulating antibody, can be monitored to determine the frequency of dosage administration. Yet other methods for determining the timing of frequency of administration will be readily apparent to one of skill in the art.

An optional method step involves the co-administration to the patient, either concurrently with, or before or after administration of the C68-derived construct, of a suitable amount of a short acting immune modulator. The selected immune modulator is defined herein as an agent capable of inhibiting the formation of neutralizing antibodies directed against the recombinant vector of this invention or capable of inhibiting cytolytic T lymphocyte (CTL) elimination of the vector. The immune modulator may interfere with the interactions between the T helper subsets (T_{H1} or T_{H2}) and B cells to inhibit neutralizing antibody formation. Alternatively, the immune modulator may inhibit the interaction between T_{H1} cells and CTLs to reduce the occurrence of CTL elimination of the vector. A variety of useful immune modulators and dosages for use of same are disclosed, for example, in Yang *et al.*, *J. Virol.*, 70(9) (Sept., 1996); International Patent Application No. WO96/12406, published May 2, 1996; and International Patent Application No. PCT/US96/03035, all incorporated herein by reference.

1. Therapeutic Transgenes

Useful therapeutic products encoded by the transgene include hormones and growth and differentiation factors including, without limitation, insulin, glucagon, growth hormone (GH), parathyroid hormone (PTH), growth hormone releasing

factor (GRF), follicle stimulating hormone (FSH), luteinizing hormone (LH), human chorionic gonadotropin (hCG), vascular endothelial growth factor (VEGF), angiopoietins, angiostatin, granulocyte colony stimulating factor (GCSF), erythropoietin (EPO), connective tissue growth factor (CTGF), basic fibroblast growth factor (bFGF), acidic fibroblast growth factor (aFGF), epidermal growth factor (EGF), transforming growth factor (TGF), platelet-derived growth factor (PDGF), insulin growth factors I and II (IGF-I and IGF-II), any one of the transforming growth factor superfamily, including TGF, activins, inhibins, or any of the bone morphogenic proteins (BMP) BMPs 1-15, any one of the heregluin/neuregulin/ARIA/neu differentiation factor (NDF) family of growth factors, nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), neurotrophins NT-3 and NT-4/5, ciliary neurotrophic factor (CNTF), glial cell line derived neurotrophic factor (GDNF), neurturin, agrin, any one of the family of semaphorins/collapsins, netrin-1 and netrin-2, hepatocyte growth factor (HGF), ephrins, noggin, sonic hedgehog and tyrosine hydroxylase.

Other useful transgene products include proteins that regulate the immune system including, without limitation, cytokines and lymphokines such as thrombopoietin (TPO), interleukins (IL) IL-1 through IL-18, monocyte chemoattractant protein, leukemia inhibitory factor, granulocyte-macrophage colony stimulating factor, Fas ligand, tumor necrosis factors and, interferons, and, stem cell factor, flk-2/flt3 ligand. Gene products produced by the immune system are also useful in the invention. These include, without limitations, immunoglobulins IgG, IgM, IgA, IgD and IgE, chimeric immunoglobulins, humanized antibodies, single chain antibodies, T cell receptors, chimeric T cell receptors, single chain T cell receptors, class I and class II MHC molecules, as well as engineered immunoglobulins and MHC molecules. Useful gene products also include complement regulatory proteins such as complement regulatory proteins, membrane cofactor protein (MCP), decay accelerating factor (DAF), CR1, CF2 and CD59.

Still other useful gene products include any one of the receptors for the hormones, growth factors, cytokines, lymphokines, regulatory proteins and immune system proteins. The invention encompasses receptors for cholesterol regulation, including the low density lipoprotein (LDL) receptor, high density lipoprotein (HDL) receptor, the very low

density lipoprotein (VLDL) receptor, and the scavenger receptor. The invention also encompasses gene products such as members of the steroid hormone receptor superfamily including glucocorticoid receptors and estrogen receptors, Vitamin D receptors and other nuclear receptors. In addition, useful gene products include transcription factors such as *jun*,
5 *fos*, max, mad, serum response factor (SRF), AP-1, AP2, *myb*, MyoD and myogenin, ETS-box containing proteins, TFE3, E2F, ATF1, ATF2, ATF3, ATF4, ZF5, NFAT, CREB, HNF-4, C/EBP, SP1, CCAAT-box binding proteins, interferon regulation factor (IRF-1), Wilms tumor protein, ETS-binding protein, STAT, GATA-box binding proteins, e.g., GATA-3, and the forkhead family of winged helix proteins.

10 Other useful gene products include, carbamoyl synthetase I, ornithine transcarbamylase, arginosuccinate synthetase, arginosuccinate lyase, arginase, fumarylacetate hydrolase, phenylalanine hydroxylase, alpha-1 antitrypsin, glucose-6-phosphatase, porphobilinogen deaminase, factor VIII, factor IX, cystathione beta-synthase, branched chain ketoacid decarboxylase, albumin, isovaleryl-coA dehydrogenase, propionyl
15 CoA carboxylase, methyl malonyl CoA mutase, glutaryl CoA dehydrogenase, insulin, beta-glucosidase, pyruvate carboxylate, hepatic phosphorylase, phosphorylase kinase, glycine decarboxylase, H-protein, T-protein, a cystic fibrosis transmembrane regulator (CFTR) sequence, and a dystrophin cDNA sequence.

Other useful gene products include non-naturally occurring
20 polypeptides, such as chimeric or hybrid polypeptides having a non-naturally occurring amino acid sequence containing insertions, deletions or amino acid substitutions. For example, single-chain engineered immunoglobulins could be useful in certain immunocompromised patients. Other types of non-naturally occurring gene sequences include antisense molecules and catalytic nucleic acids, such as ribozymes, which could be used to reduce overexpression
25 of a target.

Reduction and/or modulation of expression of a gene are particularly desirable for treatment of hyperproliferative conditions characterized by hyperproliferating cells, as are cancers and psoriasis. Target polypeptides include those polypeptides which are produced exclusively or at higher levels in hyperproliferative cells as compared to normal
30 cells. Target antigens include polypeptides encoded by oncogenes such as *myb*, *myc*, *fyn*, and

the translocation gene bcr/abl, ras, src, P53, neu, trk and EGRF. In addition to oncogene products as target antigens, target polypeptides for anti-cancer treatments and protective regimens include variable regions of antibodies made by B cell lymphomas and variable regions of T cell receptors of T cell lymphomas which, in some embodiments, are also used
5 as target antigens for autoimmune disease. Other tumor-associated polypeptides can be used as target polypeptides such as polypeptides which are found at higher levels in tumor cells including the polypeptide recognized by monoclonal antibody 17-1A and folate binding polypeptides. Such target polypeptides and their ligands are also useful in forming fusion partners with a C68 protein of the invention.

10 Other suitable therapeutic polypeptides and proteins include those which may be useful for treating individuals suffering from autoimmune diseases and disorders by conferring a broad based protective immune response against targets that are associated with autoimmunity including cell receptors and cells which produce self-directed antibodies. T cell mediated autoimmune diseases include Rheumatoid arthritis (RA),
15 multiple sclerosis (MS), Sjögren's syndrome, sarcoidosis, insulin dependent diabetes mellitus (IDDM), autoimmune thyroiditis, reactive arthritis, ankylosing spondylitis, scleroderma, polymyositis, dermatomyositis, psoriasis, vasculitis, Wegener's granulomatosis, Crohn's disease and ulcerative colitis. Each of these diseases is characterized by T cell receptors (TCRs) that bind to endogenous antigens and initiate the inflammatory cascade associated
20 with autoimmune diseases.

The C68-derived constructs of the invention are particularly well suited for therapeutic regimens in which multiple deliveries of transgenes is desired, e.g., in regimens involving redelivery of the same transgene or in combination regimens involving delivery of other transgenes. Such regimens may involve administration of a C68-derived
25 construct, followed by re-administration with a vector from the same serotype adenovirus. Particularly desirable regimens involve administration of a C68-derived construct of the invention, in which the serotype of the viral vector delivered in the first administration differs from the serotype of the viral vector utilized in one or more of the subsequent administrations. For example, a therapeutic regimen involves administration of a C68-
30 derived vector and repeat administration with one or more adenoviral vectors of the same or

different serotypes. In another example, a therapeutic regimen involves administration of an adenoviral vector followed by repeat administration with a C68-derived vector of the invention which differs from the serotype of the first delivered adenoviral vector, and optionally further administration with another vector which is the same or, preferably, differs from the serotype of the vector in the prior administration steps. These regimens are not limited to delivery of adenoviral vectors constructed using the C68-derived capsids of the invention. Rather, these regimens can readily utilize constructs, including non-viral targeting proteins and viral vectors, from other adenoviral serotypes, including, without limitation, other chimpanzee adenoviral serotypes (e.g., C1, etc), other non-human primate adenoviral serotypes, or human adenoviral serotypes, in combination with one or more of the C68-derived constructs of the invention. Examples of such chimpanzee, other non-human primate and human adenoviral serotypes are discussed elsewhere in this document. Further, these therapeutic regimens may involve either simultaneous or sequential delivery of C68-derived constructs of the invention in combination with non-adenoviral vectors, non-viral vectors, and/or a variety of other therapeutically useful compounds or molecules. The present invention is not limited to these therapeutic regimens, a variety of which will be readily apparent to one of skill in the art.

B. Ad-Mediated Delivery of Immunogenic Transgenes

The C68-derived constructs of the invention, including viral vectors and proteins, may also be employed as immunogenic compositions. As used herein, an immunogenic composition is a composition to which a humoral (e.g., antibody) or cellular (e.g., a cytotoxic T cell) response is mounted to a transgene product delivered by the immunogenic composition following delivery to a mammal, and preferably a primate. The present invention provides a recombinant C68-derived Ad that can contain in any of its adenovirus sequence deletions a gene encoding a desired immunogen, or a C68 protein capable of targeting an immunogenic molecule. The C68-derived adenovirus is well suited for use as a live recombinant virus vaccine in different animal species compared to an adenovirus of human origin, but is not limited to such a use. The recombinant adenoviruses and C68 proteins can be used as prophylactic or therapeutic vaccines against any pathogen for

which the antigen(s) crucial for induction of an immune response and able to limit the spread of the pathogen has been identified and for which the cDNA is available.

Such vaccinal (or other immunogenic) compositions are formulated in a suitable delivery vehicle, as described above. Generally, doses for the immunogenic compositions are in the range defined above for therapeutic compositions. The levels of immunity of the selected gene can be monitored to determine the need, if any, for boosters. Following an assessment of antibody titers in the serum, optional booster immunizations may be desired.

Optionally, a vaccinal composition of the invention may be formulated to contain other components, including, e.g. adjuvants, stabilizers, pH adjusters, preservatives and the like. Such components are well known to those of skill in the vaccine art. Examples of suitable adjuvants include, without limitation, liposomes, alum, monophosphoryl lipid A, and any biologically active factor, such as cytokine, an interleukin, a chemokine, a ligands, and optimally combinations thereof. Certain of these biologically active factors can be expressed *in vivo*, e.g., via a polynucleotide, plasmid or viral vector. For example, such an adjuvant can be administered with a priming DNA vaccine encoding an antigen to enhance the antigen-specific immune response compared with the immune response generated upon priming with a DNA vaccine encoding the antigen only.

The recombinant adenoviruses are administered in a "an immunogenic amount", that is, an amount of recombinant adenovirus that is effective in a route of administration to transfect the desired cells and provide sufficient levels of expression of the selected gene to induce an immune response. Where protective immunity is provided, the recombinant adenoviruses are considered to be vaccine compositions useful in preventing infection and/or recurrent disease.

Alternatively, or in addition, the vectors of the invention may contain, or C68-capsid or other protein can be utilized to target a transgene encoding a peptide, polypeptide or protein which induces an immune response to a selected immunogen. The C68-derived viruses of this invention are expected to be highly efficacious at inducing cytolytic T cells and antibodies to the inserted heterologous antigenic protein expressed by the vector.

1. Immunogenic Transgenes

For example, immunogens may be selected from a variety of viral families. Example of desirable viral families against which an immune response would be desirable include, the picornavirus family, which includes the genera rhinoviruses, which are responsible for about 50% of cases of the common cold; the genera enteroviruses, which include polioviruses, coxsackieviruses, echoviruses, and human enteroviruses such as hepatitis A virus; and the genera aphthoviruses, which are responsible for foot and mouth diseases, primarily in non-human animals. Within the picornavirus family of viruses, target antigens include the VP1, VP2, VP3, VP4, and VPG. Another viral family includes the calcivirus family, which encompasses the Norwalk group of viruses, which are an important causative agent of epidemic gastroenteritis. Still another viral family desirable for use in targeting antigens for inducing immune responses in humans and non-human animals is the togavirus family, which includes the genera alphavirus, which include Sindbis viruses, Ross River virus, and Venezuelan, Eastern & Western Equine encephalitis, and rubivirus, including Rubella virus. The flaviviridae family includes dengue, yellow fever, Japanese encephalitis, St. Louis encephalitis and tick borne encephalitis viruses. Other target antigens may be generated from the Hepatitis C or the coronavirus family, which includes a number of non-human viruses such as infectious bronchitis virus (poultry), porcine transmissible gastroenteric virus (pig), porcine hemagglutinating encephalomyelitis virus (pig), feline infectious peritonitis virus (cats), feline enteric coronavirus (cat), canine coronavirus (dog), and human respiratory coronaviruses, which may cause the common cold and/or non-A, B or C hepatitis. Within the coronavirus family, target antigens include the E1 (also called M or matrix protein), E2 (also called S or Spike protein), E3 (also called HE or hemagglutinin-esterase) glycoprotein (not present in all coronaviruses), or N (nucleocapsid). Still other antigens may be targeted against the rhabdovirus family, which includes the genera vesiculovirus (e.g., Vesicular Stomatitis Virus), and the general lyssavirus (e.g., rabies). Within the rhabdovirus family, suitable antigens may be derived from the G protein or the N protein. The family filoviridae, which includes hemorrhagic fever viruses such as Marburg and Ebola virus may be a suitable source of antigens. The paramyxovirus family includes parainfluenza Virus Type 1, parainfluenza Virus Type 3, bovine parainfluenza Virus Type 3,

rubulavirus (mumps virus), parainfluenza Virus Type 2, parainfluenza virus Type 4, Newcastle disease virus (chickens), rinderpest, morbillivirus, which includes measles and canine distemper, and pneumovirus, which includes respiratory syncytial virus. The influenza virus is classified within the family orthomyxovirus and is a suitable source of antigen (e.g., the HA protein, the N1 protein). The bunyavirus family includes the genera bunyavirus (California encephalitis, La Crosse), phlebovirus (Rift Valley Fever), hantavirus (puremala is a hemahagin fever virus), nairovirus (Nairobi sheep disease) and various unassigned bungaviruses. The arenavirus family provides a source of antigens against LCM and Lassa fever virus. The reovirus family includes the genera reovirus, rotavirus (which causes acute gastroenteritis in children), orbiviruses, and cultivirus (Colorado Tick fever, Lebombo (humans), equine encephalosis, blue tongue).

The retrovirus family includes the sub-family oncorivirinal which encompasses such human and veterinary diseases as feline leukemia virus, HTLVI and HTLVII, lentivirinal (which includes human immunodeficiency virus (HIV), simian immunodeficiency virus (SIV), feline immunodeficiency virus (FIV), equine infectious anemia virus, and spumavirinal). Among the lentiviruses, many suitable antigens have been described and can readily be selected. Examples of suitable HIV and SIV antigens include, without limitation the gag, pol, Vif, Vpx, VPR, Env, Tat, Nef, and Rev proteins, as well as various fragments thereof. For example, suitable fragments of the Env protein may include any of its subunits such as the gp120, gp160, gp41, or smaller fragments thereof, e.g., of at least about 8 amino acids in length. Similarly, fragments of the tat protein may be selected. [See, US Patent 5,891,994 and US Patent 6,193,981.] See, also, the HIV and SIV proteins described in D.H. Barouch et al, J. Virol., 75(5):2462-2467 (March 2001), and R.R. Amara, et al, *Science*, 292:69-74 (6 April 2001). In another example, the HIV and/or SIV immunogenic proteins or peptides may be used to form fusion proteins or other immunogenic molecules. See, e.g., the HIV-1 Tat and/or Nef fusion proteins and immunization regimens described in WO 01/54719, published August 2, 2001, and WO 99/16884, published April 8, 1999. The invention is not limited to the HIV and/or SIV immunogenic proteins or peptides described herein. In addition, a variety of modifications to these proteins have been described or could readily be made by one of skill in the art. See, e.g., the modified gag protein that is described

in US Patent 5,972,596. Further, any desired HIV and/or SIV immunogens may be delivered alone or in combination. Such combinations may include expression from a single vector or from multiple vectors. Optionally, another combination may involve delivery of one or more expressed immunogens with delivery of one or more of the immunogens in protein form.

5 Such combinations are discussed in more detail below.

The papovavirus family includes the sub-family polyomaviruses (BKU and JCU viruses) and the sub-family papillomavirus (associated with cancers or malignant progression of papilloma). The adenovirus family includes viruses (EX, AD7, ARD, O.B.) which cause respiratory disease and/or enteritis. The parvovirus family feline parvovirus (feline enteritis), feline panleucopeniavirus, canine parvovirus, and porcine parvovirus. The herpesvirus family includes the sub-family alphaherpesvirinae, which encompasses the genera simplexvirus (HSV1, HSV2), varicellovirus (pseudorabies, varicella zoster) and the sub-family betaherpesvirinae, which includes the genera cytomegalovirus (HCMV, muromegalovirus) and the sub-family gammaherpesvirinae, which includes the genera lymphocryptovirus, EBV (Burkitts lymphoma), infectious rhinotracheitis, Marek's disease virus, and rhadinovirus. The poxvirus family includes the sub-family chordopoxvirinae, which encompasses the genera orthopoxvirus (Variola (Smallpox) and Vaccinia (Cowpox)), parapoxvirus, avipoxvirus, capripoxvirus, leporipoxvirus, suipoxvirus, and the sub-family entomopoxvirinae. The hepadnavirus family includes the Hepatitis B virus. One unclassified virus which may be suitable source of antigens is the Hepatitis delta virus. Still other viral sources may include avian infectious bursal disease virus and porcine respiratory and reproductive syndrome virus. The alphavirus family includes equine arteritis virus and various Encephalitis viruses.

10
15
20

The present invention may also encompass immunogens which are useful to immunize a human or non-human animal against other pathogens including bacteria, fungi, parasitic microorganisms or multicellular parasites which infect human and non-human vertebrates, or from a cancer cell or tumor cell. Examples of bacterial pathogens include pathogenic gram-positive cocci include pneumococci; staphylococci; and streptococci. Pathogenic gram-negative cocci include meningococcus; gonococcus. Pathogenic enteric gram-negative bacilli include enterobacteriaceae; pseudomonas, acinetobacteria and

25
30

eikenella; melioidosis; salmonella; shigella; haemophilus; moraxella; *H. ducreyi* (which causes chancroid); brucella; *Francisella tularensis* (which causes tularemia); yersinia (pasteurella); streptobacillus moniliformis and spirillum; Gram-positive bacilli include listeria monocytogenes; erysipelotheix rhusiopathiae; *Corynebacterium diphtheria* (diphtheria);
 5 cholera; *B. anthracis* (anthrax); donovanosis (granuloma inguinale); and bartonellosis.

Diseases caused by pathogenic anaerobic bacteria include tetanus; botulism; other clostridia; tuberculosis; leprosy; and other mycobacteria. Pathogenic spirochetal diseases include syphilis; treponematoses: yaws, pinta and endemic syphilis; and leptospirosis. Other infections caused by higher pathogen bacteria and pathogenic fungi include actinomycosis;

10 nocardiosis; cryptococcosis, blastomycosis, histoplasmosis and coccidioidomycosis; candidiasis, aspergillosis, and mucormycosis; sporotrichosis; paracoccidioidomycosis, petriellidiosis, torulopsosis, mycetoma and chromomycosis; and dermatophytosis. Rickettsial infections include Typhus fever, Rocky Mountain spotted fever, Q fever, and Rickettsialpox.

Examples of mycoplasma and chlamydial infections include: mycoplasma pneumoniae;

15 lymphogranuloma venereum; psittacosis; and perinatal chlamydial infections. Pathogenic eukaryotes encompass pathogenic protozoans and helminths and infections produced thereby include: amebiasis; malaria; leishmaniasis; trypanosomiasis; toxoplasmosis; *Pneumocystis carinii*; *Trichans*; *Toxoplasma gondii*; babesiosis; giardiasis; trichinosis; filariasis; schistosomiasis; nematodes; trematodes or flukes; and cestode (tapeworm) infections.

20 Many of these organisms and/or toxins produced thereby have been identified by the Centers for Disease Control [(CDC), Department of Health and Human Services, USA], as agents which have potential for use in biological attacks. For example, some of these biological agents, include, *Bacillus anthracis* (anthrax), *Clostridium botulinum* and its toxin (botulism), *Yersinia pestis* (plague), variola major (smallpox), *Francisella tularensis* (tularemia), and viral hemorrhagic fever, all of which are currently classified as
 25 Category A agents; *Coxiella burnetti* (Q fever); Brucella species (brucellosis), *Burkholderia mallei* (glanders), *Ricinus communis* and its toxin (ricin toxin), *Clostridium perfringens* and its toxin (epsilon toxin), *Staphylococcus* species and their toxins (enterotoxin B), all of which are currently classified as Category B agents; and Nippan virus, multidrug-resistant
 30 tuberculosis, yellow fever, tickborne hemorrhagic fever viruses, tickborne encephalitis

viruses, and hantaviruses, which are currently classified as Category C agents. In addition, other organisms, which are so classified or differently classified, may be identified and/or used for such a purpose in the future. It will be readily understood that the viral vectors and other constructs described herein are useful to deliver antigens from these organisms, viruses, their toxins or other by-products, which will prevent and/or treat infection or other adverse reactions with these biological agents.

Administration of the vectors and proteins of the invention to deliver immunogens against the variable region of the T cells elicit an immune response including CTLs to eliminate those T cells. In RA, several specific variable regions of TCRs which are involved in the disease have been characterized. These TCRs include V-3, V-14, V-17 and V α -17. Thus, delivery of a nucleic acid sequence that encodes at least one of these polypeptides will elicit an immune response that will target T cells involved in RA. In MS, several specific variable regions of TCRs which are involved in the disease have been characterized. These TCRs include V-7 and V α -10. Thus, delivery of a nucleic acid sequence that encodes at least one of these polypeptides will elicit an immune response that will target T cells involved in MS. In scleroderma, several specific variable regions of TCRs which are involved in the disease have been characterized. These TCRs include V-6, V-8, V-14 and V α -16, V α -3C, V α -7, V α -14, V α -15, V α -16, V α -28 and V α -12. Thus, delivery of a recombinant chimpanzee adenovirus that encodes at least one of these polypeptides will elicit an immune response that will target T cells involved in scleroderma.

C. Ad-Mediated Delivery Methods

The therapeutic levels, or levels of immunity, of the selected gene can be monitored to determine the need, if any, for boosters. Following an assessment of CD8+ T cell response, or optionally, antibody titers, in the serum, optional booster immunizations may be desired. Optionally, the C68-derived constructs of the invention may be delivered in a single administration or in various combination regimens, e.g., in combination with a regimen or course of treatment involving other active ingredients or in a prime-boost regimen. A variety of such regimens have been described in the art and may be readily selected.

For example, prime-boost regimens may involve the administration of a DNA (e.g., plasmid) based vector to prime the immune system to second, booster, administration

with a traditional antigen, such as a protein or a recombinant virus carrying the sequences encoding such an antigen. See, e.g., WO 00/11140, published March 2, 2000, incorporated by reference. Alternatively, an immunization regimen may involve the administration of a recombinant chimpanzee adenoviral vector of the invention to boost the immune response to
5 a vector (either viral or DNA-based) carrying an antigen, or a protein. In still another alternative, an immunization regimen involves administration of a protein followed by booster with a vector encoding the antigen.

In one embodiment, the invention provides a method of priming and boosting an immune response to a selected antigen by delivering a plasmid DNA vector carrying said
10 antigen, followed by boosting with a recombinant chimpanzee adenoviral vector of the invention. In one embodiment, the prime-boost regimen involves the expression of multiproteins from the prime and/or the boost vehicle. See, e.g., R.R. Amara, *Science*, **292**:69-74 (6 April 2001) which describes a multiprotein regimen for expression of protein subunits useful for generating an immune response against HIV and SIV. For example, a
15 DNA prime may deliver the Gag, Pol, Vif, VPX and Vpr and Env, Tat, and Rev from a single transcript. Alternatively, the SIV Gag, Pol and HIV-1 Env is delivered in a recombinant adenovirus construct of the invention. Still other regimens are described in WO 99/16884 and WO 01/54719.

However, the prime-boost regimens are not limited to immunization for HIV
20 or to delivery of these antigens. For example, priming may involve delivering with a first chimp vector of the invention followed by boosting with a second chimp vector, or with a composition containing the antigen itself in protein form. In one example, the prime-boost regimen can provide a protective immune response to the virus, bacteria or other organism from which the antigen is derived. In another desired embodiment, the prime-boost regimen
25 provides a therapeutic effect that can be measured using convention assays for detection of the presence of the condition for which therapy is being administered.

The priming composition may be administered at various sites in the body in a dose dependent manner, which depends on the antigen to which the desired immune response is being targeted. The invention is not limited to the amount or situs of injection(s) or to the
30 pharmaceutical carrier. Rather, the regimen may involve a priming and/or boosting step,

each of which may include a single dose or dosage that is administered hourly, daily, weekly or monthly, or yearly. As an example, the mammals may receive one or two doses containing between about 10 μg to about 50 μg of plasmid in carrier. A desirable amount of a DNA composition ranges between about 1 μg to about 10,000 μg of the DNA vector. Dosages may vary from about 1 μg to 1000 μg DNA per kg of subject body weight. The amount or site of delivery is desirably selected based upon the identity and condition of the mammal.

The dosage unit of the vector suitable for delivery of the antigen to the mammal is described herein. The vector is prepared for administration by being suspended or dissolved in a pharmaceutically or physiologically acceptable carrier such as isotonic saline; isotonic salts solution or other formulations that will be apparent to those skilled in such administration. The appropriate carrier will be evident to those skilled in the art and will depend in large part upon the route of administration. The compositions of the invention may be administered to a mammal according to the routes described above, in a sustained release formulation using a biodegradable biocompatible polymer, or by on-site delivery using micelles, gels and liposomes. Optionally, the priming step of this invention also includes administering with the priming composition, a suitable amount of an adjuvant, such as are defined herein.

Preferably, a boosting composition is administered about 2 to about 27 weeks after administering the priming composition to the mammalian subject. The administration of the boosting composition is accomplished using an effective amount of a boosting composition containing or capable of delivering the same antigen as administered by the priming DNA vaccine. The boosting composition may be composed of a recombinant viral vector derived from the same viral source (e.g., adenoviral sequences of the invention) or from another source. Alternatively, the "boosting composition" can be a composition containing the same antigen as encoded in the priming DNA vaccine, but in the form of a protein or peptide, which composition induces an immune response in the host. In another embodiment, the boosting composition contains a DNA sequence encoding the antigen under the control of a regulatory sequence directing its expression in a mammalian cell, e.g., vectors such as well-known bacterial or viral vectors. The primary requirements of the boosting

composition are that the antigen of the composition is the same antigen, or a cross-reactive antigen, as that encoded by the priming composition.

In another embodiment, the chimpanzee adenoviral vectors and C68 targeting proteins of the invention are also well suited for use in a variety of other immunization and therapeutic regimens. Such regimens may involve delivery of C68 constructs of the invention simultaneously or sequentially with Ad constructs of different serotype capsids, regimens in which C68-derived constructs of the invention are delivered simultaneously or sequentially with non-Ad vectors, regimens in which the adenoviral vectors of the invention are delivered simultaneously or sequentially with proteins, peptides, and/or other biologically useful therapeutic or immunogenic compounds. Such uses will be readily apparent to one of skill in the art.

V. Method for Rapid Screening of Bacterial Transformants

An elegant selection method is provided by the present invention, which permits the rapid screening of constructs produced by homologous recombination or direct cloning methods. As used herein, these constructs are preferably viruses, but may include other types of vectors, such as a cosmid, episome, plasmid, or other genetic element that delivers a heterologous molecule to cells.

In one desired embodiment, the method utilizes the gene encoding green fluorescent protein (GFP), to provide a green-white selection method in which the presence of a recombinant is detected by the absence of GFP expression (i.e., the recombinants are observed as white in a green background). Alternatively, the method may utilize another suitable marker genes, including, without limitation, other fluorescent proteins and luciferase.

In one example, the method is used for production of a recombinant construct from homologous recombination of co-transected vectors into a selected host cell. As used herein, a host cell may be readily selected from an biological organism, including prokaryotic and eukaryotic cells, such as those discussed in the section related to production of a recombinant viral particle. Selection of the host cell is not a limitation of the present invention.

Suitably, each of the vectors contains the marker gene (e.g., GFP) under the control of a promoter that directs expression thereof in a host cell. Alternatively, each of the parental

vectors may contain a different marker gene that allows them to be distinguished not only from the recombinant construct produced, but also from each other. Preferably, where prokaryotic GFP is utilized, it is under the control of a prokaryotic promoter such as the promoter from lacZ. However, other suitable prokaryotic or non-prokaryotic promoters may be readily selected from among the promoters described herein and known to those of skill in the art. Advantageously, the GFP protein is placed in the portion of the vectors that are eliminated during homologous recombination and thus, the GFP protein is absent from the recombinant vector produced. In this manner, the presence of unrecombined parental vectors are readily detected under a phase contrast fluorescent microscope (or other suitable detection means) as expressing the marker gene and the recombinant constructs lack expression of the marker. In the methods in which both parent vectors utilize GFP, the recombinant appears as white in a background of green.

In another example, the method is used for production of a recombinant construct involving homologous recombination, in which the host cell stably contains at least one of the parental constructs to be utilized for production of the recombinant construct. In this embodiment, the host cell can be subjected to a single transfection. In still other embodiments, the method of the invention may be utilized for triple transfections. As with the double transfection described above, the parental constructs may contain the same marker gene or may contain different marker genes.

In another example, the method of the invention is used from production of a recombinant construct by direct cloning. Suitably, in this embodiment, the marker gene is present in that portion of the parent construct which is deleted during the cloning process. For example, the marker gene expression cassette (i.e., the gene, promoter, and any other necessary regulatory sequences) is engineered into the E1- or E3-region of an adenoviral vector, into which a transgene or minigene cassette will be cloned. The success of direct cloning into the target region can be readily detected by the absence of marker gene expression.

Optionally, the method of the invention can be readily assembled in the form of a kit which is available in a commercially useful format for production of recombinant constructs, e.g., recombinant adenoviruses. Typically, such kits will include plasmid backbones

containing a desired viral genome containing a marker gene inserted at a point upstream or downstream of the recombination site, as appropriate, or a plasmid backbone containing the marker gene inserted at the splice site for direct cloning of a heterologous gene. Such a kit can further include appropriate culture media, host cells, a test control, instructions, and other suitable materials.

In the examples below, this method is used in production of adenoviruses. However, it will be readily understood that this method may be readily adapted for use in generating other types of adenoviral, or non-adenoviral viral vectors.

The following examples are provided to illustrate the invention and do not limit the scope thereof. One skilled in the art will appreciate that although specific reagents and conditions are outlined in the following examples, modifications can be made which are meant to be encompassed by the spirit and scope of the invention.

Example 1 - Creation of an E1 deleted vector based on Chimpanzee Adenovirus C68 Using Green-white Selection Of Recombinants

A replication defective version of C68 was isolated for use in gene transfer. The classic strategy of creating a recombinant with E1 deleted, by homologous recombination in an E1 expressing cell line was pursued. The first step was creation of a plasmid containing m.u. 0 through 1.3 followed by addition of a minigene expressing enhanced green fluorescent protein (GFP) from a CMV promoter and C68 sequence spanning 9-16.7 m.u. This linearized plasmid was cotransfected into an E1 expressing cell line with Ssp I-digested C68 plasmid (SspI cuts at 3.6 m.u. leaving 4644 bp for homologous recombination). Experiments were initially conducted with 293 cells which harbor E1 from human Ad5 with the hope that this would suffice for transcomplementation. Indeed, plaques formed which represented the desired recombinant. The resulting vector was called C68-CMV-GFP.

The strategy for generating recombinants was modified to enable efficient and rapid isolation of recombinants. First, the alkaline phosphatase DNA in the initial shuttle vector was replaced with a prokaryotic GFP gene driven by the prokaryotic promoter from lacZ.

This allowed efficient screening of bacterial transformations when attempting to incorporate a desired eukaryotic RNA pol II transcriptional unit into the shuttle vector. The resulting transformation can be screened for expression of GFP; white colonies are recombinants while green colonies are residual parental plasmid.

5 A green-white selection has been used to screen the products of cotransfection for the isolation of human Ad5 recombinants (A.R. Davis et al, *Gene Thera.*, 5:1148-1152 (1998)). In the present system, and in contrast to Davis, the initial shuttle vector was revised to include extended 3' sequences from 9 to 26 MU. This vector was cotransfected with viral DNA from the original C68-CMV-GFP isolate that had been restricted with Xba I, which cuts at MU
10 16.5 allowing for 9.5 Kb of overlap for homologous recombination. The resulting plaques were screened under a phase contrast fluorescent microscope for non-fluorescing isolates that represent the desired recombinants. This greatly simplified screening in comparison to the standard methods based on structure or transgene expression. Thus, this method may be readily adapted for use in generating other types of adenoviral, or non-adenoviral viral
15 vectors.

A. Shuttle Plasmid

To construct a plasmid shuttle vector for creation of recombinant C68 virus, the plasmid pSP72 (Promega, Madison, WI) was modified by digestion with Bgl II followed by filling-in of the ends with Klenow enzyme (Boehringer Mannheim, Indianapolis, IN) and
20 ligation with a synthetic 12 bp Pac I linker (New England Biolabs, Beverly, MA) to yield pSP72-Pac. A 456 bp Pac I/SnaB I fragment spanning map unit (m.u. or MU) 0-1.3 of the C68 genome was isolated from the pNEB-BamE plasmid containing BamHI E fragment of the C68 genome and cloned into Pac I and EcoR V treated pSP72-Pac to yield pSP-C68-MU 0-1.3. A minigene cassette consisting of the cytomegalovirus early promoter driving lacZ with
25 a SV40 poly A signal was separated from pCMV β (Clontech, Palo Alto, CA) as a 4.5 kb EcoRI/SalI fragment and ligated to pSP-C68-MU 0-1.3 restricted with the same set of enzymes, resulting in pSP-C68-MU 0-1.3-CMVLacZ.

For the initial step in the isolation of the 9-16.7 MU region of C68, both pGEM-3Z (Promega, Madison, MI) and pBS-C68-BamF were double-digested with BamHI
30 and Sph I enzymes. Then the 293 bp fragment from pBS-C68-BamF was ligated with pGEM-

3Z backbone to form pGEM-C68-MU 9-9.8. A 2.4 kb fragment including the C68 MU 9.8-16.7 was obtained from the pBS-C68 BamHB clone after XbaI digestion, filling in reaction and subsequent BamHI treatment and cloned into BamHI/SmaI double digested pGEM-C68-MU 9-9.8 to generate pGEM-C68-MU 9-16.7. The C68 9-16.7 m.u. region was isolated from pGEM-C68-MU 9-16.7 by digestion with EcoRI, filling in of the ends with Klenow enzyme (Boehringer Mannheim, Indianapolis, IN), ligation of a synthetic 12 bp HindIII linker (NEB) and then digestion with HindIII. This 2.7 kb fragment spanning the C68 MU 9-16.7 was cloned into the HindIII site of pSP-C68-MU 0-1.3-CMVlacZ to form the final shuttle plasmid pC68-CMV-LacZ. In addition, an 820 bp alkaline phosphatase (AP) cDNA fragment was isolated from pAdCMVALP (K. J. Fisher, et al., *J. Virol.*, 70:520-532 (1996)) and exchanged for lacZ at Not I sites of pC68-CMV-lacZ, resulting in pC68-CMV-AP.

B. Construction of Recombinant Virus

To create the E1-deleted recombinant C68-CMVEGFP vector, a pC68-CMV-EGFP shuttle plasmid was first constructed by replacing the lacZ transgene in pC68-CMV-lacZ with the enhanced green fluorescent protein (EGFP) gene. The replacement cloning process was carried out as the follows. An additional NotI restriction site was introduced into the 5' end of the EGFP coding sequence in the pEGFP-1 (Clontech, Palo Alto, CA) by BamHI digestion, filling in reaction and ligation of a 8 bp synthetic NotI linker (NEB). After NotI restriction of both constructs, the EGFP sequence was isolated from the modified pEGFP-1 and used to replace the lacZ gene in the pC68-CMV-lacZ. The pC68-CMVEGFP construct (3 µg) was co-transfected with Ssp I-digested C68 genomic DNA (1 µg) into 293 cells for homologous recombination as previously described (G. Gao, et al., *J. Virol.*, 70:8934-8943 (1996)). Green plaques visualized by fluorescent microscopy were isolated for 2 rounds of plaque purification, expansion and purification by CsCl gradient sedimentation (G. Gao, et al., cited above).

The invention provides a uniquely modified version of the green/white selection process (A. R. Davis, et al., *Gene Thera.*, 5:1148-1152 (1998)). The present example illustrates use of this method for construction of recombinant C68 vectors. A 7.2 kb fragment spanning 9 to 36 MU was isolated from the pBSC68-BamB plasmid by treatment with AgeI and BsiwI restriction endonucleases and cloned into Asp718 and AgeI sites of

pC68-CMV-AP shuttle plasmid, resulting in a new plasmid called pC68CMV-AP-MU36. A further modification was made to remove 26 to 36 m.u. from pC68CMV-AP-MU36 by Eco47III and NruI digestions. The new shuttle plasmid called pC68CMV-AP-MU26 has a shorter region for homologous recombination (i.e., 16.7-26 MU) 3' to the minigene. To make a recombinant C68 vector, alkaline phosphatase (AP) is replaced with the gene of interest. The resulting pC68CMV-Nugene-MU26 construct is co-transfected with Xba I (16.5 MU) restricted C68-CMVGFP viral DNA into 293 cells, followed by top agar overlay. The recombinant virus plaques (white) are generated through the homologous recombination in the region of 16.7-26 MU which is shared between pC68CMV-Nugene construct and C68 viral backbone; the recombinants which form white plaques are selected from green plaques of uncut C68-CMVGFP virus.

The green/white selection mechanism was also introduced to the process of cloning of the gene of interest into the pC68 shuttle plasmid. The AP gene in both pC68CMV-AP-MU36 and pC68CMV-AP-MU26 was replaced with a cassette of prokaryotic GFP gene driven by the lacZ promoter isolated from pGFP-MU31 (Clontech, Palo Alto, CA). Thus, white colonies of bacterial transformants will contain the recombinant plasmid. This green/white selection process for bacterial colonies circumvented the need for making and characterizing large numbers of miniprep DNAs and so further enhanced the efficiency in creating recombinant C68 vectors.

Example 2 - Chimpanzee C68 Virus Stock and Replication

Examples 3 through 5 which follow provide additional characterization of the chimpanzee C68. It will be appreciated by one of skill in the art that this information can be readily used in the construction of novel recombinant chimpanzee adenoviral constructs.

The C68 virus stock was obtained from ATCC (Rockville, MD) and propagated in 293 cells (ATCC) cultured in DMEM (Sigma, St. Louis, MO) supplemented with 10% fetal calf serum (FCS; Sigma or Hyclone, Logan, UT) and 1% Penicillin-Streptomycin (Sigma). Infection of 293 cells was carried out in DMEM supplemented with 2% FCS for the first 24 hours, after which FCS was added to bring the final concentration to 10%. Infected cells were harvested when 100% of the cells exhibited virus-induced cytopathic effect (CPE),

collected, and concentrated by centrifugation. Cell pellets were resuspended in 10mM Tris (pH 8.0), and lysed by 3 cycles of freezing and thawing. Virus preparations were obtained following 2 ultra centrifuge steps on cesium chloride density gradients and stocks of virus were diluted to 1×10^{12} particles/ml in 10mM Tris/100mM NaCl/50% glycerol and stored at
 5 -70 °C.

Example 3 - Cloning and sequencing of viral genomic DNA

Genomic DNA was isolated from the purified virus preparation following standard methods and digested with a panel of 16 restriction enzymes following the manufacturer's
 10 recommendations. Except as noted, all restriction and modifying enzymes were obtained from Boehringer Mannheim, Indianapolis, IN. Genomic DNA was digested with BamHI, PstI, Sall, HindIII or XbaI and the fragments were subcloned into plasmids (K. L. Berkner and P.A. Sharp, *Nucl. Acids Res.*, 11:6003-20 (1983)). After deproteination, synthetic 10bp PacI linkers (New England Biolabs, Beverly, MA) were double digested with PacI and
 15 BamHI, or PstI.

The PstI, BamHI and HindIII clones generated from C68 are illustrated in Figure 1, parts C, D and E, respectively. The fragments indicated by the shaded boxes were not cloned, but the sequence of the entire genome has been determined through sequencing overlapping clones and viral DNA directly (unshaded boxes). The cloned fragments and insert sizes are
 20 described in Table 1. In the following table, pBS = pBluescript SK+ clone; pNEB = pNEB 193 clone; pBR = pBR322 clone; No prefix = fragment not cloned

Table 1. C68 plasmid clones and insert sizes

Construct Name	Insert Size (base pairs)	Fragment 5' End	Fragment 3' End	5' End Map Unit	3' End Map Unit
Pst-I Fragments					
C68-Pst-A	6768	24784	31551	67.9%	86.4%
pBS:C68-Pst-B	6713	4838	11550	13.2%	31.6%
pBS:C68-Pst-C	5228	14811	20038	40.6%	54.9%
pBS:C68-Pst-D	2739	12072	14810	33.1%	40.6%
pBS:C68-Pst-E	2647	20039	22685	54.9%	32.1%
pBS:C68-Pst-F	1951	32046	33996	87.8%	93.1%
pNEB:C68-Pst-G	1874	1	1874	0.0%	5.1%
pBS:C68-Pst-H	1690	23094	24783	63.2%	67.9%
pBS:C68-Pst-I	1343	33997	35339	93.1%	96.8%
pNEB:C68-Pst-J	1180	35340	36519	96.8%	100.0%
pBS:C68-Pst-K	1111	2763	3873	7.6%	10.6%
pBS:C68-Pst-L	964	3874	4837	10.6%	13.2%
pBS:C68-Pst-M	888	1875	2762	5.1%	7.6%
pBS:C68-Pst-N	408	22686	23093	62.1%	63.2%
C68-Pst-O	380	31666	32045	86.7%	87.7%
pBS:C68-Pst-P	285	11551	11835	31.6%	32.4%
C68-Pst-Q	236	11836	12071	32.4%	33.1%
pBS:C68-Pst-R	114	31552	31665	86.4%	86.7%
BamHI Fragments					
C68-Bam-A	16684	19836	36519	54.3%	100.0%
pBS:C68-Bam-B	8858	3582	12439	9.8%	34.1%
pBS:C68-Bam-C	4410	12440	16849	34.1%	46.1%
pBS:C68-Bam-D	2986	16850	19835	46.1%	54.3%
pNEB:C68-Bam-E	2041	1	2041	0.0%	5.6%
pBS:C68-Bam-F	1540	2042	3581	5.6%	9.8%
HindIII Fragments					
pBR:C68-Hind-B	9150	23471	32620	64.3%	89.3%

Chimpanzee adenovirus, C68, was obtained from ATCC and propagated in human 293 cells. Viral genomic DNA was isolated from purified virions using established procedures (A. R. Davis, *et al.*, *Gene Thera.*, **5**:1148-1152 (1998)) and digested with a panel of restriction enzymes; the data were consistent with previous studies (data not shown) (G. R. Kitchingman, *Gene*, **20**:205-210 (1982); Q. Li and G. Wadell, *Arch Virol.* **101**:65-77 (1998); R. Wigand, *et al.*, *Intervirology.* **30**:1-9 (1989)). Restriction fragments spanning the entire genome of C68 were subcloned into plasmids. A schematic drawing of the C68 genome is shown in Figure 1A, and the Pst-I, BamHI and HindIII fragments that were cloned into plasmid vectors are indicated by the unshaded boxes, in Figs. 1B, 1C, and 1D, respectively.

The cloned fragments, fragment sizes and genomic position are also listed in Table 1. Both plasmid clones and genomic DNA were used as templates for sequencing. The genome was sequenced by primer walking in both directions and each base was included in an average of approximately four reactions.

The C68 genome is 36521 bp in length [see, US Patent 6,083,716]. Preliminary comparison with GenBank sequences indicated varying degrees of similarity with other human and animal adenoviruses along the entire length of the viral genome. Regions with homology to all of the previously described adenoviral genetic units, early regions 1-4 and the major late genes, were found in the C68 genome (Fig. 1A). DNA homology between C68 and the human adenoviruses that have been completely sequenced, Ad2 (NC001405), Ad5 (NC001405), Ad12 (NC001460), Ad17 (NC002067) and Ad40 (NC01464), was used to order the clones. The open reading frames (ORF) were determined and the genes were identified based on homology to other human adenoviruses. All of the major adenoviral early and late genes are present in C68. The inverted terminal repeats (ITR=s) are 130 bp in length.

Example 4 - Analysis of C68 sequence

The complete nucleotide sequence of every member of the Mastadenovirus genus accessible from GenBank, including isolates from different species, were screened for identity to C68. The Ad4 minigenome was assembled from the following GenBank sequences: Left-hand ITR (J01964); E1A region (M14918); DNA pol and pTP (X74508, 74672); VA RNA-I, II (U10682); 52, 55K (U52535); pVII (U70921); hexon (X84646);

endoprotease (M16692); DNA-binding protein (M12407); fiber (X76547); right-hand ITR (J01965). The Ad7 composite genome was created from the following sequence data: Mu 3-21 (X03000); VA RNA-I, II, pTP & 52, 55K (U52574); penton (AD001675); pVI, hexon and endoprotease (AF065065); DNA-binding protein (K02530); E3 and fiber region (AF104384);
5 right-hand ITR (V00037).

The amino acid sequence alignment was generated with Clustal X, edited with Jalview (<http://www.ebi.ac.uk/~michele/jalview/>), and analyzed with Boxshade (http://www.ch.embnet.org/software/BOX_form.html). Publicly available hexon protein sequences from all human adenovirus serotypes were initially aligned to identify the set
10 showing the highest homology to C68.

The nucleotide sequence and predicted amino acid sequences of all significant open reading frames in the C68 genome were compared to known DNA and protein sequences. The nucleotide sequence of C68 was compared to sequences of Ad 2, 4, 5, 7, 12, 17 and 40. In agreement with previous restriction analysis (Kitchingman, cited above) C68 is most
15 similar to human Ad4 (subgroup E).

The E1A region of C68 extends from the TATA box at nt 480 to the poly A addition site at 1521. The consensus splice donor and acceptor sites are in the analogous position of the human Ad counterparts, and the 28.2K and 24.8K proteins are similar in size to the human Ad proteins. The ORF for the smallest E1A protein of C68 is predicted to encode 101
20 residues as opposed to approximately 60 amino acids for other adenoviruses. There is a TTA codon at residue 60 for C68 where other adenoviruses often have a TGA stop codon. The first 60 residues of C68 E1A 100R protein have 85% identity to the Ad4 homolog.

The C68 genome encodes genes for the four E1B proteins, 20.5K, 54.7K, 10.1K and 18.5K as well as pIX. All five C68 encoded proteins are similar in size to that of other Ad
25 E1B and pIX proteins. The Ad4 homolog of the E1B 21K protein has only 142 amino acids, where C68 has 186 residues and other human adenoviruses have 163-178 residues. The C68 and Ad4 proteins share 95% identity over the first 134 aa, then the similarity ends and the Ad4 protein terminates at 142 amino acids.

The C68 genome encodes homologs of the E2A 55K DNA binding protein and the
30 Iva2 maturation protein, as well as the E2B terminal protein and the DNA polymerase. All of

the E2 region proteins are similar in size to their human Ad counterparts, and the E2B proteins are particularly well conserved. The C68 E2B 123.6K DNA polymerase is predicted to be 1124 residues, while Ad4 is predicted to have 1193 although the other human adenoviruses have smaller polymerases. Residues 1-71 of the Ad4 polymerase have no
5 similarity to any other Ad polymerase, and it is possible that this protein actually initiates at an internal ATG codon. From amino acids 72-1193, Ad4 and C68 polymerases have 96% amino acid identity.

The E3 regions of human adenoviruses sequenced so far exhibit considerable sequence and coding capacity variability. Ad40 has five E3 region genes, Ad12 has six, C68
10 and Ad5 have seven, Ad38 has eight and Ad3 as well as Ad7 (subgroup B human adenoviruses) have nine putative E3 region genes. The Ad4 E3 region has not yet been sequenced. In comparison with the E3 region of Ad35, all 7 E3 gene homologs were identified in the C68 genome (C. F. Basler and M.S. Horwitz, *Virology*, **215**: 165-177 (1996)).

15 The C68 E4 region has 6 ORFs and each is homologous to proteins in the human Ad5, 12 and 40 E4 region. The E4 nomenclature is confusing because the ORF2 homologs of C68, Ad12 and Ad40 are approximately 130 residues, while in Ad5 there are two ORFs encoding proteins of 64 and 67 residues with homology, respectively, to the amino and carboxy terminal ends of the larger ORF2 proteins. ORF5 has been omitted in our nomenclature
20 because the 5th ORF in the E4 region is homologous to the widely studied ORF6 protein of human Ad5.

The major late promoter and the tri-partite leader sequences of the C68 genome were located. ORFs with the potential to encode the 15 major late proteins were located. All of the C68 late proteins are similar in size to their human Ad counterparts. The percent amino
25 acid identity between chimpanzee and human Ad late proteins varies considerably. The C68 fiber protein is predicted to have 90% amino acid identity with the Ad4 protein, but much less similarity to the other human Ad fiber proteins. The CAR binding site in the fiber knob is present in C68.

Example 5 - Virus neutralizing antibody assays

Several studies were performed to determine if there is cross-reactivity between type specific antisera of C68 and human adenovirus. The neutralizing activity of sera was tested as follows. Panels of sera from normal human subjects (N=50), rhesus monkeys (N=52) and chimpanzees (N=20) were evaluated for neutralizing antibodies against Ad5 and C68 based vectors using 293 cells as an indicator cell line. Sera collected from individual humans, rhesus monkeys, or chimpanzees were inactivated at 56 °C for 30 minutes. A serial dilution of each sample (1:10, 1:20, 1:40, 1:80, 1:160, 1:320 in 100 µl of DMEM containing 10% FCS) was added to equal amounts of H5.010CMVEGFP (1000 PFU/well) or C68CMVEGFP virus and incubated at 4 °C for two hrs. One hundred and fifty microliters of the mixture were transferred onto 2 x 10⁵ 293 cells in 96 well flat bottom plates. Control wells were infected with equal amounts of virus (without addition of serum). Samples were incubated at 37 °C in 5% CO₂ for 48 hrs and examined under a fluorescent microscope. Sample dilutions that showed >50% reduction of green-fluorescent foci as compared to infected controls were scored positive for neutralizing antibodies.

As expected, approximately 35% of normal human subjects demonstrated neutralizing antibody against Ad5, a frequency much higher than observed in sera of rhesus monkeys and chimpanzee. Neutralizing antibody to C68 was observed in 80% of chimpanzee and only 2% of normal human subjects or rhesus monkeys. Titers of neutralizing antibodies in the non-target species were generally low.

To further evaluate cross-reactivity of C68 with human adenovirus vectors, mice were immunized with 2 x 10⁷ plaque forming units (pfu) of Ad 2, 4, 5, 7 and 12 as well as C68. Sera were harvested 2 weeks later and tested for antibodies that neutralized either Ad5 or C68 vectors. Neutralizing antibody to Ad5 vector was only detected in animals immunized with Ad5. Importantly, the only animals with neutralizing antibody to C68 vector were those immunized with C68 vector; none of the human serotypes tested, including Ad4, generated antibodies in mice that neutralized C68 *in vitro*.

Important to the utility of C68 vector in human trials is the absence of neutralizing antibody in the human population. In our study, a screen of 50 normal human subjects failed to detect any significant neutralizing antibodies (>1:10) using the same assay that showed

neutralizing antibodies in >50% of chimpanzees. Furthermore, sera of mice immunized with multiple human Ad serotypes including Ad4, did not neutralize infection with C68.

Example 6 - Structural analysis of hexon proteins

5 The absence of neutralizing antibodies between C68 and human serotypes compelled us to more carefully evaluate structural differences in the regions of hexon presumed to harbor type specific epitopes. Previous studies have suggested that these epitopes are located within the 7 hypervariable regions of hexon determined by Crawford-Miksza and Schnurr (*J. Virol*, 70:1836-1844 (1996)). A comparison of the amino acid sequences of hexon proteins
10 between C68 and several human adenoviruses is shown in Figure 3. Indeed, C68 is substantially dissimilar in significant regions of these hypervariable sequences.

Example 7 - Construction of C68-Derived Capsid Containing a Human Fiber Gene

15 To generate a C68-derived vector with an altered tropism, the chimeric fiber gene construct containing the Ad5 fiber knob fused to the C68 tail and shaft is incorporated into a plasmid carrying the C68 genome. For the precise replacement of the wild-type C68 fiber gene, a plasmid carrying the green fluorescent protein driven by a CMV promoter is used for modification of C68 fiber. The resulting transfer vector contains a CMV promoter driven
20 green fluorescent protein (GFP) expression cassette inserted into the E3 region, the chimeric C68/Ad5 fiber gene, and E4. This transfer vector was used for incorporation of GFP cassette and modified fiber gene into the backbone of an E3 deleted C68 infectious plasmid via homologous recombination in *E. coli*. The viral genome was released from the plasmid by PacI digestion and used to transfect 293 cells. The chimeric C68-derived virus is produced
25 about 3 weeks following transfection using techniques described herein. Similar techniques can be readily utilized to construct other C68-derived capsids.

 All publications cited in this specification are incorporated herein by reference. While the invention has been described with reference to a particularly preferred embodiment, it will be appreciated that modifications can be made without departing from
30 the spirit of the invention. Such modifications are intended to fall within the scope of the appended claims.

What is claimed is:

1. A C68 chimpanzee adenoviral capsid protein which is substantially free of other viral proteins with which it is naturally associated, said protein selected from the group consisting of:

(a) a hexon protein characterized by the amino acid sequence of SEQ ID NO:17;

(b) a penton protein characterized by the amino acid sequence of SEQ ID NO:12;

(c) a fiber protein characterized by the amino acid sequence of SEQ ID NO:27;

(d) a unique fragment of any of (a) to (c) comprising 8 amino acid residues in length.

2. The C68 chimpanzee adenoviral capsid protein according to claim 1, wherein the fragment of hexon protein (a) is selected from the group consisting of:

(i) about amino acids 125 to 443 of SEQ ID NO:16;

(ii) about amino acids 131 to about 441 of SEQ ID NO:16;

(iii) about amino acids 138 to about 441 of SEQ ID NO:16;

(iv) about amino acids 138 to about 163 of SEQ ID NO:16;

(v) about amino acids 170 to about 176 of SEQ ID NO:16;

(vi) about amino acids 195 to about 203 of SEQ ID NO:16;

(vii) about amino acids 233 to about 246 of SEQ ID NO:16;

(viii) about amino acids 253 to about 264 of SEQ ID NO:16;

(ix) about amino acids 287 to about 297 of SEQ ID NO:16; and

(x) about amino acids 404 to about 430 of SEQ ID NO:16.

3. The C68 chimpanzee adenoviral capsid protein according to claim 1, wherein the fragment of fiber protein (c) is about amino acids 247 to 425 of SEQ ID NO:27.

4. A novel adenovirus serotype comprising a unique fragment of the C68 hexon protein according to claim 1 fused to a heterologous adenovirus hexon peptide.

5. The novel adenovirus serotype according to claim 4, wherein said fragment is selected from the group consisting of:

- (a) about amino acids 125 to about 443 of SEQ ID NO:16;
- (b) about amino acids 131 to about 441 of SEQ ID NO:16;
- (c) about amino acids 138 to about 441 of SEQ ID NO:16;
- (d) about amino acids 138 to about 163 of SEQ ID NO:16;
- (e) about amino acids 170 to about 176 of SEQ ID NO:16;
- (f) about amino acids 195 to about 203 of SEQ ID NO:16;
- (g) about amino acids 233 to about 246 of SEQ ID NO:16;
- (h) about amino acids 253 to about 264 of SEQ ID NO:16;
- (i) about amino acids 287 to about 297 of SEQ ID NO:16; and
- (j) about amino acids 404 to about 430 of SEQ ID NO:16.

6. The novel adenovirus serotype according to claim 4, wherein the C68 fragment is a loop region of the hexon.

7. The novel adenovirus serotype according to claim 4, wherein at least one loop region of the C68 hexon protein is substituted with a loop region from a heterologous adenovirus serotype.

8. A recombinant adenovirus comprising a capsid of the novel adenovirus serotype of any of claims 2 to 7 encapsidating a molecule for delivery to a target cell.

9. The recombinant virus according to claim 8, wherein the molecule comprises an adenovirus 5' inverted terminal repeat sequence (ITRs), a minigene, and an adenovirus 3' ITR.

10. The recombinant adenovirus according to claim 9, wherein the adenovirus ITRs are from a serotype heterologous to C68.
11. An adenoviral capsid comprising a C68 capsid protein according to claim 1 and a protein from a heterologous adenoviral capsid protein.
12. The adenoviral capsid according to claim 11, wherein the capsid comprises the C68 hexon and at least the penton or the fiber is from a different adenovirus serotype.
13. The adenoviral capsid according to claim 12, wherein the penton and fiber are from a human adenovirus serotype.
14. The adenoviral capsid according to claim 12, wherein the penton is from C68 and the fiber is from a different adenovirus serotype.
15. The adenoviral capsid according to claim 12, wherein the fiber is from C68 and the penton is from a different adenovirus serotype.
16. The adenoviral capsid according to claim 11, wherein the C68 protein is selected from the penton and the fiber and the hexon is from a different adenovirus serotype.
17. A recombinant adenovirus comprising an adenoviral capsid according to any of claims 11 to 16, said capsid encapsidating a molecule for delivery to a target cell.
18. A pharmaceutical composition comprising a physiologically acceptable carrier and a recombinant virus according to claim 17.
19. Use of a recombinant adenovirus according to claim 8 or claim 17 for delivering a molecule to a target cell.
20. Use according to claim 19, wherein said molecule is an immunogen.

21. Use according to claim 19, wherein said molecule is a therapeutic molecule.
22. A pseudotyped adenovirus comprising a C68 adenoviral capsid having an amino acid sequence as provided in SEQ ID NO:16 encapsidating a molecule comprising adenovirus 5' inverted terminal repeat sequences (ITR), a minigene, and adenovirus 3' ITR, wherein each of the 5' and 3' ITRs are from a serotype heterologous to C68.
23. A pharmaceutical composition comprising a physiologically acceptable carrier and a pseudotyped adenovirus according to claim 22.
24. Use of a pseudotyped adenovirus according to claim 22 for delivering a molecule to a target cell.
25. Use according to claim 22, wherein said molecule is an immunogen.
26. Use according to claim 22, wherein said molecule is a therapeutic molecule.
27. A method for rapid screening of recombinant constructs comprising the steps of:
 - (a) providing a first parent construct having a green fluorescent protein (GFP), operably linked to regulatory sequences which permit its expression in a host cell, downstream of the portion of the first parent construct to be in the recombinant construct;
 - (b) providing a second parent construct having a GFP operably linked to regulatory sequences which permit its expression in a host cell upstream of the portion of the second parent construct to be in the recombinant construct;
 - (c) culturing a selected host cell with the first parent construct and second parent construct under conditions which permits them to homologously recombine; and
 - (d) screening for the absence expression of GFP expression, which indicates the presence of the recombinant construct.

28. The method according to claim 27, wherein the regulatory sequence comprises the prokaryotic promoter from lacZ.

29. The method according to claim 27, wherein the first parent construct is a plasmid.

30. The method according to claim 27, wherein the second parent constructs is a plasmid.

31. The method according to claim 27, wherein the first parent construct and the second parent construct are provided by co-transfection into the cell.

32. The method according to claim 27, wherein one of the first parent construct or the second parent construct is stably contained within the host cell.


33. A host cell comprising:

- (a) a first parent construct having a green fluorescent protein (GFP), operably linked to regulatory sequences which permit its expression in a host cell, downstream of the portion of the first parent construct to be in the recombinant construct; and
- (b) a second parent construct having a GFP operably linked to regulatory sequences which permit its expression in a host cell upstream of the portion of the second parent construct to be in the recombinant construct.

1/4

BEST AVAILABLE COPY

FIG. 1



2/4

BEST AVAILABLE COPY

FIG. 2

C68 - E 130	NTCQWTKADGETA-TE	NTCYGNA PVQGIN	ITKDGIOLGTD-T	DDOPI
Ad4 - E 130	NTCQW--K-DSBS	KMHT-GAAAMPCTGKKIEADGIPRIDSTS	ITKDGIOLGTD-T	DDOPI
Ad16 - E 130	NTCQW--K-DSBS	KMHT-GVAAAMPCTGKKIEADGIPRIDSTS	ITKDGIOLGTD-T	DDOPI
Ad3 - B 130	NTSQWIVTNGNAVTT	TTNT-GIASMKCN	ITKDGIOLGTD-T	DDOPI
Ad7 - B 130	NTSQWIVTAGEEAVTT	TTNT-GIASMKCN	ITKDGIOLGTD-T	DDOPI
Ad2 - C 130	NSCEWEQTEDSGRAVAE	DEEEEEEEQNARDQATKTHVACAPSGSET	ITKDGIOLGTD-T	DDOPI
		-----HVR1-----	-----HVR2-----	
C68 - E 177	YADKTYQPEPQVGDAEWH	ITGTDEKYGGRALKPDTKMKPCYGSFAKPTNKEGGQA-N	VK-TGTG-TTKEYDIDMAF	
Ad4 - E 177	YADKTYQPEPQVGDAEWH	ITGTDEKYGGRALKPDTKMKPCYGSFAKPTNKEGGQA-N	VK-TGTG-TTKEYDIDMAF	
Ad16 - E 177	YADKTYQPEPQVGDAEWH	ITGTDEKYGGRALKPDTKMKPCYGSFAKPTNKEGGQA-N	VK-TGTG-TTKEYDIDMAF	
Ad3 - B 183	YADKTYQPEPQVGDAEWH	ITGTDEKYGGRALKPDTKMKPCYGSFAKPTNKEGGQA-N	VK-TGTG-TTKEYDIDMAF	
Ad7 - B 181	YADKTYQPEPQVGDAEWH	ITGTDEKYGGRALKPDTKMKPCYGSFAKPTNKEGGQA-N	VK-TGTG-TTKEYDIDMAF	
Ad2 - C 205	YADKTYQPEPQVGDAEWH	ITGTDEKYGGRALKPDTKMKPCYGSFAKPTNKEGGQA-N	VK-TGTG-TTKEYDIDMAF	
		-----HVR3-----	-----HVR4-----	
C68 - E 251	FDNRSAA	AAGLAPEIIVLYTENVDLETPDTHIVYKAGTDDSSSINLQQQAMPNRPNYIGFRDNFIFGLWYVNSTGNN		
Ad4 - E 253	FDNRSAA	AAGLAPEIIVLYTENVDLETPDTHIVYKAGTDDSSSINLQQQAMPNRPNYIGFRDNFIFGLWYVNSTGNN		
Ad16 - E 253	FDNRSAA	AAGLAPEIIVLYTENVDLETPDTHIVYKAGTDDSSSINLQQQAMPNRPNYIGFRDNFIFGLWYVNSTGNN		
Ad3 - B 259	FDNRSAA	AAGLAPEIIVLYTENVDLETPDTHIVYKAGTDDSSSINLQQQAMPNRPNYIGFRDNFIFGLWYVNSTGNN		
Ad7 - B 130	FDNRSAA	AAGLAPEIIVLYTENVDLETPDTHIVYKAGTDDSSSINLQQQAMPNRPNYIGFRDNFIFGLWYVNSTGNN		
Ad2 - C 277	FDNRSAA	AAGLAPEIIVLYTENVDLETPDTHIVYKAGTDDSSSINLQQQAMPNRPNYIGFRDNFIFGLWYVNSTGNN		
		-----HVR5-----		
C68 - E 327	GVLAGQASQLNAVVDLQDRNTELSYQLLLDLSLGDRTTRYFSMWNAQVDSYDPDVRRIENHGVDELPNYCFPLDVGRTDT			
Ad4 - E 328	GVLAGQASQLNAVVDLQDRNTELSYQLLLDLSLGDRTTRYFSMWNAQVDSYDPDVRRIENHGVDELPNYCFPLDVGRTDT			
Ad16 - E 328	GVLAGQASQLNAVVDLQDRNTELSYQLLLDLSLGDRTTRYFSMWNAQVDSYDPDVRRIENHGVDELPNYCFPLDVGRTDT			
Ad3 - B 338	GVLAGQASQLNAVVDLQDRNTELSYQLLLDLSLGDRTTRYFSMWNAQVDSYDPDVRRIENHGVDELPNYCFPLDVGRTDT			
Ad7 - B 334	GVLAGQASQLNAVVDLQDRNTELSYQLLLDLSLGDRTTRYFSMWNAQVDSYDPDVRRIENHGVDELPNYCFPLDVGRTDT			
Ad2 - C 357	GVLAGQASQLNAVVDLQDRNTELSYQLLLDLSLGDRTTRYFSMWNAQVDSYDPDVRRIENHGVDELPNYCFPLDVGRTDT			
		-----HVR6-----		
C68 - E 407	YQGIKANG	DOTTWTKDSS-VNDANEIGKGNPFAM		
Ad4 - E 408	YQGIKANG	DOTTWTKDSS-VNDANEIGKGNPFAM		
Ad16 - E 408	YQGIKANG	DOTTWTKDSS-VNDANEIGKGNPFAM		
Ad3 - B 418	YQGIKANG	DOTTWTKDSS-VNDANEIGKGNPFAM		
Ad7 - B 414	YQGIKANG	DOTTWTKDSS-VNDANEIGKGNPFAM		
Ad2 - C 437	YQGIKANG	DOTTWTKDSS-VNDANEIGKGNPFAM		
		-----HVR7-----		

3/4

FIG. 3

Pan-9 fiber knob	(1)	TLWTPDPSPNCQILAENDAKLTLCLTKCGSQILATVSVLVGSG-NLNP
Ad 2 fiber knob	(1)	TLWTPDPSPNCRIHSDNDCKFTLVLTCKGSQVLATVAALAVSG--DLSS
Ad 5 fiber knob	(1)	TLWTPAPSPNCRNLNAEKDAKLTLVLTKCGSQILATVSVLAVKG--SLAP
Pan-9 fiber knob	(50)	ITGTVSSAQVFLRFDANGVLLTEHSTLKKYWGVRQGSIDGTPYTNAVGF
Ad 2 fiber knob	(49)	MTGTVASVSIFLRFDQNGVLMENSSSLKKHYWNFRNGNSTNANPYTNAVGF
Ad 5 fiber knob	(49)	ISGTVQSAHLIIRFDENGVLNNSFLDPEYWNFRNGDLTEGTAYTNAVGF
Pan-9 fiber knob	(100)	MPNLKAYPKSQSSTTKNNIVGQVYMNGDVSKPMLLTITLNGTDDS-----
Ad 2 fiber knob	(99)	MPNLLAYPKTSQTAKNNIVSQVYLHGDKTKPMLTITLNGTSESTETSE
Ad 5 fiber knob	(99)	MPNLSAYPKSHGKTAKSNIVSQVYLNCGDKTKPVTTLTITLNGTQET-GDTT
Pan-9 fiber knob	(145)	NSTYSMSFSYTW-T-NGSVVGATFGANSYTFFSYIAQE
Ad 2 fiber knob	(149)	VSTYSMSFTWSWE-SGKYTTETTFATNSYTFFSYIAQE
Ad 5 fiber knob	(148)	PSAYSMSFSWDWS-GHNYINEIFATSSYTFFSYIAQE

Hu5 Pan-9	APKGAPNPCEWDEAAATALEINLEEEEDDDNEDEVDEQAEEQKTHVFGQAPYSGINITKEGIQIGVEGQT-- APKGAPNTCQWTYKADG-----ETATEKTYTYGNAPVQGINITKDGIGLGTDTDD--
Hu5 Pan-9	--PKYADKTFQPEPQIGESQWYETEIN--HAAGRVLKKTTPMKPCYGSYAKPTNENGGQGILVKQQN--G -QPIYADKTYQPEPQVGVGDAEWHDITGTDEKYGGRALKPDTKMKPCYGSFAKPTNKEGGQANVKTGTG--T
Hu5 Pan-9	KLESQVEMQFFSTTEATAGNGDNLTPKVVLVYSEDVDIETPDTHISYMPPTIKEGNSRELMSGQSMNRPNY TKEYDIDMAFFDNRSAAG---LAPEIVLYTENVDLETPDTHIVYKAGTDDSSSSINLGQQAMPNRPNY
Hu5 Pan-9	IAFRDNFIGLMYYNSTGNMGVLAGQASQLNAVVDLQDRNTELSYQLLLDSIGDRTRYFSMWNQAVDSYDP IGFRDNFIGLMYYNSTGNMGVLAGQASQLNAVVDLQDRNTELSYQLLLDSIGDRTRYFSMWNQAVDSYDP
Hu5 Pan-9	DVRIIENHGTEDELPNYCFPLGGVINTETLTKVKPKTG-----QENGWEKDATEFSDKNEIRVGNNFAMEI DVRIIENHGVEDELPNYCFPLDAVGRDITYQGIKAN-----GTDQTTWTKDDSVN-DANEIGKGNPFAMEI

FIG. 4

SEQUENCE LISTING

<110> The Trustees of the University of Pennsylvania
 Gao, Guangping
 Wilson, James M.

<120> Method for Rapid Screening of Bacterial Transformants and
 Novel Simian Adenovirus Proteins

<130> UPN-N2630PCT

<150> US 60/300,501
 <151> 2001-06-22

<150> US 60/385,632
 <151> 2002-06-04

<160> 41

<170> PatentIn version 3.1

<210> 1
 <211> 101
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 1

Met	Arg	His	Leu	Arg	Asp	Leu	Pro	Asp	Glu	Lys	Ile	Ile	Ile	Ala	Ser
1				5					10					15	
Gly	Asn	Glu	Ile	Leu	Glu	Leu	Val	Val	Pro	Ser	Leu	Thr	Gln	Met	Met
			20					25					30		
Arg	Pro	Pro	Leu	Gln	Ser	Pro	Leu	Arg	His	Pro	Gln	Lys	Leu	Ala	His
			35				40					45			
Leu	His	Leu	Arg	Ile	Leu	Leu	Asp	Gln	Phe	Leu	Leu	Glu	Pro	Leu	Gly
	50					55					60				
Gly	Glu	Gln	Leu	Trp	Asn	Val	Trp	Met	Thr	Cys	Tyr	Arg	Val	Gly	Leu
65					70					75					80
Asn	Leu	Trp	Thr	Cys	Val	Pro	Gly	Asn	Ala	Pro	Gly	Thr	Lys	Cys	His
				85					90					95	
Thr	Cys	Val	Phe	Thr											
				100											

<210> 2
 <211> 257
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 2

Met	Arg	His	Leu	Arg	Asp	Leu	Pro	Asp	Glu	Lys	Ile	Ile	Ile	Ala	Ser
1				5					10					15	
Gly	Asn	Glu	Ile	Leu	Glu	Leu	Val	Val	Asn	Ala	Met	Met	Gly	Asp	Asp
			20					25					30		

Pro Pro Glu Pro Pro Thr Pro Phe Glu Thr Pro Ser Leu His Asp Leu
 35 40 45
 Tyr Asp Leu Glu Val Asp Val Pro Glu Asp Asp Pro Asn Glu Glu Ala
 50 55 60
 Val Asn Asp Phe Phe Ser Asp Ala Ala Leu Leu Ala Ala Glu Glu Ala
 65 70 75 80
 Ser Ser Ser Ser Ser Asp Ser Asp Ser Ser Leu His Thr Pro Arg Pro
 85 90 95
 Gly Arg Gly Glu Lys Lys Ile Pro Glu Leu Lys Gly Glu Glu Met Asp
 100 105 110
 Leu Arg Cys Tyr Glu Glu Cys Leu Pro Pro Ser Asp Asp Glu Asp Glu
 115 120 125
 Gln Ala Ile Gln Asn Ala Ala Ser Gln Gly Val Gln Ala Ala Ser Glu
 130 135 140
 Ser Phe Ala Leu Asp Cys Pro Pro Leu Pro Gly His Gly Cys Lys Ser
 145 150 155 160
 Cys Glu Phe His Arg Met Asn Thr Gly Asp Lys Ala Val Leu Cys Ala
 165 170 175
 Leu Cys Tyr Met Arg Ala Tyr Asn His Cys Val Tyr Ser Pro Val Ser
 180 185 190
 Asp Ala Asp Asp Glu Thr Pro Thr Thr Lys Ser Thr Ser Ser Pro Pro
 195 200 205
 Glu Ile Gly Thr Ser Pro Pro Glu Asn Ile Val Arg Pro Val Pro Val
 210 215 220
 Arg Ala Thr Gly Arg Arg Ala Ala Val Glu Cys Leu Asp Asp Leu Leu
 225 230 235 240
 Gln Gly Gly Val Glu Pro Leu Asp Leu Cys Thr Arg Lys Arg Pro Arg
 245 250 255

His

<210> 3
 <211> 226
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 3

Met Arg His Leu Arg Asp Leu Pro Asp Glu Lys Ile Ile Ile Ala Ser
 1 5 10 15
 Gly Asn Glu Ile Leu Glu Leu Val Val Asn Ala Met Met Gly Asp Asp
 20 25 30
 Pro Pro Glu Pro Pro Thr Pro Phe Glu Thr Pro Ser Leu His Asp Leu
 35 40 45

Tyr Asp Leu Glu Val Asp Val Pro Glu Asp Asp Pro Asn Glu Glu Ala
 50 55 60
 Val Asn Asp Phe Phe Ser Asp Ala Ala Leu Leu Ala Ala Glu Glu Ala
 65 70 75 80
 Ser Ser Ser Ser Ser Asp Ser Asp Ser Ser Leu His Thr Pro Arg Pro
 85 90 95
 Gly Arg Gly Glu Lys Lys Ile Pro Glu Leu Lys Gly Glu Glu Met Asp
 100 105 110
 Leu Arg Cys Tyr Glu Glu Cys Leu Pro Pro Ser Asp Asp Glu Asp Glu
 115 120 125
 Gln Ala Ile Gln Asn Ala Ala Ser Gln Gly Val Gln Ala Ala Ser Glu
 130 135 140
 Ser Phe Ala Leu Asp Cys Pro Pro Leu Pro Gly His Gly Cys Pro Val
 145 150 155 160
 Ser Asp Ala Asp Asp Glu Thr Pro Thr Thr Lys Ser Thr Ser Ser Pro
 165 170 175
 Pro Glu Ile Gly Thr Ser Pro Pro Glu Asn Ile Val Arg Pro Val Pro
 180 185 190
 Val Arg Ala Thr Gly Arg Arg Ala Ala Val Glu Cys Leu Asp Asp Leu
 195 200 205
 Leu Gln Gly Gly Val Glu Pro Leu Asp Leu Cys Thr Arg Lys Arg Pro
 210 215 220
 Arg His
 225

<210> 4
 <211> 186
 <212> PRT
 <213> chimpanzee C68 adenovirus protein
 <400> 4

Met Glu Ile Trp Thr Val Leu Glu Asp Phe His Lys Thr Arg Gln Leu
 1 5 10 15
 Leu Glu Asn Ala Ser Asn Gly Val Ser Tyr Leu Trp Arg Phe Cys Phe
 20 25 30
 Gly Gly Asp Leu Ala Arg Leu Val Tyr Arg Ala Lys Gln Asp Tyr Ser
 35 40 45
 Glu Gln Phe Glu Val Ile Leu Arg Glu Cys Ser Gly Leu Phe Asp Ala
 50 55 60
 Leu Asn Leu Gly His Gln Ser His Phe Asn Gln Arg Ile Ser Arg Ala
 65 70 75 80
 Leu Asp Phe Thr Thr Pro Gly Arg Thr Thr Ala Ala Val Ala Phe Phe
 85 90 95

Ala Phe Ile Leu Asp Lys Trp Ser Gln Glu Thr His Phe Ser Arg Asp
 100 105 110
 Tyr Gln Leu Asp Phe Leu Ala Val Ala Leu Trp Arg Thr Trp Lys Cys
 115 120 125
 Gln Arg Leu Asn Ala Ile Ser Gly Tyr Leu Pro Val Gln Pro Leu Asp
 130 135 140
 Thr Leu Arg Ile Leu Asn Leu Gln Glu Ser Pro Arg Ala Arg Gln Arg
 145 150 155 160
 Arg Gln Gln Gln Gln Gln Glu Glu Asp Gln Glu Glu Asn Pro Arg Ala
 165 170 175
 Gly Leu Asp Pro Pro Ala Glu Glu Glu Glu
 180 185

<210> 5
 <211> 498
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 5

Met Glu Ser Arg Asn Pro Phe Gln Gln Gly Leu Pro Ala Gly Phe Leu
 1 5 10 15
 Ser Ser Ser Phe Val Glu Asn Met Glu Val Pro Ala Pro Glu Cys Asn
 20 25 30
 Leu Arg Leu Leu Ala Gly Thr Ala Ala Arg His Ser Glu Asp Pro Glu
 35 40 45
 Ser Pro Gly Glu Ser Gln Gly Thr Pro Thr Ser Pro Ala Ala Ala Ala
 50 55 60
 Gly Gly Gly Ser Arg Arg Glu Pro Glu Ser Arg Pro Gly Pro Ser Gly
 65 70 75 80
 Gly Gly Gly Gly Val Ala Asp Leu Phe Pro Glu Leu Arg Arg Val Leu
 85 90 95
 Thr Arg Ser Ser Ser Gly Arg Glu Arg Gly Ile Lys Arg Glu Arg His
 100 105 110
 Asp Glu Thr Asn His Arg Thr Glu Leu Thr Val Gly Leu Met Ser Arg
 115 120 125
 Lys Arg Pro Glu Thr Val Trp Trp His Glu Val Gln Ser Thr Gly Thr
 130 135 140
 Asp Glu Val Ser Val Met His Glu Arg Phe Ser Leu Glu Gln Val Lys
 145 150 155 160
 Thr Cys Trp Leu Glu Pro Glu Asp Asp Trp Glu Val Ala Ile Arg Asn
 165 170 175
 Tyr Ala Lys Leu Ala Leu Arg Pro Asp Lys Lys Tyr Lys Ile Thr Lys
 180 185 190

Leu Ile Asn Ile Arg Asn Ala Cys Tyr Ile Ser Gly Asn Gly Ala Glu
 195 200 205
 Val Glu Ile Cys Leu Gln Glu Arg Val Ala Phe Arg Cys Cys Met Met
 210 215 220
 Asn Met Tyr Pro Gly Val Val Gly Met Asp Gly Val Thr Phe Met Asn
 225 230 235 240
 Met Arg Phe Arg Gly Asp Gly Tyr Asn Gly Thr Val Phe Met Ala Asn
 245 250 255
 Thr Lys Leu Thr Val His Gly Cys Ser Phe Phe Gly Phe Asn Asn Thr
 260 265 270
 Cys Ile Glu Ala Trp Gly Gln Val Gly Val Arg Gly Cys Ser Phe Ser
 275 280 285
 Ala Asn Trp Met Gly Val Val Gly Arg Thr Lys Ser Met Leu Ser Val
 290 295 300
 Lys Lys Cys Leu Phe Glu Arg Cys His Leu Gly Val Met Ser Glu Gly
 305 310 315 320
 Glu Ala Arg Ile Arg His Cys Ala Ser Thr Glu Thr Gly Cys Phe Val
 325 330 335
 Leu Cys Lys Gly Asn Ala Lys Ile Lys His Asn Met Ile Cys Gly Ala
 340 345 350
 Ser Asp Glu Arg Gly Tyr Gln Met Leu Thr Cys Ala Gly Gly Asn Ser
 355 360 365
 His Met Leu Ala Thr Val His Val Ala Ser His Ala Arg Lys Pro Trp
 370 375 380
 Pro Glu Phe Glu His Asn Val Met Thr Arg Cys Asn Met His Leu Gly
 385 390 395 400
 Ser Arg Arg Gly Met Phe Met Pro Tyr Gln Cys Asn Leu Asn Tyr Val
 405 410 415
 Lys Val Leu Leu Glu Pro Asp Ala Met Ser Arg Val Ser Leu Thr Gly
 420 425 430
 Val Phe Asp Met Asn Val Glu Val Trp Lys Ile Leu Arg Tyr Asp Glu
 435 440 445
 Ser Lys Thr Arg Cys Arg Ala Cys Glu Cys Gly Gly Lys His Ala Arg
 450 455 460
 Phe Gln Pro Val Cys Val Asp Val Thr Glu Asp Leu Arg Pro Asp His
 465 470 475 480
 Leu Val Leu Pro Cys Thr Gly Thr Glu Phe Gly Ser Ser Gly Glu Glu
 485 490 495
 Ser Asp

<210> 6
 <211> 169
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 6

```

Met Glu Ser Arg Asn Pro Phe Gln Gln Gly Leu Pro Ala Gly Phe Leu
1           5           10           15

Ser Ser Ser Phe Val Glu Asn Met Glu Val Pro Ala Pro Glu Cys Asn
          20           25           30

Leu Arg Leu Leu Ala Gly Thr Ala Ala Arg His Ser Glu Asp Pro Glu
          35           40           45

Ser Pro Gly Glu Ser Gln Gly Thr Pro Thr Ser Pro Ala Ala Ala Ala
          50           55           60

Gly Gly Gly Ser Arg Arg Glu Pro Glu Ser Arg Pro Gly Pro Ser Gly
65           70           75           80

Gly Gly Gly Gly Val Ala Asp Leu Phe Pro Glu Leu Arg Arg Val Leu
          85           90           95

Thr Arg Val Ser Leu Thr Gly Val Phe Asp Met Asn Val Glu Val Trp
          100          105          110

Lys Ile Leu Arg Tyr Asp Glu Ser Lys Thr Arg Cys Arg Ala Cys Glu
          115          120          125

Cys Gly Gly Lys His Ala Arg Phe Gln Pro Val Cys Val Asp Val Thr
          130          135          140

Glu Asp Leu Arg Pro Asp His Leu Val Leu Pro Cys Thr Gly Thr Glu
145          150          155          160

Phe Gly Ser Ser Gly Glu Glu Ser Asp
          165

```

<210> 7
 <211> 93
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 7

```

Met Glu Ser Arg Asn Pro Phe Gln Gln Gly Leu Pro Ala Gly Phe Leu
1           5           10           15

Ser Ser Ser Phe Val Glu Asn Met Glu Val Pro Ala Pro Glu Cys Asn
          20           25           30

Leu Arg Leu Leu Ala Gly Thr Ala Ala Arg His Ser Glu Asp Pro Glu
          35           40           45

Ser Pro Gly Glu Ser Gln Gly Thr Pro Thr Ser Pro Ala Ala Ala Ala
          50           55           60

```

Gly Gly Gly Ser Arg Arg Glu Pro Glu Ser Arg Pro Gly Pro Ser Gly
 65 70 75 80

Gly Gly Gly Gly Val Ala Asp Leu Pro Cys Val Trp Met
 85 90

<210> 8
 <211> 142
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 8

Met Ser Gly Ser Gly Ser Phe Glu Gly Gly Val Phe Ser Pro Tyr Leu
 1 5 10 15

Thr Gly Arg Leu Pro Ser Trp Ala Gly Val Arg Gln Asn Val Met Gly
 20 25 30

Ser Thr Val Asp Gly Arg Pro Val Gln Pro Ala Asn Ser Ser Thr Leu
 35 40 45

Thr Tyr Ala Thr Leu Ser Ser Ser Ser Leu Asp Ala Ala Ala Ala
 50 55 60

Ala Ala Ala Ser Ala Ala Ser Ala Val Arg Gly Met Ala Met Gly Ala
 65 70 75 80

Gly Tyr Tyr Gly Thr Leu Val Ala Asn Ser Ser Ser Thr Asn Asn Pro
 85 90 95

Ala Ser Leu Asn Glu Glu Lys Leu Leu Leu Leu Met Ala Gln Leu Glu
 100 105 110

Ala Leu Thr Gln Arg Leu Gly Glu Leu Thr Gln Gln Val Ala Gln Leu
 115 120 125

Gln Glu Gln Thr Arg Ala Ala Val Ala Thr Val Lys Ser Lys
 130 135 140

<210> 9
 <211> 448
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 9

Met Glu Thr Lys Gly Arg Arg Ser Gly Ala Val Phe Asp Gln Pro Asp
 1 5 10 15

Glu Pro Glu Ala His Pro Arg Lys Arg Pro Ala Arg Arg Ala Pro Leu
 20 25 30

His Arg Asp Gly Asp His Pro Asp Ala Asp Ala Ala Thr Leu Glu Gly
 35 40 45

Pro Asp Pro Gly Cys Ala Gly Arg Pro Ser Ser Gly Ala Ile Leu Pro
 50 55 60

Gln Pro Ser Gln Pro Ala Lys Arg Gly Gly Leu Leu Asp Arg Asp Ala
 65 70 75 80
 Val Glu His Ile Thr Glu Leu Trp Asp Arg Leu Glu Leu Leu Gln Gln
 85 90 95
 Thr Leu Ser Lys Met Pro Met Ala Asp Gly Leu Lys Pro Leu Lys Asn
 100 105 110
 Phe Ala Ser Leu Gln Glu Leu Leu Ser Leu Gly Gly Glu Arg Leu Leu
 115 120 125
 Ala Glu Leu Val Arg Glu Asn Met His Val Arg Glu Met Met Asn Glu
 130 135 140
 Val Ala Pro Leu Leu Arg Glu Asp Gly Ser Cys Leu Ser Leu Asn Tyr
 145 150 155 160
 His Leu Gln Pro Val Ile Gly Val Ile Tyr Gly Pro Thr Gly Cys Gly
 165 170 175
 Lys Ser Gln Leu Leu Arg Asn Leu Leu Ser Ala Gln Leu Ile Ser Pro
 180 185 190
 Ala Pro Glu Thr Val Phe Phe Ile Ala Pro Gln Val Asp Met Ile Pro
 195 200 205
 Pro Ser Glu Leu Lys Ala Trp Glu Met Gln Ile Cys Glu Gly Asn Tyr
 210 215 220
 Ala Pro Gly Ile Glu Gly Thr Phe Val Pro Gln Ser Gly Thr Leu Arg
 225 230 235 240
 Pro Lys Phe Ile Lys Met Ala Tyr Asp Asp Leu Thr Gln Asp His Asn
 245 250 255
 Tyr Asp Val Ser Asp Pro Arg Asn Val Phe Ala Gln Ala Ala Ala His
 260 265 270
 Gly Pro Ile Ala Ile Ile Met Asp Glu Cys Met Glu Asn Leu Gly Gly
 275 280 285
 His Lys Gly Val Ala Lys Phe Phe His Ala Phe Pro Ser Lys Leu His
 290 295 300
 Asp Lys Phe Pro Lys Cys Thr Gly Tyr Thr Val Leu Val Val Leu His
 305 310 315 320
 Asn Met Asn Pro Arg Arg Asp Leu Gly Gly Asn Ile Ala Asn Leu Lys
 325 330 335
 Ile Gln Ala Lys Met His Leu Ile Ser Pro Arg Met His Pro Ser Gln
 340 345 350
 Leu Asn Arg Phe Val Asn Thr Tyr Thr Lys Gly Leu Pro Val Ala Ile
 355 360 365
 Ser Leu Leu Leu Lys Asp Ile Val Gln His His Ala Leu Arg Pro Cys
 370 375 380
 Tyr Asp Trp Val Ile Tyr Asn Thr Thr Pro Glu His Glu Ala Leu Gln
 385 390 395 400

Trp Ser Tyr Leu His Pro Arg Asp Gly Leu Met Pro Met Tyr Leu Asn
405 410 415

Ile Gln Ala His Leu Tyr Arg Val Leu Glu Lys Ile His Arg Val Leu
420 425 430

Asn Asp Arg Asp Arg Trp Ser Arg Ala Tyr Arg Ala Arg Lys Ile Lys
435 440 445

```
<210> 10
<211> 200
<212> PRT
<213> chimpanzee C68 adenovirus protein
```

```
<220>
<221> MISC_FEATURE
<222> (137)..(137)
<223> xaa can be any amino acid
```

```
<220>
<221> MISC_FEATURE
<222> (155)..(155)
<223> xaa can be any amino acid
```

<400> 10

Met Arg Ala Asp Gly Glu Glu Leu Asp Leu Leu Pro Pro Ile Gly Gly
1 5 10 15

Met Ala Val Asp Val Met Glu Val Glu Met Pro Thr Ala Arg Arg Thr
20 25 30

Leu Val Leu Val Phe Ile Gln Ala Ala Thr Val Leu Ala Thr Leu His
35 40 45

Gly Met His Val Leu His Glu Leu Tyr Leu Ser Ser Phe Asp Glu Glu
50 55 60

Phe Gln Trp Glu Val Glu Ser Trp Arg Leu His Leu Val Leu Tyr Tyr
65 70 75 80

Val Val Val Val Gly Leu Ala Leu Phe Cys Leu Asp Gly Gly His Ala
85 90 95

Asp Glu Pro Ala Arg Glu Ala Gly Pro Asp Leu Gly Ala Ser Gly Ser
100 105 110

Glu Ser Glu Asp Glu Gly Ala Gln Ala Gly Ala Val Gln Gly Pro Glu
115 120 125

Thr Leu Arg Ser Gln Val Ser Gly Xaa Arg Arg Arg Ala Val Asp Leu
130 135 140

Gln Glu Phe Phe Gln Gly Ala Arg Glu Val Xaa Met Val Leu Asp Leu
145 150 155 160

His Arg Ala Ile Gly Gly Glu Leu His Gly Leu Gln Gly Pro Val Pro
165 170 175

Leu Gly Cys Asp His Arg Pro Pro Phe Leu Leu Gly Arg Leu Gly Arg
180 185 190

Arg Gly Arg Cys Leu Phe His Gly
 195 200

<210> 11
 <211> 391
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 11

Met His Pro Val Leu Arg Gln Met Arg Pro His His Pro Pro Pro Gln
 1 5 10 15
 Gln Gln Pro Pro Pro Gln Pro Ala Leu Leu Pro Pro Pro Gln Gln Gln
 20 25 30
 Leu Pro Ala Thr Thr Ala Ala Ala Val Ser Gly Ala Gly Gln Ser
 35 40 45
 Tyr Asp His Gln Leu Ala Leu Glu Glu Gly Glu Gly Leu Ala Arg Leu
 50 55 60
 Gly Ala Ser Ser Pro Glu Arg His Pro Arg Val Gln Met Lys Arg Asp
 65 70 75 80
 Ala Arg Glu Ala Tyr Val Pro Lys Gln Asn Leu Phe Arg Asp Arg Ser
 85 90 95
 Gly Glu Glu Pro Glu Glu Met Arg Ala Ala Arg Phe His Ala Gly Arg
 100 105 110
 Glu Leu Arg Arg Gly Leu Asp Arg Lys Arg Val Leu Arg Asp Glu Asp
 115 120 125
 Phe Glu Ala Asp Glu Leu Thr Gly Ile Ser Pro Ala Arg Ala His Val
 130 135 140
 Ala Ala Ala Asn Leu Val Thr Ala Tyr Glu Gln Thr Val Lys Glu Glu
 145 150 155 160
 Ser Asn Phe Gln Lys Ser Phe Asn Asn His Val Arg Thr Leu Ile Ala
 165 170 175
 Arg Glu Glu Val Thr Leu Gly Leu Met His Leu Trp Asp Leu Leu Glu
 180 185 190
 Ala Ile Val Gln Asn Pro Thr Ser Lys Pro Leu Thr Ala Gln Leu Phe
 195 200 205
 Leu Val Val Gln His Ser Arg Asp Asn Glu Ala Phe Arg Glu Ala Leu
 210 215 220
 Leu Asn Ile Thr Glu Pro Glu Gly Arg Trp Leu Leu Asp Leu Val Asn
 225 230 235 240
 Ile Leu Gln Ser Ile Val Val Gln Glu Arg Gly Leu Pro Leu Ser Glu
 245 250 255
 Lys Leu Ala Ala Ile Asn Phe Ser Val Leu Ser Leu Gly Lys Tyr Tyr
 260 265 270

Ala	Arg	Lys 275	Ile	Tyr	Lys	Thr	Pro 280	Tyr	Val	Pro	Ile	Asp 285	Lys	Glu	Val
Lys	Ile	Asp 290	Gly	Phe	Tyr	Met 295	Arg	Met	Thr	Leu	Lys 300	Val	Leu	Thr	Leu
Ser 305	Asp	Asp	Leu	Gly	Val 310	Tyr	Arg	Asn	Asp	Arg 315	Met	His	Arg	Ala	Val 320
Ser	Ala	Ser	Arg	Arg 325	Arg	Glu	Leu	Ser	Asp 330	Gln	Glu	Leu	Met	His 335	Ser
Leu	Gln	Arg	Ala 340	Leu	Thr	Gly	Ala	Gly 345	Thr	Glu	Gly	Glu	Ser 350	Tyr	Phe
Asp	Met	Gly 355	Ala	Asp	Leu	His	Trp 360	Gln	Pro	Ser	Arg	Arg 365	Ala	Leu	Glu
Ala	Ala 370	Ala	Gly	Pro	Tyr	Val 375	Glu	Glu	Val	Asp	Asp 380	Glu	Val	Asp	Glu
Glu 385	Gly	Glu	Tyr	Leu	Glu 390	Asp									

```
<210> 12
<211> 534
<212> PRT
<213> chimpanzee C68 adenovirus protein
```

<400> 12

Met 1	Met	Arg	Arg	Ala 5	Tyr	Pro	Glu	Gly	Pro 10	Pro	Pro	Ser	Tyr	Glu 15	Ser
Val	Met	Gln	Gln 20	Ala	Met	Ala	Ala	Ala 25	Ala	Met	Gln	Pro	Pro 30	Leu	Glu
Ala	Pro	Tyr 35	Val	Pro	Pro	Arg	Tyr 40	Leu	Ala	Pro	Thr	Glu 45	Gly	Arg	Asn
Ser	Ile 50	Arg	Tyr	Ser	Glu	Leu 55	Ala	Pro	Leu	Tyr	Asp 60	Thr	Thr	Arg	Leu
Tyr 65	Leu	Val	Asp	Asn	Lys 70	Ser	Ala	Asp	Ile	Ala 75	Ser	Leu	Asn	Tyr	Gln 80
Asn	Asp	His	Ser	Asn 85	Phe	Leu	Thr	Thr	Val 90	Val	Gln	Asn	Asn	Asp 95	Phe
Thr	Pro	Thr	Glu 100	Ala	Ser	Thr	Gln	Thr 105	Ile	Asn	Phe	Asp	Glu 110	Arg	Ser
Arg	Trp	Gly 115	Gly	Gln	Leu	Lys	Thr 120	Ile	Met	His	Thr	Asn 125	Met	Pro	Asn
Val	Asn 130	Glu	Phe	Met	Tyr	Ser 135	Asn	Lys	Phe	Lys	Ala 140	Arg	Val	Met	Val
Ser 145	Arg	Lys	Thr	Pro	Asn 150	Gly	Val	Thr	Val	Thr 155	Glu	Asp	Tyr	Asp	Gly 160

Ser Gln Asp Glu Leu Lys Tyr Glu Trp Val Glu Phe Glu Leu Pro Glu
 165 170 175
 Gly Asn Phe Ser Val Thr Met Thr Ile Asp Leu Met Asn Asn Ala Ile
 180 185 190
 Ile Asp Asn Tyr Leu Ala Val Gly Arg Gln Asn Gly Val Leu Glu Ser
 195 200 205
 Asp Ile Gly Val Lys Phe Asp Thr Arg Asn Phe Arg Leu Gly Trp Asp
 210 215 220
 Pro Val Thr Glu Leu Val Met Pro Gly Val Tyr Thr Asn Glu Ala Phe
 225 230 235 240
 His Pro Asp Ile Val Leu Leu Pro Gly Cys Gly Val Asp Phe Thr Glu
 245 250 255
 Ser Arg Leu Ser Asn Leu Leu Gly Ile Arg Lys Arg Gln Pro Phe Gln
 260 265 270
 Glu Gly Phe Gln Ile Met Tyr Glu Asp Leu Glu Gly Gly Asn Ile Pro
 275 280 285
 Ala Leu Leu Asp Val Asp Ala Tyr Glu Lys Ser Lys Glu Asp Ala Ala
 290 295 300
 Ala Glu Ala Thr Ala Ala Val Ala Thr Ala Ser Thr Glu Val Arg Gly
 305 310 315 320
 Asp Asn Phe Ala Ser Ala Ala Ala Val Ala Ala Ala Glu Ala Ala Glu
 325 330 335
 Thr Glu Ser Lys Ile Val Ile Gln Pro Val Glu Lys Asp Ser Lys Asn
 340 345 350
 Arg Ser Tyr Asn Val Leu Pro Asp Lys Ile Asn Thr Ala Tyr Arg Ser
 355 360 365
 Trp Tyr Leu Ala Tyr Asn Tyr Gly Asp Pro Glu Lys Gly Val Arg Ser
 370 375 380
 Trp Thr Leu Leu Thr Thr Ser Asp Val Thr Cys Gly Val Glu Gln Val
 385 390 395 400
 Tyr Trp Ser Leu Pro Asp Met Met Gln Asp Pro Val Thr Phe Arg Ser
 405 410 415
 Thr Arg Gln Val Ser Asn Tyr Pro Val Val Gly Ala Glu Leu Leu Pro
 420 425 430
 Val Tyr Ser Lys Ser Phe Phe Asn Glu Gln Ala Val Tyr Ser Gln Gln
 435 440 445
 Leu Arg Ala Phe Thr Ser Leu Thr His Val Phe Asn Arg Phe Pro Glu
 450 455 460
 Asn Gln Ile Leu Val Arg Pro Pro Ala Pro Thr Ile Thr Thr Val Ser
 465 470 475 480
 Glu Asn Val Pro Ala Leu Thr Asp His Gly Thr Leu Pro Leu Arg Ser
 485 490 495

Ser Ile Arg Gly Val Gln Arg Val Thr Val Thr Asp Ala Arg Arg Arg
 500 505 510

Thr Cys Pro Tyr Val Tyr Lys Ala Leu Gly Ile Val Ala Pro Arg Val
 515 520 525

Leu Ser Ser Arg Thr Phe
 530

<210> 13
 <211> 201
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<220>
 <221> MISC_FEATURE
 <222> (70)..(70)
 <223> Xaa can be any amino acid

<400> 13

Met Ser Ile Leu Ile Ser Pro Ser Asn Asn Thr Gly Trp Gly Leu Arg
 1 5 10 15

Ala Pro Ser Lys Met Tyr Gly Gly Ala Arg Gln Arg Ser Thr Gln His
 20 25 30

Pro Val Arg Val Arg Gly His Phe Arg Ala Pro Trp Gly Ala Leu Lys
 35 40 45

Gly Arg Val Arg Ser Arg Thr Thr Val Asp Asp Val Ile Asp Gln Val
 50 55 60

Val Ala Asp Ala Arg Xaa Tyr Thr Pro Ala Ala Ala Pro Val Ser Thr
 65 70 75 80

Val Asp Ala Val Ile Asp Ser Val Val Ala Asp Ala Arg Arg Tyr Ala
 85 90 95

Arg Ala Lys Ser Arg Arg Arg Arg Ile Ala Arg Arg His Arg Ser Thr
 100 105 110

Pro Ala Met Arg Ala Ala Arg Ser Leu Val Ala Gln Gly Gln Ala His
 115 120 125

Gly Thr Gln Gly His Val Gln Gly Gly Gln Thr Arg Gly Phe Arg Arg
 130 135 140

Gln Arg Arg Gln Asp Pro Glu Thr Arg Gly His Gly Gly Gly Ser Gly
 145 150 155 160

His Arg Gln His Val Pro Pro Ala Ala Arg Glu Arg Val Leu Gly Ala
 165 170 175

Arg Arg Arg His Arg Cys Ala Arg Ala Arg Ala His Pro Pro Pro Ser
 180 185 190

His Leu Lys Met Phe Thr Ser Arg Cys
 195 200

<210> 14
 <211> 356
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<220>
 <221> MISC_FEATURE
 <222> (111)..(111)
 <223> Xaa can be any amino acid

<220>
 <221> MISC_FEATURE
 <222> (183)..(183)
 <223> Xaa can be any amino acid

<220>
 <221> MISC_FEATURE
 <222> (212)..(212)
 <223> Xaa can be any amino acid

<220>
 <221> MISC_FEATURE
 <222> (220)..(220)
 <223> Xaa can be any amino acid

<400> 14

Met	Ser	Lys	Arg	Lys	Phe	Lys	Glu	Glu	Met	Leu	Gln	Val	Ile	Ala	Pro	1	5	10	15
Glu	Ile	Tyr	Gly	Pro	Ala	Val	Val	Lys	Glu	Glu	Arg	Lys	Pro	Arg	Lys	20	25	30	
Ile	Lys	Arg	Val	Lys	Lys	Asp	Lys	Lys	Glu	Glu	Glu	Ser	Asp	Val	Asp	35	40	45	
Gly	Leu	Val	Glu	Phe	Val	Arg	Glu	Phe	Ala	Pro	Arg	Arg	Arg	Val	Gln	50	55	60	
Trp	Arg	Gly	Arg	Lys	Val	Gln	Pro	Val	Leu	Arg	Pro	Gly	Thr	Thr	Val	65	70	75	
Val	Phe	Thr	Pro	Gly	Glu	Arg	Ser	Gly	Thr	Ala	Ser	Lys	Arg	Ser	Tyr	85	90	95	
Asp	Glu	Val	Tyr	Gly	Asp	Asp	Asp	Ile	Leu	Glu	Gln	Ala	Ala	Xaa	Arg	100	105	110	
Leu	Gly	Glu	Phe	Ala	Tyr	Gly	Lys	Arg	Ser	Arg	Ser	Ala	Pro	Lys	Glu	115	120	125	
Glu	Ala	Val	Ser	Ile	Pro	Leu	Asp	His	Gly	Asn	Pro	Thr	Pro	Ser	Leu	130	135	140	
Lys	Pro	Val	Thr	Leu	Gln	Gln	Val	Leu	Pro	Thr	Ala	Ala	Pro	Arg	Arg	145	150	155	
Gly	Phe	Lys	Arg	Glu	Gly	Glu	Asp	Leu	Tyr	Pro	Thr	Met	Gln	Leu	Met	165	170	175	
Val	Pro	Lys	Arg	Gln	Lys	Xaa	Glu	Asp	Val	Leu	Glu	Thr	Met	Lys	Val	180	185	190	

Asp Pro Asp Val Gln Pro Glu Val Lys Val Arg Pro Ile Lys Gln Val
 195 200 205
 Ala Pro Gly Xaa Gly Val Gln Thr Val Asp Ile Xaa Ile Pro Thr Glu
 210 215 220
 Pro Met Glu Thr Gln Thr Glu Pro Met Ile Lys Pro Ser Thr Ser Thr
 225 230 235 240
 Met Glu Val Gln Thr Asp Pro Trp Met Pro Ser Ala Pro Ser Arg Arg
 245 250 255
 Pro Arg Arg Lys Tyr Gly Ala Ala Ser Leu Leu Met Pro Asn Tyr Ala
 260 265 270
 Leu His Pro Ser Ile Ile Pro Thr Pro Gly Tyr Arg Gly Thr Arg Phe
 275 280 285
 Tyr Arg Gly His Thr Ser Ser Arg Arg Arg Lys Thr Thr Thr Arg Arg
 290 295 300
 Ser Pro Ser Pro His Arg Arg Cys Asn His Pro Cys Arg Pro Gly Ala
 305 310 315 320
 Glu Ser Val Pro Pro Arg Pro Arg Thr Ser Asp Pro Ala Ala Arg Ala
 325 330 335
 Leu Pro Pro Glu His Arg His Leu Asn Phe Arg Gln Leu Cys Arg Ser
 340 345 350
 Met Ala Leu Thr
 355

<210> 15
 <211> 257
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<220>
 <221> MISC_FEATURE
 <222> (202)..(202)
 <223> Xaa can be any amino acid

<220>
 <221> MISC_FEATURE
 <222> (210)..(210)
 <223> Xaa can be any amino acid

<220>
 <221> MISC_FEATURE
 <222> (256)..(256)
 <223> Xaa can be any amino acid

<400> 15

Met Asp Ser Asp Ala Pro Gly Pro Val Met Cys Phe Arg Arg Gln Met
 1 5 10 15
 Glu Asp Ile Asn Phe Ser Ser Leu Ala Pro Arg His Gly Thr Arg Pro
 20 25 30

Phe Met Gly Thr Trp Ser Asp Ile Gly Thr Ser Gln Leu Asn Gly Gly
 35 40 45
 Ala Phe Asn Trp Ser Ser Leu Trp Ser Gly Leu Lys Asn Phe Gly Ser
 50 55 60
 Thr Leu Lys Thr Tyr Gly Ser Lys Ala Trp Asn Ser Thr Thr Gly Gln
 65 70 75 80
 Ala Leu Arg Asp Lys Leu Lys Glu Gln Asn Phe Gln Gln Lys Val Val
 85 90 95
 Asp Gly Leu Ala Ser Gly Ile Asn Gly Val Val Asp Leu Ala Asn Gln
 100 105 110
 Ala Val Gln Arg Gln Ile Asn Ser Arg Leu Asp Pro Val Pro Pro Ala
 115 120 125
 Gly Ser Val Glu Met Pro Gln Val Glu Glu Glu Leu Pro Pro Leu Asp
 130 135 140
 Lys Arg Gly Glu Lys Arg Pro Arg Pro Asp Ala Glu Glu Thr Leu Leu
 145 150 155 160
 Thr His Thr Asp Glu Pro Pro Pro Tyr Glu Glu Ala Val Lys Leu Gly
 165 170 175
 Leu Pro Thr Thr Arg Pro Ile Ala Pro Leu Ala Thr Gly Val Leu Lys
 180 185 190
 Pro Glu Lys Pro Ala Thr Leu Asp Leu Xaa Pro Pro Gln Pro Ser Arg
 195 200 205
 Pro Xaa Thr Val Ala Lys Pro Leu Pro Pro Val Ala Val Ala Arg Ala
 210 215 220
 Arg Pro Gly Gly Thr Ala Arg Pro His Ala Asn Trp Gln Ser Thr Leu
 225 230 235 240
 Asn Ser Ile Val Gly Leu Gly Val Gln Ser Val Lys Arg Arg Arg Xaa
 245 250 255

Tyr

<210> 16
 <211> 933
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<220>
 <221> MISC_FEATURE
 <222> (826)..(826)
 <223> Xaa can be any amino acid

<220>
 <221> MISC_FEATURE
 <222> (922)..(922)
 <223> Xaa can be any amino acid

<400> 16

```

Met Ala Thr Pro Ser Met Leu Pro Gln Trp Ala Tyr Met His Ile Ala
1      5      10      15

Gly Gln Asp Ala Ser Glu Tyr Leu Ser Pro Gly Leu Val Gln Phe Ala
20     25     30

Arg Ala Thr Asp Thr Tyr Phe Ser Leu Gly Asn Lys Phe Arg Asn Pro
35     40     45

Thr Val Ala Pro Thr His Asp Val Thr Thr Asp Arg Ser Gln Arg Leu
50     55     60

Thr Leu Arg Phe Val Pro Val Asp Arg Glu Asp Asn Thr Tyr Ser Tyr
65     70     75     80

Lys Val Arg Tyr Thr Leu Ala Val Gly Asp Asn Arg Val Leu Asp Met
85     90     95

Ala Ser Thr Tyr Phe Asp Ile Arg Gly Val Leu Asp Arg Gly Pro Ser
100    105    110

Phe Lys Pro Tyr Ser Gly Thr Ala Tyr Asn Ser Leu Ala Pro Lys Gly
115    120    125

Ala Pro Asn Thr Cys Gln Trp Thr Tyr Lys Ala Asp Gly Glu Thr Ala
130    135    140

Thr Glu Lys Thr Tyr Thr Tyr Gly Asn Ala Pro Val Gln Gly Ile Asn
145    150    155    160

Ile Thr Lys Asp Gly Ile Gln Leu Gly Thr Asp Thr Asp Asp Gln Pro
165    170    175

Ile Tyr Ala Asp Lys Thr Tyr Gln Pro Glu Pro Gln Val Gly Asp Ala
180    185    190

Glu Trp His Asp Ile Thr Gly Thr Asp Glu Lys Tyr Gly Gly Arg Ala
195    200    205

Leu Lys Pro Asp Thr Lys Met Lys Pro Cys Tyr Gly Ser Phe Ala Lys
210    215    220

Pro Thr Asn Lys Glu Gly Gly Gln Ala Asn Val Lys Thr Gly Thr Gly
225    230    235    240

Thr Thr Lys Glu Tyr Asp Ile Asp Met Ala Phe Phe Asp Asn Arg Ser
245    250    255

Ala Ala Ala Ala Gly Leu Ala Pro Glu Ile Val Leu Tyr Thr Glu Asn
260    265    270

Val Asp Leu Glu Thr Pro Asp Thr His Ile Val Tyr Lys Ala Gly Thr
275    280    285

Asp Asp Ser Ser Ser Ser Ile Asn Leu Gly Gln Gln Ala Met Pro Asn
290    295    300

Arg Pro Asn Tyr Ile Gly Phe Arg Asp Asn Phe Ile Gly Leu Met Tyr
305    310    315    320

```

Tyr Asn Ser Thr Gly Asn Met Gly Val Leu Ala Gly Gln Ala Ser Gln
 325 330 335
 Leu Asn Ala Val Val Asp Leu Gln Asp Arg Asn Thr Glu Leu Ser Tyr
 340 345 350
 Gln Leu Leu Leu Asp Ser Leu Gly Asp Arg Thr Arg Tyr Phe Ser Met
 355 360 365
 Trp Asn Gln Ala Val Asp Ser Tyr Asp Pro Asp Val Arg Ile Ile Glu
 370 375 380
 Asn His Gly Val Glu Asp Glu Leu Pro Asn Tyr Cys Phe Pro Leu Asp
 385 390 395 400
 Ala Val Gly Arg Thr Asp Thr Tyr Gln Gly Ile Lys Ala Asn Gly Thr
 405 410 415
 Asp Gln Thr Thr Trp Thr Lys Asp Asp Ser Val Asn Asp Ala Asn Glu
 420 425 430
 Ile Gly Lys Gly Asn Pro Phe Ala Met Glu Ile Asn Ile Gln Ala Asn
 435 440 445
 Leu Trp Arg Asn Phe Leu Tyr Ala Asn Val Ala Leu Tyr Leu Pro Asp
 450 455 460
 Ser Tyr Lys Tyr Thr Pro Ala Asn Val Thr Leu Pro Thr Asn Thr Asn
 465 470 475 480
 Thr Tyr Asp Tyr Met Asn Gly Arg Val Val Ala Pro Ser Leu Val Asp
 485 490 495
 Ser Tyr Ile Asn Ile Gly Ala Arg Trp Ser Leu Asp Pro Met Asp Asn
 500 505 510
 Val Asn Pro Phe Asn His His Arg Asn Ala Gly Leu Arg Tyr Arg Ser
 515 520 525
 Met Leu Leu Gly Asn Gly Arg Tyr Val Pro Phe His Ile Gln Val Pro
 530 535 540
 Gln Lys Phe Phe Ala Ile Lys Ser Leu Leu Leu Leu Pro Gly Ser Tyr
 545 550 555 560
 Thr Tyr Glu Trp Asn Phe Arg Lys Asp Val Asn Met Ile Leu Gln Ser
 565 570 575
 Ser Leu Gly Asn Asp Leu Arg Thr Asp Gly Ala Ser Ile Ser Phe Thr
 580 585 590
 Ser Ile Asn Leu Tyr Ala Thr Phe Phe Pro Met Ala His Asn Thr Ala
 595 600 605
 Ser Thr Leu Glu Ala Met Leu Arg Asn Asp Thr Asn Asp Gln Ser Phe
 610 615 620
 Asn Asp Tyr Leu Ser Ala Ala Asn Met Leu Tyr Pro Ile Pro Ala Asn
 625 630 635 640
 Ala Thr Asn Val Pro Ile Ser Ile Pro Ser Arg Asn Trp Ala Ala Phe
 645 650 655

Arg Gly Trp Ser Phe Thr Arg Leu Lys Thr Lys Glu Thr Pro Ser Leu
 660 665 670
 Gly Ser Gly Phe Asp Pro Tyr Phe Val Tyr Ser Gly Ser Ile Pro Tyr
 675 680 685
 Leu Asp Gly Thr Phe Tyr Leu Asn His Thr Phe Lys Lys Val Ser Ile
 690 695 700
 Thr Phe Asp Ser Ser Val Ser Trp Pro Gly Asn Asp Arg Leu Leu Thr
 705 710 715 720
 Pro Asn Glu Phe Glu Ile Lys Arg Thr Val Asp Gly Glu Gly Tyr Asn
 725 730 735
 Val Ala Gln Cys Asn Met Thr Lys Asp Trp Phe Leu Val Gln Met Leu
 740 745 750
 Ala His Tyr Asn Ile Gly Tyr Gln Gly Phe Tyr Val Pro Glu Gly Tyr
 755 760 765
 Lys Asp Arg Met Tyr Ser Phe Phe Arg Asn Phe Gln Pro Met Ser Arg
 770 775 780
 Gln Val Val Asp Glu Val Asn Tyr Lys Asp Tyr Gln Ala Val Thr Leu
 785 790 795 800
 Ala Tyr Gln His Asn Asn Ser Gly Phe Val Gly Tyr Leu Ala Pro Thr
 805 810 815
 Met Arg Gln Gly Gln Pro Tyr Pro Ala Xaa Tyr Pro Tyr Pro Leu Ile
 820 825 830
 Gly Lys Ser Ala Val Thr Ser Val Thr Gln Lys Lys Phe Leu Cys Asp
 835 840 845
 Arg Val Met Trp Arg Ile Pro Phe Ser Ser Asn Phe Met Ser Met Gly
 850 855 860
 Ala Leu Thr Asp Leu Gly Gln Asn Met Leu Tyr Ala Asn Ser Ala His
 865 870 875 880
 Ala Leu Asp Met Asn Phe Glu Val Asp Pro Met Asp Glu Ser Thr Leu
 885 890 895
 Leu Tyr Val Val Phe Glu Val Phe Asp Val Val Arg Val His Gln Pro
 900 905 910
 His Arg Gly Val Ile Glu Ala Val Tyr Xaa Arg Thr Pro Phe Ser Ala
 915 920 925
 Gly Asn Ala Thr Thr
 930

<210> 17

<211> 513

<212> PRT

<213> chimpanzee C68 adenovirus protein

<220> <221> MISC FEATURE
 <222> (511) .. (511)
 <223> Xaa can be any amino acid

<400> 17

```

Met Ala Gly Arg Gly Gly Ser Gln Ser Glu Arg Arg Arg Glu Arg Thr
1          5          10          15

Pro Glu Arg Gly Arg Gly Ser Ala Ser His Pro Pro Ser Arg Gly Gly
          20          25          30

Glu Ser Pro Ser Pro Pro Pro Leu Pro Pro Lys Arg His Thr Tyr Arg
          35          40          45

Arg Val Ala Ser Asp Gln Glu Glu Glu Ile Val Val Val Ser Glu
          50          55          60

Asn Ser Arg Ser Pro Ser Pro Ser Pro Thr Ser Pro Pro Pro Leu Pro
65          70          75          80

Pro Lys Lys Lys Pro Arg Lys Thr Lys His Val Val Leu Gln Asp Val
          85          90          95

Ser Gln Asp Ser Glu Asp Glu Arg Gln Ala Glu Glu Glu Leu Ala Ala
          100          105          110

Val Gly Phe Ser Tyr Pro Pro Val Arg Ile Thr Glu Lys Asp Gly Lys
          115          120          125

Arg Ser Phe Glu Thr Leu Asp Glu Ser Asp Pro Leu Ala Ala Ala Ala
          130          135          140

Ser Ala Lys Met Met Val Lys Asn Pro Met Ser Leu Pro Ile Val Ser
          145          150          155          160

Ala Trp Glu Lys Gly Met Glu Ile Met Thr Met Leu Met Asp Arg Tyr
          165          170          175

Arg Val Glu Thr Asp Leu Lys Ala Asn Phe Gln Leu Met Pro Glu Gln
          180          185          190

Gly Glu Val Tyr Arg Arg Ile Cys His Leu Tyr Ile Asn Glu Glu His
          195          200          205

Arg Gly Ile Pro Leu Thr Phe Thr Ser Asn Lys Thr Leu Thr Thr Met
          210          215          220

Met Gly Arg Phe Leu Gln Gly Phe Val His Ala His Ser Gln Ile Ala
          225          230          235          240

His Lys Asn Trp Glu Cys Thr Gly Cys Ala Leu Trp Leu His Gly Cys
          245          250          255

Thr Glu Ala Glu Gly Lys Leu Arg Cys Leu His Gly Thr Thr Met Ile
          260          265          270

Gln Lys Glu His Met Ile Glu Met Asp Val Ala Ser Glu Asn Gly Gln
          275          280          285

Arg Ala Leu Lys Glu Asn Pro Asp Arg Ala Lys Ile Thr Gln Asn Arg
          290          295          300

```


Trp Gly Arg Ser Val Val Gln Leu Ala Asn Asn Asp Ala Arg Cys Cys
 305 310 315 320
 Val His Asp Ala Gly Cys Ala Thr Asn Gln Phe Ser Ser Lys Ser Cys
 325 330 335
 Gly Val Phe Phe Thr Glu Gly Ala Lys Ala Gln Gln Ala Phe Arg Gln
 340 345 350
 Leu Glu Ala Phe Met Lys Ala Met Tyr Pro Gly Met Asn Ala Asp Gln
 355 360 365
 Ala Gln Met Met Leu Ile Pro Leu His Cys Asp Cys Asn His Lys Pro
 370 375 380
 Gly Cys Val Pro Thr Met Gly Arg Gln Thr Cys Lys Met Thr Pro Phe
 385 390 395 400
 Gly Met Ala Asn Ala Glu Asp Leu Asp Val Glu Ser Ile Thr Asp Ala
 405 410 415
 Thr Val Leu Ala Ser Val Lys His Pro Ala Leu Met Val Phe Gln Cys
 420 425 430
 Cys Asn Pro Val Tyr Arg Asn Ser Arg Ala Gln Asn Ala Gly Pro Asn
 435 440 445
 Cys Asp Phe Lys Ile Ser Ala Pro Asp Leu Leu Gly Ala Leu Gln Leu
 450 455 460
 Thr Arg Lys Leu Trp Thr Asp Ser Phe Pro Asp Thr Pro Leu Pro Lys
 465 470 475 480
 Leu Leu Ile Pro Glu Phe Lys Trp Leu Ala Lys Tyr Gln Phe Arg Asn
 485 490 495
 Val Ser Leu Pro Ala Gly His Ala Glu Thr Arg Lys Asn Pro Xaa Asp
 500 505 510

Phe

<210> 18
 <211> 222
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 18

Met Pro Arg Gly Asn Lys Lys Leu Lys Val Glu Leu Pro Pro Val Glu
 1 5 10 15
 Asp Leu Glu Glu Asp Trp Glu Asn Ser Ser Gln Ala Glu Glu Glu
 20 25 30
 Met Glu Glu Asp Trp Asp Ser Thr Gln Ala Glu Glu Asp Ser Leu Gln
 35 40 45
 Asp Ser Leu Glu Glu Asp Glu Glu Glu Ala Glu Glu Val Glu Glu
 50 55 60

Ala Ala Ala Ala Arg Pro Ser Ser Ser Ala Gly Glu Lys Ala Ser Ser
 65 70 75 80
 Thr Asp Thr Ile Ser Ala Pro Gly Arg Gly Pro Ala Arg Pro His Ser
 85 90 95
 Arg Trp Asp Glu Thr Gly Arg Phe Pro Asn Pro Thr Thr Gln Thr Ala
 100 105 110
 Pro Thr Thr Ser Lys Lys Arg Gln Gln Gln Gln Lys Lys Thr Ser Arg
 115 120 125
 Lys Pro Ala Ala Arg Lys Ser Thr Ala Ala Ala Ala Gly Gly Leu Arg
 130 135 140
 Ile Ala Ala Asn Glu Pro Ala Gln Thr Arg Glu Leu Arg Asn Arg Ile
 145 150 155 160
 Phe Pro Thr Leu Tyr Ala Ile Phe Gln Gln Ser Arg Gly Gln Glu Gln
 165 170 175
 Glu Leu Lys Val Lys Asn Arg Ser Leu Arg Ser Leu Thr Arg Ser Cys
 180 185 190
 Leu Tyr His Lys Ser Glu Asp Gln Leu Gln Arg Thr Leu Glu Asp Ala
 195 200 205
 Glu Ala Leu Phe Asn Lys Tyr Cys Ala Leu Thr Leu Lys Glu
 210 215 220

<210> 19
 <211> 227
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 19

Met Ser Lys Glu Ile Pro Thr Pro Tyr Met Trp Ser Tyr Gln Pro Gln
 1 5 10 15
 Met Gly Leu Ala Ala Gly Ala Ala Gln Asp Tyr Ser Thr Arg Met Asn
 20 25 30
 Trp Leu Ser Ala Gly Pro Ala Met Ile Ser Arg Val Asn Asp Ile Arg
 35 40 45
 Ala His Arg Asn Gln Ile Leu Leu Glu Gln Ser Ala Leu Thr Ala Thr
 50 55 60
 Pro Arg Asn His Leu Asn Pro Arg Asn Trp Pro Ala Ala Leu Val Tyr
 65 70 75 80
 Gln Glu Ile Pro Gln Pro Thr Thr Val Leu Leu Pro Arg Asp Ala Gln
 85 90 95
 Ala Glu Val Gln Leu Thr Asn Ser Gly Val Gln Leu Ala Gly Gly Ala
 100 105 110
 Thr Leu Cys Arg His Arg Pro Ala Gln Gly Ile Lys Arg Leu Val Ile
 115 120 125

Arg Gly Arg Ser Thr Gln Leu Asn Asp Glu Val Val Ser Ser Ser Leu
 130 135 140
 Gly Leu Arg Pro Asp Gly Val Phe Gln Leu Ala Gly Ser Gly Arg Ser
 145 150 155 160
 Ser Phe Thr Pro Arg Gln Ala Val Leu Thr Leu Glu Ser Ser Ser Ser
 165 170 175
 Gln Pro Arg Ser Gly Gly Ile Gly Thr Leu Gln Phe Val Glu Glu Phe
 180 185 190
 Thr Pro Ser Val Tyr Phe Asn Pro Phe Ser Gly Ser Pro Gly His Tyr
 195 200 205
 Pro Asp Glu Phe Ile Pro Asn Phe Asp Ala Ile Ser Glu Ser Val Asp
 210 215 220
 Gly Tyr Asp
 225

<210> 20
 <211> 106
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 20

Met Ser His Gly Gly Ala Ala Asp Leu Ala Arg Leu Arg His Leu Asp
 1 5 10 15
 His Cys Arg Arg Phe Arg Cys Phe Ala Arg Asp Leu Ala Glu Phe Ala
 20 25 30
 Tyr Phe Glu Leu Pro Glu Glu His Pro Gln Gly Pro Ala His Gly Val
 35 40 45
 Arg Ile Val Val Glu Gly Gly Leu Asp Ser His Leu Leu Arg Ile Phe
 50 55 60
 Ser Gln Arg Pro Ile Leu Val Glu Arg Glu Gln Gly Gln Thr Leu Leu
 65 70 75 80
 Thr Leu Tyr Cys Ile Cys Asn His Pro Gly Leu His Glu Ser Leu Cys
 85 90 95
 Cys Leu Leu Cys Thr Glu Tyr Asn Lys Ser
 100 105

<210> 21
 <211> 146
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<220> <221> MISC_FEATURE
 <222> (62)..(62)
 <223> Xaa can be any amino acid

<400> 21

```

Met Lys Val Phe Val Val Cys Cys Val Leu Ser Ile Ile Lys Ala Glu
1          5          10          15

Thr Ala Thr Thr Pro Asp Phe Arg Val Ser Lys Leu Gln Leu Phe Gln
          20          25          30

Pro Phe Leu Pro Gly Thr Tyr Gln Cys Val Ser Gly Pro Cys His His
          35          40          45

Thr Phe His Leu Ile Pro Asn Thr Thr Ala Ser Leu Pro Xaa Thr Asn
          50          55          60

Asn Gln Thr Asn Leu His Gln Arg His Arg Arg Asp Leu Ser Glu Ser
65          70          75          80

Asn Thr Thr Thr His Thr Gly Gly Glu Leu Arg Gly Gln Pro Thr Ser
          85          90          95

Gly Ile Tyr Tyr Gly Pro Trp Glu Val Val Gly Leu Ile Thr Leu Gly
          100          105          110

Leu Val Ala Gly Gly Leu Leu Val Leu Cys Tyr Leu Tyr Leu Pro Cys
          115          120          125

Cys Ser Tyr Leu Val Val Leu Cys Cys Trp Phe Lys Lys Trp Gly Arg
130          135          140

Ser Pro
145

```

<210> 22

<211> 176

<212> PRT

<213> chimpanzee C68 adenovirus protein

<220>

<221> MISC_FEATURE

<222> (28)..(28)

<223> Xaa can be any amino acid

<400> 22

```

Met Gly Lys Ile Thr Leu Val Ser Cys Gly Ala Leu Val Ala Val Leu
1          5          10          15

Leu Ser Ile Val Gly Leu Gly Gly Ala Ala Val Xaa Lys Glu Lys Ala
          20          25          30

Asp Pro Cys Leu His Phe Asn Pro Asn Lys Cys Gln Leu Ser Phe Gln
          35          40          45

Pro Asp Gly Asn Arg Cys Ala Val Leu Ile Lys Cys Gly Trp Glu Cys
50          55          60

Glu Asn Val Arg Ile Glu Tyr Asn Asn Lys Thr Arg Asn Asn Thr Leu
65          70          75          80

Ala Ser Val Trp Gln Pro Gly Asp Pro Glu Trp Tyr Thr Val Ser Val
          85          90          95

```

Pro Gly Ala Asp Gly Ser Pro Arg Thr Val Asn Asn Thr Phe Ile Phe
 100 105 110
 Ala His Met Cys Asp Thr Val Met Trp Met Ser Lys Gln Tyr Asp Met
 115 120 125
 Trp Pro Pro Thr Lys Glu Asn Ile Val Val Phe Ser Ile Ala Tyr Ser
 130 135 140
 Leu Cys Thr Ala Leu Ile Thr Ala Ile Val Cys Leu Ser Ile His Met
 145 150 155 160
 Leu Ile Ala Ile Arg Pro Arg Asn Asn Ala Glu Lys Glu Lys Gln Pro
 165 170 175

<210> 23
 <211> 204
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 23

Met Ala Ser Val Lys Phe Leu Leu Leu Phe Ala Ser Leu Ile Ala Val
 1 5 10 15
 Ile His Gly Met Ser Asn Glu Lys Ile Thr Ile Tyr Thr Gly Thr Asn
 20 25 30
 His Thr Leu Lys Gly Pro Glu Lys Ala Thr Glu Val Ser Trp Tyr Cys
 35 40 45
 Tyr Phe Asn Glu Ser Asp Val Ser Thr Glu Leu Cys Gly Asn Asn Asn
 50 55 60
 Lys Lys Asn Glu Ser Ile Thr Leu Ile Lys Phe Gln Cys Gly Ser Asp
 65 70 75 80
 Leu Thr Leu Ile Asn Ile Thr Arg Asp Tyr Val Gly Met Tyr Tyr Gly
 85 90 95
 Thr Thr Ala Gly Ile Ser Asp Met Glu Phe Tyr Gln Val Ser Val Ser
 100 105 110
 Glu Pro Thr Thr Pro Arg Met Thr Thr Thr Thr Lys Thr Thr Pro Val
 115 120 125
 Thr Thr Met Gln Leu Thr Thr Asn Asn Ile Phe Ala Met Arg Gln Met
 130 135 140
 Val Asn Asn Ser Thr Gln Pro Thr Pro Pro Ser Glu Glu Ile Pro Lys
 145 150 155 160
 Ser Met Ile Gly Ile Ile Val Ala Val Val Val Cys Met Leu Ile Ile
 165 170 175
 Ala Leu Cys Met Val Tyr Tyr Ala Phe Cys Tyr Arg Lys His Arg Leu
 180 185 190
 Asn Asp Lys Leu Glu His Leu Leu Ser Val Glu Phe
 195 200

<210> 24
 <211> 91
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 24

```

Met Ile Pro Arg Gln Phe Leu Ile Thr Ile Leu Ile Cys Leu Leu Gln
1           5           10           15

Val Cys Ala Thr Leu Ala Leu Val Ala Asn Ala Ser Pro Asp Cys Ile
          20           25           30

Gly Pro Phe Ala Ser Tyr Val Leu Phe Ala Phe Thr Thr Cys Ile Cys
          35           40           45

Cys Cys Ser Ile Val Cys Leu Leu Ile Thr Phe Phe Gln Phe Ile Asp
          50           55           60

Trp Ile Phe Val Arg Ile Ala Tyr Leu Arg His His Pro Gln Tyr Arg
65           70           75           80

Asp Gln Arg Val Ala Arg Leu Leu Arg Leu Leu
          85           90
  
```

<210> 25
 <211> 143
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<220>
 <221> MISC_FEATURE
 <222> (5)..(5)
 <223> Xaa can be any amino acid

<400> 25

```

Met Arg Ala Val Xaa Leu Leu Ala Leu Leu Leu Val Leu Pro Arg
1           5           10           15

Pro Val Asp Pro Arg Ser Pro Thr Gln Ser Pro Glu Glu Val Arg Lys
          20           25           30

Cys Lys Phe Gln Glu Pro Trp Lys Phe Leu Lys Cys Tyr Arg Gln Lys
          35           40           45

Ser Asp Met His Pro Ser Trp Ile Met Ile Ile Gly Ile Val Asn Ile
          50           55           60

Leu Ala Cys Thr Leu Ile Ser Phe Val Ile Tyr Pro Cys Phe Asp Phe
65           70           75           80

Gly Trp Asn Ser Pro Glu Ala Leu Tyr Leu Pro Pro Glu Pro Asp Thr
          85           90           95

Pro Pro Gln Gln Pro Gln Ala His Ala Leu Pro Pro Leu Gln Pro Arg
          100          105          110

Pro Gln Tyr Met Pro Ile Leu Asp Tyr Glu Ala Glu Pro Gln Arg Pro
          115          120          125
  
```

Met Leu Pro Ala Ile Ser Tyr Phe Asn Leu Thr Gly Gly Asp Asp
 130 135 140

<210> 26
 <211> 135
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 26

Met Thr Asp Pro Leu Ala Asn Asn Asn Val Asn Asp Leu Leu Leu Asp
 1 5 10 15

Met Asp Gly Arg Ala Ser Glu Gln Arg Leu Ala Gln Leu Arg Ile Arg
 20 25 30

Gln Gln Gln Glu Arg Ala Val Lys Glu Leu Gln Asp Ala Val Ala Ile
 35 40 45

His Gln Cys Lys Arg Gly Ile Phe Cys Leu Val Lys Gln Ala Lys Ile
 50 55 60

Ser Tyr Glu Val Thr Pro Asn Asp His Arg Leu Ser Tyr Glu Leu Leu
 65 70 75 80

Gln Gln Arg Gln Lys Phe Thr Cys Leu Val Gly Val Asn Pro Ile Val
 85 90 95

Ile Thr Gln Gln Ser Gly Asp Thr Lys Gly Cys Ile His Cys Ser Cys
 100 105 110

Asp Ser Pro Asp Cys Val His Thr Leu Ile Lys Thr Leu Cys Gly Leu
 115 120 125

Arg Asp Leu Leu Pro Met Asn
 130 135

<210> 27
 <211> 425
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 27

Met Ser Lys Lys Arg Val Arg Val Asp Asp Asp Phe Asp Pro Val Tyr
 1 5 10 15

Pro Tyr Asp Ala Asp Asn Ala Pro Thr Val Pro Phe Ile Asn Pro Pro
 20 25 30

Phe Val Ser Ser Asp Gly Phe Gln Glu Lys Pro Leu Gly Val Leu Ser
 35 40 45

Leu Arg Leu Ala Asp Pro Val Thr Thr Lys Asn Gly Glu Ile Thr Leu
 50 55 60

Lys Leu Gly Glu Gly Val Asp Leu Asp Ser Ser Gly Lys Leu Ile Ser
 65 70 75 80

Asn Thr Ala Thr Lys Ala Ala Ala Pro Leu Ser Phe Ser Asn Asn Thr
 85 90 95
 Ile Ser Leu Asn Met Asp His Pro Phe Tyr Thr Lys Asp Gly Lys Leu
 100 105 110
 Ser Leu Gln Val Ser Pro Pro Leu Asn Ile Leu Arg Thr Ser Ile Leu
 115 120 125
 Asn Thr Leu Ala Leu Gly Phe Gly Ser Gly Leu Gly Leu Arg Gly Ser
 130 135 140
 Ala Leu Ala Val Gln Leu Val Ser Pro Leu Thr Phe Asp Thr Asp Gly
 145 150 155 160
 Asn Ile Lys Leu Thr Leu Asp Arg Gly Leu His Val Thr Thr Gly Asp
 165 170 175
 Ala Ile Glu Ser Asn Ile Ser Trp Ala Lys Gly Leu Lys Phe Glu Asp
 180 185 190
 Gly Ala Ile Ala Thr Asn Ile Gly Asn Gly Leu Glu Phe Gly Ser Ser
 195 200 205
 Ser Thr Glu Thr Gly Val Asp Asp Ala Tyr Pro Ile Gln Val Lys Leu
 210 215 220
 Gly Ser Gly Leu Ser Phe Asp Ser Thr Gly Ala Ile Met Ala Gly Asn
 225 230 235 240
 Lys Glu Asp Asp Lys Leu Thr Leu Trp Thr Thr Pro Asp Pro Ser Pro
 245 250 255
 Asn Cys Gln Ile Leu Ala Glu Asn Asp Ala Lys Leu Thr Leu Cys Leu
 260 265 270
 Thr Lys Cys Gly Ser Gln Ile Leu Ala Thr Val Ser Val Leu Val Val
 275 280 285
 Gly Ser Gly Asn Leu Asn Pro Ile Thr Gly Thr Val Ser Ser Ala Gln
 290 295 300
 Val Phe Leu Arg Phe Asp Ala Asn Gly Val Leu Leu Thr Glu His Ser
 305 310 315 320
 Thr Leu Lys Lys Tyr Trp Gly Tyr Arg Gln Gly Asp Ser Ile Asp Gly
 325 330 335
 Thr Pro Tyr Thr Asn Ala Val Gly Phe Met Pro Asn Leu Lys Ala Tyr
 340 345 350
 Pro Lys Ser Gln Ser Ser Thr Thr Lys Asn Asn Ile Val Gly Gln Val
 355 360 365
 Tyr Met Asn Gly Asp Val Ser Lys Pro Met Leu Leu Thr Ile Thr Leu
 370 375 380
 Asn Gly Thr Asp Asp Ser Asn Ser Thr Tyr Ser Met Ser Phe Ser Tyr
 385 390 395 400
 Thr Trp Thr Asn Gly Ser Tyr Val Gly Ala Thr Phe Gly Ala Asn Ser
 405 410 415

Tyr Thr Phe Ser Tyr Ile Ala Gln Glu
 420 425

<210> 28
 <211> 83
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 28

Ile Thr Val Ile Pro Thr Thr Glu Asp Asn Pro Gln Leu Leu Ser Cys
 1 5 10 15

Glu Val Gln Met Arg Glu Cys Pro Glu Gly Phe Ile Ser Leu Thr Asp
 20 25 30

Pro Arg Leu Ala Arg Ser Glu Thr Val Trp Asn Val Glu Thr Lys Ser
 35 40 45

Met Ser Ile Thr Asn Gly Ile Gln Met Phe Lys Ala Val Arg Gly Glu
 50 55 60

Arg Val Val Tyr Ser Met Ser Trp Glu Gly Gly Gly Lys Ile Thr Ala
 65 70 75 80

Arg Ile Leu

<210> 29
 <211> 301
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 29

Met Ser Glu Ser Asn Cys Ile Met Thr Arg Ser Arg Thr Arg Ser Ala
 1 5 10 15

Ala Ser Arg His His Pro Tyr Arg Pro Ala Pro Leu Pro Arg Cys Glu
 20 25 30

Glu Thr Glu Thr Arg Ala Ser Leu Val Glu Asp His Pro Val Leu Pro
 35 40 45

Asp Cys Asp Thr Leu Ser Met His Asn Val Ser Ser Val Arg Gly Leu
 50 55 60

Pro Cys Ser Ala Gly Phe Ala Val Leu Gln Glu Phe Pro Val Pro Trp
 65 70 75 80

Asp Met Val Leu Thr Pro Glu Glu Leu Arg Val Leu Lys Arg Cys Met
 85 90 95

Ser Ile Cys Leu Cys Cys Ala Asn Ile Asp Leu Phe Ser Ser Gln Met
 100 105 110

Ile His Gly Tyr Glu Arg Trp Val Leu His Cys His Cys Arg Asp Pro
 115 120 125

Gly Ser Leu Arg Cys Met Ala Gly Gly Ala Val Leu Ala Leu Trp Phe
 130 135 140

Arg Arg Ile Ile Arg Gly Cys Met Phe Asn Gln Arg Val Met Trp Tyr
 145 150 155 160
 Arg Glu Val Val Asn Arg His Met Pro Lys Glu Ile Met Tyr Val Gly
 165 170 175
 Ser Val Phe Trp Arg Gly His His Leu Ile Tyr Leu Arg Ile Trp Tyr
 180 185 190
 Asp Gly His Val Gly Ser Ile Leu Pro Ala Met Ser Phe Gly Trp Ser
 195 200 205
 Val Leu Asn Tyr Gly Leu Leu Asn Asn Leu Val Val Leu Cys Cys Thr
 210 215 220
 Tyr Cys Ser Asp Leu Ser Glu Ile Arg Met Arg Cys Cys Ala Arg Arg
 225 230 235 240
 Thr Arg Arg Leu Met Leu Arg Ala Val Gly Ile Met Leu Arg Glu Ser
 245 250 255
 Leu Asp Pro Asp Pro Leu Ser Ser Ser Leu Thr Glu Arg Arg Arg Gln
 260 265 270
 Arg Leu Leu Arg Gly Leu Met Arg His His Arg Pro Ile Pro Phe Ala
 275 280 285
 Asp Tyr Asp Ser His Arg Arg Ser Ser Ala Ser Ser Arg
 290 295 300

<210> 30
 <211> 121
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 30

Met Val Leu Pro Val Leu Pro Ser Pro Ala Val Thr Glu Thr Gln Gln
 1 5 10 15
 Asn Cys Ile Ile Trp Leu Gly Leu Ala His Ser Thr Val Val Asp Val
 20 25 30
 Ile Arg Ala Ile Arg His Asp Gly Ile Phe Ile Thr Pro Glu Ala Leu
 35 40 45
 Asp Leu Leu His Gly Leu Arg Glu Trp Leu Phe Tyr Asn Phe Asn Thr
 50 55 60
 Glu Arg Ser Lys Arg Arg Asp Arg Arg Arg Arg Ser Val Cys Ser Ala
 65 70 75 80
 Arg Thr Arg Phe Cys Tyr Ser Lys Tyr Glu Asn Val Arg Lys Gln Leu
 85 90 95
 His His Asp Thr Val Ala Asn Thr Ile Ser Arg Val Pro Pro Ser Pro
 100 105 110
 Val Ser Ala Gly Pro Leu Thr Thr Leu
 115 120

<210> 31
 <211> 117
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<220> <221> MISC_FEATURE
 <222> (45)..(45)
 <223> Xaa can be any amino acid

<400> 31

```

Met Arg Val Cys Leu Arg Met Pro Val Glu Gly Ala Leu Arg Glu Leu
1           5           10           15

Phe Ile Met Ala Gly Leu Asp Leu Pro His Glu Leu Val Arg Ile Ile
          20           25           30

Gln Gly Trp Lys Asn Glu Asn Tyr Leu Gly Met Val Xaa Glu Cys Asn
          35           40           45

Met Met Ile Glu Glu Leu Glu Asn Pro Pro Ala Phe Ala Ile Val Leu
 50           55           60

Phe Leu Asp Val Arg Val Glu Ala Leu Leu Glu Ala Thr Val Glu His
65           70           75           80

Leu Glu Asn Arg Ile Thr Phe Asp Leu Ala Val Ile Phe His Gln His
          85           90           95

Ser Gly Gly Glu Arg Cys His Leu Arg Asp Leu His Phe Glu Val Leu
          100          105          110

Arg Asp Arg Leu Asp
          115

```

<210> 32
 <211> 129
 <212> PRT
 <213> chimpanzee C68 adenovirus protein

<400> 32

```

Met Leu Glu Arg Thr Ala Cys Ile Tyr Phe Ile Val Val Pro Glu Ala
1           5           10           15

Leu Asn Val His Leu Glu Asp Phe Ser Phe Val Asp Phe Leu Lys Asn
          20           25           30

Cys Leu Gly Asp Phe Leu Ser Ser Tyr Leu Glu Asp Ile Thr Gly Ser
          35           40           45

Ser Gln His Ala Tyr Ser Ser Leu Ala Phe Gly Asn Ala His Trp Gly
 50           55           60

Gly Leu Arg Phe Ile Cys Thr Val Ala Cys Pro Asn Leu Ile Pro Gly
65           70           75           80

Gly Pro Met Ala Lys Asn Phe Gly Glu Asp Met Lys Glu Tyr Leu Gln
          85           90           95

```

Leu Leu Leu Arg Glu Glu Leu Arg Asp Arg Gly Arg Asp Phe Asp Ile
 100 105 110

Pro Leu Val Asn Leu Leu Gln Val Asn Gln Glu Gln Asn Ile Leu Glu
 115 120 125

Leu

<210> 33
 <211> 36521
 <212> DNA
 <213> chimpanzee C68 adenovirus

<220>
 <221> misc_feature
 <222> (8268)..(8268)
 <223> can be a or c or g or t

<220>
 <221> misc_feature
 <222> (8322)..(8322)
 <223> can be a or c or g or t

<220>
 <221> misc_feature
 <222> (8535)..(8535)
 <223> can be a or c or g or t

<220>
 <221> misc_feature
 <222> (16753)..(16753)
 <223> can be a or c or g or t

<220>
 <221> misc_feature
 <222> (28095)..(28095)
 <223> can be a or c or g or t

<220>
 <221> misc_feature
 <222> (29373)..(29373)
 <223> can be a or c or g or t

<220>
 <221> misc_feature
 <222> (30447)..(30447)
 <223> can be a or c or g or t

<220>
 <221> misc_feature
 <222> (31015)..(31015)
 <223> can be a or c or g or t

<400> 33
 ccttcttcaa taatatacct tcaaactttt tgtgcgcggtt aatatgcaaa tgaggcggtt 60
 gaatttgggg aggaagggcg gtgattgggtc gagggatgag cgaccgtag gggcggggag 120
 agtgacgttt tgatgacgtg gttgagagga ggagccagtt tgcaagttct cgtgggaaaa 180
 gtgacgtcaa acgaggtgtg gtttgaacac ggaaatactc aattttcccg cgctctctga 240

caggaaatga ggtgtttctg ggcggatgca agtgaaaacg ggccattttc gcgcgaaaac	300
tgaatgagga agtgaaaatc tgagtaattt cgcgtttatg gcagggagga gtatttgccg	360
agggccgagt agactttgac cgatcacgtg ggggtttcga ttaccgtgtt tttcacctaa	420
atttccgcgt acggtgtcaa agtccggtgt ttttacgtag gtgtcagctg atcgccaggg	480
tatttaaacc tgcgctctcc agtcaagagg ccactcttga gtgccagcga gaagagtttt	540
ctcctccgcg ccgcgagtca gatctacact ttgaaagatg aggcacctga gagacctgcc	600
cgatgagaaa atcatcatcg cttccgggaa cgagattctg gaactggtgg taaatgccat	660
gatgggcgac gaccctccgg agccccccac cccatttgag acaccttcgc tgcacgattt	720
gtatgatctg gaggtggatg tgcccagga cgatcccaat gaggaggcgg taaatgattt	780
ttttagcgat gccgcgctgc tagctgccga ggaggcttcg agctctagct cagacagcga	840
ctcttcaactg cataccccta gaccggcag aggtgagaaa aagatccccg agcttaaagg	900
ggaagagatg gacttgcgct gctatgagga atgcttgccc ccgagcgatg atgaggacga	960
gcaggcgatc cagaacgcag cgagccaggg agtgcaagcc gccagcgaga gctttgcgct	1020
ggactgcccg cctctgcccg gacacggctg taagtcttgt gaatttcacg gcatgaatac	1080
tggagataaa gctgtgttgt gtgcactttg ctatatgaga gcttacaacc attgtgttta	1140
cagtaagtgt gattaagttg aacttttagag ggaggcagag agcagggtga ctgggcgatg	1200
actggtttat ttatgtatat atgttcttta tataggcccc gtctctgacg cagatgatga	1260
gacccccact acaaagtcca cttcgtcacc ccagaaaatt ggcacatctc cacctgagaa	1320
tattgttaga ccagttcctg ttagagccac tgggaggaga gcagctgtgg aatgtttgga	1380
tgacttgcta cagggtgggg ttgaaccttt ggacttgtgt acccggaac gccccaggca	1440
ctaagtgcc aacatgtgtg tttacttgag gtgatgtcag tatttatagg gtgtggagtg	1500
caataaaaaa tgtgttgact ttaagtgcgt ggtttatgac tcaggggtgg ggactgtgag	1560
tatataagca ggtgcagacc tgtgtggtta gctcagagcg gcatggagat ttggacggtc	1620
ttggaagact ttcacaagac tagacagctg ctagagaacg cctcgaacgg agtctcttac	1680
ctgtggagat tctgcttcgg tggcgacctg gctaggctag tctacagggc caaacaggat	1740
tatagtgaac aatttgaggt tattttgaga gagtgttctg gtctttttga cgctcttaac	1800
ttggggccatc agtctcactt taaccagagg atttcgagag cccttgattt tactactcct	1860
ggcagaacca ctgcagcagt agcctttttt gcttttatctc ttgacaaatg gagtcaagaa	1920
acccatttca gcagggatta ccagctggat ttcttagcag tagctttgtg gagaacatgg	1980
aagtgccagc gcctgaatgc aatctccggc tacttgccgg tacagccgct agacactctg	2040
aggatcctga atctccagga gagtcccagg gcacgccaac gtcgccagca gcagcagcag	2100

gaggaggatc	aagaagagaa	cccgagagcc	ggcctggacc	ctccggcgga	ggaggaggag	2160
tagctgacct	gtttcctgaa	ctgcgccggg	tgctgactag	gtcttcgagt	ggtcggggaga	2220
gggggattaa	gcgggagagg	catgatgaga	ctaatacacag	aactgaactg	actgtgggtc	2280
tgatgagtcg	caagcgccca	gaaacagtgt	ggtggcatga	ggtgcagtcg	actggcacag	2340
atgaggtgtc	ggtgatgcat	gagaggtttt	ctctagaaca	agtcaagact	tgttggtttag	2400
agcctgagga	tgattgggag	gtagccatca	ggaattatgc	caagctggct	ctgaggccag	2460
acaagaagta	caagattact	aagctgataa	atatcagaaa	tgcttgctac	atctcaggga	2520
atggggctga	agtggagatc	tgtctccagg	aaaggggtggc	tttcagatgc	tgcatgatga	2580
atatgtaccc	gggagtgggtg	ggcatggatg	gggttacctt	tatgaacatg	aggttcaggg	2640
gagatgggta	taatggcacg	gtcttttatgg	ccaataccaa	gctgacagtc	catggctgct	2700
ccttcttttg	gtttaataac	acctgcatcg	aggcctgggg	tcaggtcggt	gtgaggggct	2760
gcagtttttc	agccaactgg	atgggggtcg	tgggcaggac	caagagtatg	ctgtccgtga	2820
agaaatgctt	gtttgagagg	tgccacctgg	gggtgatgag	cgagggcgaa	gccagaatcc	2880
gccactgcgc	ctctaccgag	acgggctgct	ttgtgctgtg	caagggcaat	gctaagatca	2940
agcataatat	gatctgtgga	gcctcggacg	agcgcggtca	ccagatgctg	acctgcgccg	3000
gcgggaacag	ccatatgctg	gccaccgtac	atgtggcttc	ccatgctcgc	aagccctggc	3060
ccgagttcga	gcacaatgtc	atgaccaggt	gcaatatgca	tctgggggtcc	cgccgaggca	3120
tgttcatgcc	ctaccagtgc	aacctgaatt	atgtgaaggt	gctgctggag	cccgatgcc	3180
tgtccagagt	gagcctgacg	gggggtgtttg	acatgaatgt	ggaggtgtgg	aagattctga	3240
gatatgatga	atccaagacc	aggtgccgag	cctgcgagtg	cggaggggaag	catgccaggt	3300
tccagcccgt	gtgtgtggat	gtgacggagg	acctgcgacc	cgatcatttg	gtgttgccct	3360
gcaccgggac	ggagttcgggt	tccagcgggg	aagaatctga	ctagagtgag	tagtgttctg	3420
gggcggggga	ggacctgcat	gagggccaga	ataactgaaa	tctgtgcttt	tctgtgtgtt	3480
gcagcagcat	gagcggaagc	ggctcctttg	agggaggggt	attcagccct	tatctgacgg	3540
ggcgtctccc	ctcctgggcg	ggagtgcgtc	agaatgtgat	gggatccacg	gtggacggcc	3600
ggcccgtgca	gcccgcgaac	tcttcaacct	tgacctatgc	aacctgagc	tcttcgtcgt	3660
tggacgcagc	tgccgccgca	gctgctgcat	ctgccgccag	cgccgtgcgc	ggaatggcca	3720
tgggcgccgg	ctactacggc	actctgggtg	ccaactcgag	ttccaccaat	aatcccgcca	3780
gcctgaacga	ggagaagctg	ttgctgctga	tggcccagct	cgaggccttg	accagcgcc	3840
tgggcgagct	gaccagcag	gtggctcagc	tgaggagca	gacgcgggcc	gcggttgcca	3900
cggtgaaatc	caaataaaaa	atgaatcaat	aaataaacgg	agacggttgt	tgattttaac	3960

acagagtctg	aatcttttatt	tgattttttcg	cgcgcggttag	gccctggacc	accggtctctg	4020
atcattgagc	acccggtgga	tctttttccag	gacccggttag	aggtgggctt	ggatgttgag	4080
gtacatgggc	atgagcccg	cccgggggtg	gaggtagctc	cattgcaggg	cctcgtgctc	4140
gggggtggtg	ttgtaaata	cccagtcata	gcagggg	agggcatggt	gttgcaaat	4200
atctttgagg	aggagactga	tggccacggg	cagccctttg	gtgtaggtgt	ttacaaatct	4260
gttgagctgg	gagggatgca	tgcgggggga	gatgaggtgc	atcttggcct	ggatcttgag	4320
attggcgatg	ttaccgcccc	gatcccgctt	ggggttcatg	ttgtgcagga	ccaccagcac	4380
ggtgtatccg	gtgcacttgg	ggaatttatc	atgcaacttg	gaagggaaag	cgtgaaagaa	4440
tttggcgacg	cctttgtgcc	cgcccagggt	ttccatgcac	tcatccatga	tgatggcgat	4500
gggcccgtgg	gcggcgccct	gggcaaagac	gtttcggggg	tcggacacat	catagttgtg	4560
gtcctgggtg	aggtcatcat	aggccatttt	aatgaatttg	gggcggaggg	tgccggactg	4620
ggggacaaag	gtaccctcga	tcccgggggc	gtagttcccc	tcacagatct	gcatctccca	4680
ggctttgagc	tcggaggggg	ggatcatgtc	cacctgcggg	gcgataaaga	acacggtttc	4740
cggggcgggg	gagatgagct	gggccgaaag	caagttccgg	agcagctggg	acttgccgca	4800
gccggtgggg	ccgtagatga	ccccgatgac	cggctgcagg	tggtagttga	gggagagaca	4860
gctgccgtcc	tcccggagga	ggggggccac	ctcgttcatc	atctcgcgca	cgtgcatggt	4920
ctcgcgcacc	agttccgcca	ggaggcgctc	tccccccagg	gataggagct	cctggagcga	4980
ggcgaagttt	ttcagcggtc	tgagtccgtc	ggccatgggc	attttgagga	gggtttgttg	5040
caagagttcc	aggcgggtcc	agagctcggt	gatgtgctct	acggcatctc	gatccagcag	5100
acctcctcgt	ttcgcggggt	gggacggctg	cgggagtatg	gcaccagacg	atgggcgtcc	5160
agcgcagcca	gggtccggtc	cttcagggt	cgcagcgctc	gcgtcagggt	ggtctccgtc	5220
acggtgaagg	ggtgcgcgcc	gggctggg	cttgcgagg	tgcgcttcag	gctcatccgg	5280
ctggtcgaaa	accgctcccc	atcggcgccc	tgcgcgctcg	ccaggtagca	attgaccatg	5340
agttcgtagt	tgagcgctc	ggccgcgtg	cctttggcgc	ggagcttacc	tttggaagtc	5400
tgcccgagg	cgggacagag	gagggacttg	agggcgtaga	gcttgggggc	gaggaagacg	5460
gactcggggg	cgtaggcgtc	cgcgccgcag	tgggcgcaga	cggctctcgca	ctccacgagc	5520
caggtgaggt	cgggctggtc	ggggtcaaaa	accagtttcc	cgccgttctt	tttgatgcgt	5580
ttcttacctt	tggtctccat	gagctcgtgt	ctccgctggg	tgacaaagag	ctgtccgtgt	5640
ccccgtagac	cgactttatg	ggccgggtcct	cgagcggtgt	gccgcgggtc	tcctcgtaga	5700
ggaaccccg	ccactccgag	acgaaagccc	gggtccaggc	cagcacgaaw	gagggcacgt	5760
gggacgggta	tcggtcgttg	tccaccagcg	ggtccacctt	ttccagggta	tgcaaacaca	5820

tgtccccctc	gtccacatcc	aggaaggtga	ttggcttgta	agtgtakgcc	acgtgaccgg	5880
gggtcccggc	cgggggggta	taaaaggggtg	cgggtccctg	ctcgtcctca	ctgtcttccg	5940
gatcgctgtc	cacgagcgcc	agctgttggg	gtaggtattc	cctctcgaag	gcgggcatga	6000
cctcggcact	caggttgtca	gtttctagaa	acgaggagga	tttgatattg	acggtgccgg	6060
cggagatgcc	tttcaagagc	ccctcgtcca	tctggtcaga	aaagacgata	tttttgttgt	6120
cgagcttggg	ggcgaaggag	ccgtakaggg	cgttgagag	gagcttggcg	atggagcgca	6180
tggctctggt	tttttccttg	tcggcgcgct	ccttggcggc	gatgttgagc	tgcacgtact	6240
cgcgcgccac	gcacttccat	tcggggaaga	cgtggtcagc	tcgtcgggca	cgattctgac	6300
ctgccagccc	cgattatgca	gggtgatgag	gtccacactg	gtggccacct	cgccgcgcag	6360
gggctcatta	ktccagcaga	ggcgtccgcc	cttgccgcgag	cagaaggggg	gcaggggggtc	6420
cagcatgacc	tcgtcggggg	ggtcggcatc	gatggtgaag	atgccgggca	ggaggtcggg	6480
gtcaaagtag	ctgatggaag	tggccagatc	gtccagggca	gcttgccatt	cgcgcacggc	6540
cagcgcgcgc	tcgtagggac	tgaggggctg	gccccagggc	atgggatggg	taagcgcgga	6600
ggcgtacatg	ccgcagatgt	cgtagacgta	gaggggctcc	tcgaggatgc	cgatgtaggt	6660
ggggtagcag	cgccccccgc	ggatgctggc	gcgcacgtag	tcatacagct	cgtgcgaggg	6720
ggcgaggagc	cccgggcccc	ggttggtgcg	actgggcttt	tcggcgcggt	agacgatctg	6780
gcggaatatg	gcatgcgagt	tggaggagat	ggtgggcctt	tggaagatgt	tgaagtgggc	6840
gtggggcagt	ccgaccgagt	cgcggatgaa	gtgggcgtag	gagtcttgca	gcttggcgac	6900
gagctcggcg	gtgactagga	cgtccagagc	gcagtagtcg	agggctctct	ggatgatgtc	6960
atacttgagc	tgtccctttt	gtttccacag	ctcgcggttg	agaaggaact	cttcgcggtc	7020
cttccagtac	tcttcgaggg	ggaacccgtc	ctgatctgca	cggtaagagc	ctagcatgta	7080
gaactgggtg	acggccttgt	aggcgcagca	gccyttctcc	acggggargg	cgtaggcctg	7140
ggcggccttg	cgcagggagg	tgtgcgtgag	ggcgaaagtg	tccctgacca	tgaccttgag	7200
gaactgggtc	ttgaagtcga	tattgtcgca	gccccctgc	tcccagagct	ggaagtccgt	7260
gcgcttcttg	taggcggggg	tgggcaaagc	gaaagtaaca	tcgttgaaga	ggatcttgcc	7320
cgcgcggggc	ataaagttgc	gagtgatgag	gaaagggttg	ggcacctcgg	cccggttggt	7380
gatgacctgg	gcggcgagca	cgatctcgtc	gaagccgttg	atgttggtgg	ccacgatgta	7440
gagttccacg	aatcgcggac	ggcccttgac	gtggggcagt	ttcttgagct	cctcgtaggt	7500
gagctcgtcg	gggtcgctga	gcccgtgctg	ctcgagcgcc	cagtcggcga	gatggggggt	7560
ggcgcggagg	aaggaagtcc	agagatccac	ggccaggggc	gtttgcagac	ggccccggta	7620
ctgacggaac	tgctgcccga	cggccatttt	ttcgggggtg	acgcagtaga	aggtgcgggg	7680

gtccccgtgc	cagcgatccc	atttgagctg	gagggcgaga	tcgagggcga	gctcgacgag	7740
ccggtcgtcc	ccggagagtt	tcatgaccag	catgaagggg	acgagctgct	tgccgaagga	7800
ccccatccag	gtgtaggttt	ccacatcgta	ggtgaggaag	agcctttcgg	tgcgaggatg	7860
cgagccgatg	gggaagaact	ggatctcctg	ccaccaattg	gaggaatggc	tgttgatgtg	7920
atggaagtag	aaatgccgac	ggcgcgccga	acactcgtgc	ttgtgtttat	acaagcggcc	7980
acagtgcctg	caacgctgca	cgggatgcac	gtgctgcacg	agctgtacct	gagttccttt	8040
gacgaggaat	ttcagtggga	agtggagtcg	tggcgccctg	atctcgtgct	gtactacgtc	8100
gtggtggtcg	gcctggccct	cttctgcctc	gatggtggtc	atgctgacga	gcccgcgcgg	8160
gaggcaggtc	cagacctcgg	cgcgagcggg	tcggagagcg	aagacgaagg	cgcgcaggcc	8220
ggagctgtcc	agggctctga	gacgctgcgg	agtcagggtca	gtgggcancg	gcggcgcgcg	8280
gttgacttgc	argagttttt	ccagggcgcg	cgggaggtcc	anatggtact	tgatctccac	8340
cgcgccattg	gtggcgaaact	ccatggcttg	caggggtccc	tgcccctggg	gtgtgaccac	8400
cgtccccctg	ttcttcttgg	gcggctgggg	cgacgggggc	ggtgcctctt	ccatgggttag	8460
aascggcggc	gaagacgcgc	gccgggcggc	aggggcggct	cggggcccgg	atgcaggggc	8520
ggcaggggca	cttcngcgcc	gcgcgcgggt	aggttctggt	actgcgccc	gagaaaactg	8580
gcgtgagcga	cgacgcgacg	gttgacgtcc	tggatctgac	gcctctgggt	gaaggccacg	8640
ggacccgtga	gtttgaacct	gaaagaaagt	tcgacagaat	caatctcggg	atcgttgacg	8700
gcggcctgcc	gcaagatctc	ttgcacgtcc	cccaggttgt	cctggtatgc	gatctcggtc	8760
atgaactgct	cgatctcctc	ctcttgaagg	tctccgcggc	cggcgcgctc	cacggtggcc	8820
gcgaagtcgt	tggagatgcg	gcccattgag	tcgagaagg	cgttcatgcc	cgcctcgttc	8880
cagacgcggc	tgtagaccac	gacgccctcg	ggatcgcggg	cgcgcatgac	cacctgggcg	8940
aggttgagct	ccacgtggcg	cgtgaagacc	gcgtagttgc	agaggcgctg	gtagaggtag	9000
ttgagcgtgg	tggcgatgtg	ctcgggtgac	aagaaataca	tgatccagcg	gcggagcggc	9060
atctcgtgta	cgtcgcccag	cgcctccaaa	cgttccatgg	cctcgtaaaa	gtccacggcg	9120
aagttgaaaa	actgggagtt	gcgcgccgag	acggtcaact	cctcctccag	aagacggatg	9180
agctcggcga	tgggtggcgcg	cacctcgcg	tcgaaggccc	ccgggagttc	ctccacttcc	9240
tcttcttctc	cctccactaa	catctcttct	acttcctcct	caggcggcag	tgggtggcggg	9300
ggagggggcc	tgcgtcgccg	gcggcgacg	ggcagacggt	cgatgaagcg	ctcgatggtc	9360
tcgccgcgcc	ggcgtcgcat	ggtctcggtg	acggcgcgcc	cgtcctcgcg	gggccgcakc	9420
gtgaagacgc	cgccgcgcat	ctccaggtgg	ccgggggggt	ccccgttgg	gcaggagagag	9480
ggcgtgacg	atgcatctta	tcaattgccc	cgtagggact	ccgcgcaagg	acctgagcgt	9540

ctcgagatcc	acgggatctg	aaaaccgctg	aacgaaggct	tcgaagccag	tcgcagtcgc	9600
aaggtakgct	gagcacgggt	tcttctggcg	ggcatgttg	gttgggagcg	gggcgggcga	9660
tgctgctggt	gatgaagttg	aaataggcgg	ttctgagacg	gcggatggtg	gcgargagca	9720
ccaggctctt	gggcccggct	tgctggatgc	gcagacggtc	ggccatgcc	caggcgtggt	9780
cctgacacct	ggccaggctc	ttgtagtagt	cctgcatgag	ccgctccaac	gggcacctcc	9840
tcctcgcccc	cgcgcccggt	catgcgcgtg	agcccgaagc	cgcgctgggg	ctggacgagc	9900
gccaggctcg	cgacgacgcg	ctcggcgagg	atggcttgct	ggatctgggt	gagggtggtc	9960
tggaagtcac	caaagtcgac	gaagcgggtg	taggctccgg	tgttgatggt	gtaggagcag	10020
ttggccatga	cggaccagtt	gacggtctgg	tggcccggac	gcacgagctc	gtggtacttg	10080
aggcgcgagt	aggcgcgctg	gtcgaagatg	tagtcgttgc	aggcgcgac	cagggtactgg	10140
tagccgatga	ggaagtgcgg	cggcggctgg	cggtagagcg	gccatcgctc	ggtggcgggg	10200
gcgccggggc	cgaggtcctc	gagcatggtg	cggtggtagc	cgtagatgta	cctggacatc	10260
cagggtgatgc	cggcggcggt	ggtggaggcg	cgcggaact	cgcgacgcg	ttccagatgt	10320
tgcgcagcgg	caggaagtag	ttcatgggtg	gcacggtctg	gcccgtgagg	cgcgcgagc	10380
cgtggatgct	ctatacgggc	aaaaacgaaa	gcggtcagcg	gctcgactcc	gtggcctgga	10440
ggctaagcga	acgggttggg	ctgcgcgtgt	accccggttc	gaatctcgaa	tcaggctgga	10500
gccgcagcta	acgtggtatt	ggcactccc	tctmgacca	agcbtgacc	aaccctccag	10560
gatacggagg	cgggtcgttt	tgcaactttt	ttttggaggc	cggatgagac	tagtaagcgc	10620
ggaaagcggc	cgaccgcgat	ggctcgtctg	ccgtagtctg	gagaagaatc	gccagggttg	10680
cgttgcggtg	tgccccggtt	cgaggccggc	cggattccgc	ggctaacgag	ggcgtggctg	10740
ccccgtcgtt	tccaagacct	catagccagc	cgacttctcc	agttacggag	cgagcccctc	10800
ttttgttttg	tttgtttttg	ccagatgcat	cccgtactgc	ggcagatgcg	ccccaccac	10860
cctccaccgc	aacaacagcc	ccctccacag	ccggcgcttc	tgcccccgcc	ccagcagcaa	10920
cttccagcca	cgaccgccgc	ggccgccgtg	agcggggctg	gacagagtta	tgatcaccag	10980
ctggccttg	aagagggcga	ggggctggcg	cgctggggg	cgctcgtcgc	ggagcggcac	11040
ccgcgcgtgc	agatgaaaag	ggacgctcgc	gaggcatacg	tgcccaagca	gaacctgttc	11100
agagacagga	gcggcgagga	gcccaggagg	atgcgcgcgg	cccgggtcca	cgcggggcgg	11160
gagctgcggc	gcggcctgga	ccgaaagagg	gtgctgaggg	acgaggattt	cgaggcggac	11220
gagctgacgg	ggatcagccc	cgcgcgcgcg	cacgtggccg	cggccaacct	ggtcacggcg	11280
tacgagcaga	ccgtgaagga	ggagagcaac	ttcaaaaaat	ccttcaacaa	ccacgtgcgc	11340
accctgatcg	cgcgcgagga	ggtgaccctg	ggcctgatgc	acctgtggga	cctgctggag	11400

gccatcgtgc agaacccac cagcaagccg ctgacggcgc agctgttctt ggtggtgcag 11460
catagtcggg acaacgaagc gttcagggag gcgctgctga atatcaccga gcccgagggc 11520
cgctggctcc tggacctggt gaacattctg cagagcatcg tgggtgcagga gcgcgggctg 11580
ccgctgtccg agaagctggc ggccatcaac ttctcggtgc tgagtttggg caagtactac 11640
gctaggaaga tctacaagac cccgtacgtg cccatagaca aggaggtgaa gatcgacggg 11700
ttttacatgc gcatgaccct gaaagtgtg accctgagcg acgatctggg ggtgtaccgc 11760
aacgacagga tgcaccgtgc ggtgagcgcc agcaggcggc gcgagctgag cgaccaggag 11820
ctgatgcata gtctgcagcg ggccctgacc ggggcccggg ccgaggggga gagctacttt 11880
gacatgggcg cggacctgca ctggcagccc agccgcggg ccttggaggc ggcggcagga 11940
ccctacgtag aagaggtgga cgatgaggtg gacgaggagg gcgagtacct ggaagactga 12000
tggcgcgacc gtatttttgc tagatgcaac aacaacagcc acctctgat cccgcgatgc 12060
gggcgcgct gcagagccag ccgtccggca ttaactctc ggacgattgg acccaggcca 12120
tgcaacgcat catggcgctg acgaccgcga accccgaagc ctttagacag cagccccagg 12180
ccaaccggct ctcggccatc ctggaggccg tggtgccctc gcgctccaac cccacgcacg 12240
agaaggtcct ggccatcgtg aacgcgctgg tggagaacaa ggccatccgc ggcgacgagg 12300
ccggcctggt gtacaacgcg ctgctggagc gcgtggcccg ctacaacagc accaacgtgc 12360
agaccaacct ggaccgcatg gtgaccgacg tgcgcgaggc cgtggcccag cgcgagcggt 12420
tccaccgcga gtccaacctg ggatccatgg tggcgctgaa cgccttctc agcaccagc 12480
ccgccaacgt gccccggg caggaggact acaccaactt catcagcgcc ctgcgcctga 12540
tggtgaccga ggtgccccag agcgaggtgt accagtccgg gccggactac ttcttccaga 12600
ccagtgcga gggcttgca accgtgaacc tgagccaggc tttcaagaac ttgcagggcc 12660
tgtggggcgt gcaggccccg gtcggggacc gcgcgacggt gtcgagcctg ctgacgccga 12720
actcgcgct gctgctgctg ctggtggccc ccttcacgga cagcggcagc atcaaccgca 12780
actcgtacct gggctacctg attaacctgt accgcgaggc catcggccag gcgcacgtgg 12840
acgagcagac ctaccaggag atcaccacg tgagccgcgc cctgggccag gacgaccgg 12900
gcaacctgga agccacctg aactttttgc tgaccaaccg gtcgcagaag atcccgcccc 12960
agtacgcgct cagcaccgag gaggagcgca tcctgcgtta cgtgcagcag aagcgtgggc 13020
ctgttctga tgaggagg ggccacccc agcgcgcgcg tcgacatgac cgcgcgcaac 13080
atggtagccc atcatgtacg ccagcaaccg cccgttcac aataaactga tggactactt 13140
gcatcgggcg gccgcatga actctgacta tttaccaac gccatcctga atccccactg 13200
gctcccgccg ccggggttct acacgggcga gtacgacatg cccgacccca atgacggggt 13260

cctgtgggac gatgtggaca gcagcgtggt ctccccccga ccgggtgcta acgagcgccc 13320
cttgtggaag aaggaaggca gcgaccgacg cccgtcctcg gcgctgtccg gccgcgaggg 13380
tgctgccgcg gcggtgcccg aggccgccag tcctttcccg agcttgccct tctcgctgaa 13440
cagtatccgc agcagcgagc tgggcaggat cacgcgcccg cgcttgctgg gcgaagagga 13500
gtacttgaat gactcgctgt tgagaccga gcgggagaag aacttcccca ataacgggat 13560
agaaagcctg gtggacaaga tgagccgctg gaagacgtat gcgcaggagc acagggacga 13620
tccccgggcg tcgcaggggg ccacgagccg gggcagcgcc gcccgtaaac gccggtggca 13680
cgacaggcag cggggacaga tgtgggacga tgaggactcc gccgacgaca gcagcgtggt 13740
ggacttgggt gggagtggta acccgttcgc tcacctgcgc ccccgatatcg ggcgcatgat 13800
gtaagagaaa ccgaaaataa atgataactca ccaaggccat ggcgaccagc gtgcgttcgt 13860
ttcttctctg ttgttgttgt atctagtatg atgaggcgtg cgtaccgga gggtcctcct 13920
ccctcgtagc agagcgtgat gcagcaggcg atggcggcgg cggcgatgca gccccgctg 13980
gaggctcctt acgtgcccc gcggtacctg gcgcctacgg aggggcggaa cagcattcgt 14040
tactcgagc tggcaccctt gtacgatacc acccggttgt acctggtgga caacaagtcg 14100
gcggacatcg cctcgctgaa ctaccagaac gaccacagca acttcctgac caccgtggtg 14160
cagaacaatg acttcacccc cacggaggcc agcaccaga ccatcaactt tgacgagcgc 14220
tcgcggtggg gcggccagct gaaaaccatc atgcacacca acatgcccac cgtgaacgag 14280
ttcatgtaca gcaacaagtt caaggcgcgg gtgatggtct cccgcaagac cccaatggg 14340
gtgacagtga cagaggatta tgatgtagt caggatgagc tgaagtatga atgggtggaa 14400
tttgagctgc ccgaaggcaa cttctcggtg accatgacca tcgacctgat gaacaacgcc 14460
atcatcgaca attacttggc ggtggggcgg cagaacgggg tgctggagag cgacatcggc 14520
gtgaagtctg acactaggaa cttcaggctg ggctgggacc ccgtgaccga gctggtcatg 14580
cccggggtgt acaccaacga ggctttccat cccgatattg tcttgctgcc cggctgcggg 14640
gtggacttca ccgagagccg cctcagcaac ctgctgggca ttcgcaagag gcagcccttc 14700
caggaaggct tccagatcat gtacgaggat ctggaggggg gcaacatccc cgcgctcctg 14760
gatgtcgacg cctatgagaa aagcaaggag gatgcagcag ctgaagcaac tgcagccgta 14820
gctaccgcct ctaccgaggt caggggcgat aattttgcaa gcgccgcagc agtggcagcg 14880
gccgaggcgg ctgaaaccga aagtaagata gtcattcagc cgggtggagaa ggatagcaag 14940
aacaggagct acaacgtact accggacaag ataaacaccg cctaccgcag ctggtacctt 15000
gcctacaact atggcgaccc cgagaagggc gtgcgctcct ggacgctgct caccacctcg 15060
gacgtcacct gcggcgtgga gcaagtctac tggctcgctgc ccgacatgat gcaagacccg 15120

gtcaccttcc gctccacgcg tcaagtttagc aactaccgcg tgggtgggcgcg cgagctcctg 15180
 cccgtctact ccaagagctt cttcaacgag caggccgtct actcgagca gctgcgcgcc 15240
 ttcacctcgc ttacgcacgt cttcaaccgc ttccccgaga accagatcct cgtccgcccc 15300
 cccgcgcccc ccattaccac cgtcagtga aacgttccctg ctctcacaga tcacgggacc 15360
 ctgccgctgc gcagcagtat ccggggagtc cagcgcggtga ccgttactga cgccagacgc 15420
 cgcacctgcc cctacgtcta caaggccctg ggcatagtcg cgccgcgctg cctctcgagc 15480
 cgcaccttct aaatgtccat tctcatctcg ccagtaata acaccggttg gggcctgcgc 15540
 gcgcccagca agatgtacgg aggcgctcgc caacgctcca cgcaacaccc cgtgcgctg 15600
 cgcgggcact tccgcgctcc ctggggcgcc ctcaaaggcc gcgtgcggtc gcgcaccacc 15660
 gtcgacgacg tgatcgacca ggtggtggcs gacgcgcgca amtacacccc cgccgcggcg 15720
 cctgtttcca cygtggacgc cgtcatcgac agcgtggtgg cggacgcgcg ccggtacgcc 15780
 cgcgccaaga gccggcgggcg gcgcatcgcc cggcggcacc ggagcacccc cgccatgcgc 15840
 gcggcgcgga gccttgttgc gcagggccag gcgcacggga cgcagggcca tgttcagggc 15900
 ggccagacgc gcggcttcag gcgccagcgc cggcaggacc cggagacgcg cggccacggc 15960
 ggcggcagcg gccatcgcca gcatgtcccg cccgcggcga gggaacgtgt actgggtgcg 16020
 cgacgccgcc accggtgtgc gcgtgcccgt gcgcacccgc cccctcgca cttgaagatg 16080
 ttcacttcgc gatgttgatg tgtcccagcg gcgaggagga tgtccaagcg caaattcaag 16140
 gaagagatgc tccaggtcat cgcgcctgag atatacggcc ctgcggtggt gaaggaggaa 16200
 agaaagcccc gcaaaatcaa gcgggtcaaa aaggacaaaa aggaagaaga aagtgatgtg 16260
 gacggattgg tggagtttgt gcgcgagttc gcccccgcc ggcgcggtgca gtggcgcggg 16320
 cggaaggtgc aaccggtgct gagaccggc accaccgtgg tcttcacgcc cggcgagcgc 16380
 tccggcaccg cttccaagcg ctctacgac gaggtgtacg gggatgatga tattctggag 16440
 caggcgcccg akcgctggg cgagtttgct tacggcaagc gcagccgttc cgcaccgaag 16500
 gaagaggcgg tgtccatccc gctggaccac ggcaacccca cgccgagcct caagcccgtg 16560
 acctgcagc aggtgctgcc gaccgcggcg ccgcgccggg ggttcaagcg cgagggcgag 16620
 gatctgtacc ccacatgca gctgatggtg cccaagcgcc agaaghtgga agacgtgctg 16680
 gagaccatga aggtggaccc ggacgtgcag cccgaggtca aggtgcggcc catcaagcag 16740
 gtggcccccg gcntgggctg gcagaccgtg gacatcwaga ttcccacgga gcccatggaa 16800
 acgcagaccg agcccatgat caagcccagc accagcacca tggaggtgca gacggatccc 16860
 tggatgccat cggctcctag tcgaagacc cggcgcaagt acggcgcggc cagcctgctg 16920
 atgcccact acgcgctgca tccttccatc atccccacgc cgggctaccg cggcacgcgc 16980

ttctaccgcg	gtcataccag	cagccgccgc	cgcaagacca	ccactcgccg	ctcgccgctcg	17040
ccgcaccgcc	gctgcaacca	cccctgccgc	cctgggtgcgg	agagtgtacc	gccgcggccg	17100
cgcacctctg	accctgccgc	gcgcgcgcta	ccacccgagc	atcgccattt	aaactttcgc	17160
cagctttgca	gatcaatggc	cctcacatga	ccgccttcgc	gttcccatta	cgggctaccg	17220
aggaagaaaa	ccgcgccgta	gaaggctggc	ggggaacggg	atgcgtcgcc	accaccaccg	17280
gcggcgggcg	gccatcagca	agcggttggg	gggaggcttc	ctgcccgcgc	tgatcccat	17340
catcgccgcg	gcgatcgggg	cgatccccgg	cattgcttcc	gtggcggtgc	aggcctctca	17400
gcgccactga	gacacacttg	gaaacatctt	gtaatagacc	ratggactct	gacgctcctg	17460
gtcctgtgat	gtgttttcgt	agacagatgg	aagacatcaa	tttttcgtcc	ctggctccgc	17520
gacacggcac	gcggccgttc	atgggcacct	ggagcgacat	cggcaccagc	caactgaacg	17580
ggggcgccctt	caattggagc	agtctctgga	gcgggcttaa	gaatttcggg	tccacgctta	17640
aaacctatgg	cagcaaggcg	tggaacagca	ccacagggca	ggcgctgagg	gataagctga	17700
aagagcagaa	cttccagcag	aagggtggtcg	atgggctcgc	ctcgggcatc	aacgggggtg	17760
tggacctggc	caaccaggcc	gtgcagcggc	agatcaacag	ccgcctggac	ccggtgccgc	17820
ccgccggctc	cgtggagatg	ccgcaggtgg	aggaggagct	gcctcccctg	gacaagcggg	17880
gcgagaagcg	accccgcccc	gatgcggagg	agacgctgct	gacgcacacg	gacgagccgc	17940
ccccgtacga	ggaggcggtg	aaactgggtc	tgcccaccac	gcggcccatc	gcgcccctgg	18000
ccaccgggggt	gctgaaaccc	gaaaagcccc	cgaccctgga	cttgcytcct	ccccagcctt	18060
ccgccccatv	tacagtggct	aagcccctgc	cgccggtggc	cgtggcccgc	gcgcgacccg	18120
ggggcaccgc	ccgccctcat	gcgaactggc	agagcactct	gaacagcatc	gtgggtctgg	18180
gagtgcagag	tgtgaagcgc	cgccgctgmt	attaaaccta	ccgtagcgct	taacttgctt	18240
gtctgtgtgt	gtatgtatta	tgtcgccgcc	gcygctgtcc	accagaagga	ggagtgaaga	18300
ggcgcgttgc	cgagttgcra	gatggccacc	ccatcgatgc	tgccccagtg	ggcgtaacatg	18360
cacatcgccg	gacaggacgc	ttcggagtag	ctgagtccgg	gtctggtgca	gtttgcccgc	18420
gccacagaca	cctacttcag	tctggggaac	aagtttagga	acccacgggt	ggcgcccacg	18480
caygatgtga	ccaccgaccg	cagccagcgg	ctgacgctgc	gcttcgtgcc	cgtggaccgc	18540
gaggacaaca	cctactcgta	caaagtgcgc	tacacgctgg	ccgtggggcg	caaccgcgtg	18600
ctggacatgg	ccagcaccta	ctttgacatc	cgcggcgtgc	tggatcgggg	ccctagcttc	18660
aaaccctact	ccggcaccgc	ctacaacagt	ctggccccc	agggagcacc	caacacttgt	18720
cagtggacat	ataaagccga	tggtgaaact	gccacagaaa	aaacctatac	atatggaaat	18780
gcacccgtgc	agggcattaa	catcacaaaa	gatgggtattc	aacttggaac	tgacaccgat	18840

gatcagccaa tctacgcaga taaaacctat cagcctgaac ctcaagtggg tgatgctgaa 18900
tggcatgaca tcaactggtag tgatgaaaag tatggaggca gagctcttaa gcctgatacc 18960
aaaatgaagc cttgttatgg ttcttttgcc aagcctacta ataaagaagg aggtcaggca 19020
aatgtgaaaa caggaacagg cactactaaa gaatatgaca tagacatggc tttctttgac 19080
aacagaagtg cggtgctgc tggcctagct ccagaaattg ttttgtatac tgaaaatgtg 19140
gatttgaaaa ctccagatac ccatattgta tacaagcag gcacagatga cagcagctct 19200
tctattaatt tgggtcagca agccatgccc aacagaccta actacattgg tttcagagac 19260
aactttatcg ggctcatgta ctacaacagc actggcaata tgggggtgct ggccggtcag 19320
gcttctcagc tgaatgctgt ggttgacttg caagacagaa acaccgagct gtcctaccag 19380
ctcttgcttg actctctggg tgacagaacc cggtatttca gtatgtggaa tcaggcgggtg 19440
gacagctatg atcctgatgt gcgcattatt gaaaatcatg gtgtggagga tgaacttccc 19500
aactattgtt tccctctgga tgctgttggc agaacagata cttatcaggg aattaaggct 19560
aatggaactg atcaaaccac atggaccaaa gatgacagt tcaatgatgc taatgagata 19620
ggcaagggtg atccattcgc catggaaatc aacatccaag ccaacctgtg gaggaacttc 19680
ctctacgcca acgtggccct gtacctgccc gactcttaca agtacacgcc ggccaatgtt 19740
accctgccc acaacaccaa cacctacgat tacatgaacg gccgggtggg ggcgccctcg 19800
ctggtggact chtacatcaa catcggggcg cgctggtcgc tggatcccat ggacaacgtg 19860
aacccttca accaccaccg caatgcgggg ctgcgctacc gctccatgct cctgggcaac 19920
gggcgctacg tgcccttcca catccagggtg cccagaaat ttttcgcat caagagcctc 19980
ctgctcctgc ccgggtccta cacctacgag tggaacttcc gcaaggacgt caacatgatc 20040
ctgcagagct ccctcgcaa cgacctgcg acggacgggg cctccatctc cttcaccagc 20100
atcaacctct acgccacctt cttcccatg gcgcacaaca cggcctccac gctcgaggcc 20160
atgctgcgca acgacaccaa cgaccagtcc ttcaacgact acctctcggc ggccaacatg 20220
ctctacccca tcccgccaa cgccaccaac gtgcccattc ccatcccctc gcgcaactgg 20280
gccgccttcc gcggctggtc cttcacgct ctcaagacca aggagacgcc ctcgctgggc 20340
tccgggttcg acccctactt cgtctactcg ggctccatcc cctacctcga cggcaccttc 20400
tacctcaacc acaccttcaa gaaggtctcc atcaccttcg actcctccgt cagctggccc 20460
ggcaacgacc ggctcctgac gcccaacgag ttcgaaatca agcgcaccgt cgacggcgag 20520
ggctacaacg tggcccagt caacatgacc aaggactgg tccctggcca gatgctggcc 20580
cactacaaca tcggctacca gggcttctac gtgcccagg gctacaagga ccgcatgtac 20640
tccttcttcc gcaacttcca gcccatgagc cgccagggtg tggacgaggt caactacaag 20700

gactaccagg ccgtcaccct ggcctaccag cacaacaact cgggcttcgt cggctacctc 20760
 gcgcccacca tgcgccaggg ccagccmtac cccgccaamt acccmtaccc gctcatcggc 20820
 aagagcgccg tcaccagcgt caccagaaaa aagttcctct gcgacagggg catgtggcgc 20880
 atcccccttct ccagcaactt catgtccatg ggcgcgctca ccgacctcgg ccagaacatg 20940
 ctctatgcca actccgcccc cgcgctagac atgaatttcg aagtcgaccc catggatgag 21000
 tccacccttc tctatgttgt ctctgaagtc ttcgacgtcg tccgagtgcg ccagccccac 21060
 cgcgcgctca tcgaggccgt ctacmtgcgc acccccttct cggccggtaa cgccaccacc 21120
 taagctcttg cttcttgcaa gccatggccg cgggctccgg cgagcaggag ctcagggcca 21180
 tcatccgcga cctgggctgc gggccmtact tcctgggcac sttcgataag cgcttcccgg 21240
 gattcatggc cccgcacaag ctggcctgcg ccacgtgcaa cagggccggc cgcgagaccg 21300
 ggggcgagca ctggctggcc ttcgcctgaa cccgcgctcg aacacctgct acctcttcga 21360
 ccccttcggg ttctcggacg agcgctcaa gcagatctac cagttcgagt acgagggcct 21420
 gctgcgccgc agcgccctgg ccaccgagga ccgctgcgtc accctggaaa agtccacca 21480
 gaccgtgcag ggtccgcgct cggccgcctg cgggctcttc tgctgcatgt tcctgcacgc 21540
 cttcgtgcac tggcccgaac gccccatgga caagaacccc accatgaact tgctgacggg 21600
 ggtgcccac ggcatgctcc agtcgccccca ggtggaaccc accctgcgcc gcaaccagga 21660
 ggcgctytac cgcttctca actcccactc cgcmactttt cgctcccacc gcgcgcgcgt 21720
 cgagaaggcc accgccttcg accgcatgaa tcaagacatg taaaccgtgt gtgtatgtta 21780
 aatgtcttta ataaacagca ctttcatgtt acacatgcat ctgagatgat ttatttagaa 21840
 atcsaaaggg ttcttccggg tctcggcatg gcccgcgggc agggacacgt tgcggaactg 21900
 gtacttggcc agccacttga actcggggat cagcagtttg ggcagcgggg tgctcggggaa 21960
 ggagtcggtc cacagcttcc gcgtcagttg cagggcgccc agcaggtcgg gcgcggagat 22020
 cttgaaatcg cagttgggac ccgcgttctg cgcgcgggag ttgcggtaca cgggggttgc 22080
 gcactggaac accatcaggg ccgggtgctt cacgctcgcc agcaccgtcg cgtcggtgat 22140
 gctctccacg tcgaggtcct cggcgttggc catcccgaag ggggtcatct tgcaggtctg 22200
 ccttcccatg gtgggcacgc acccgggctt gtggttgcaa tcgcagtgcg gggggatcag 22260
 catcatctgg gcctggtcgg cgttcatccc cgggtacatg gccttcatga aagcctccaa 22320
 ttgcctgaac gcctgctggg ccttggctcc ctcggtgaag aagacccgc aggacttgct 22380
 agagaactgg ttggtggcgc acccggcgtc gtgcacgcag cagcgcgct cgttgttggc 22440
 cagctgcacc acgctgcgcc ccagcggtt ctgggtgatc ttggcccggt cgggggttctc 22500
 cttcagcgcg cgctgcccg tctcgcctgc cacatccatc tcgatcatgt gtccttctg 22560

gatcatggtg	gtcccggtgca	ggcaccgcag	cttgccctcg	gcctcggtgc	acccggtgcag	22620
ccacagcgcg	cacccggtgc	actcccagtt	cttggtggcg	atctgggaat	gcgcgtgcac	22680
gaagccctgc	aggaagcggc	ccatcatggt	ggtcagggtc	ttgttgctag	tgaaggtcag	22740
cggaatgccg	cgggtgctcct	cgttgatgta	caggtggcag	atgcggcggt	acacctcgcc	22800
ctgctcgggc	atcagctgga	agttggcttt	caggtcggtc	tccacgcggt	agcgggtccat	22860
cagcatagtc	atgattttcca	tacccttctc	ccaggccgag	acgatgggca	ggctcatagg	22920
gttcttcacc	atcatcttag	cgctagcagc	cgcgccagc	gggtcgctct	cgtccagggt	22980
ctcaaagctc	cgcttgccgt	ccttctcggt	gatccgcacc	ggggggtagc	tgaagcccac	23040
ggccgccagc	tcctcctcgg	cctgtctttc	gtcctcgctg	tcctggctga	cgtcctgcag	23100
gaccacatgc	ttggtcttgc	ggggtttctt	cttgggcggc	agcggcggcg	gagatggttg	23160
agatggcgag	ggggagcgcg	agttctcgct	caccactact	atctcttctt	cttcttggtc	23220
cgaggccacg	cggcggtagg	tatgtctctt	cgggggcaga	ggcggaggcg	acgggctctc	23280
gccgccgcga	cttggcggat	ggctggcaga	gccccttcgg	cgttcggggg	tgcgctcccg	23340
gcggcgctct	gactgacttc	ctccgcggcc	ggccattgtg	ttctcctagg	gaggaacaac	23400
aagcatggag	actcagccat	cgccaacctc	gccatctgcc	cccaccgccg	acgagaagca	23460
gcagcagcag	aatgaaagct	taaccgcccc	gccgcccagc	cccgccacct	ccgacgcggc	23520
cgtcccagac	atgcaagaga	tggaggaatc	catcgagatt	gacctgggct	atgtgacgcc	23580
cgcggagcac	gaggaggagc	tggcagtgcg	cttttcacaa	gaagagatac	accaagaaca	23640
gccagagcag	gaagcagaga	atgagcagag	tcaggctggg	ctcgagcatg	acggcgacta	23700
cctccacctg	agcggggggg	aggacgcgct	catcaagcat	ctggcccggc	aggccaccat	23760
cgtcaaggat	gcgctgctcg	accgcaccga	ggtgcccctc	agcgtggagg	agctcagccg	23820
cgcctacgag	ttgaacctct	tctcgccgcg	cgtgcccccc	aagcgccagc	ccaatggcac	23880
ctgcgagccc	aaccgcgccc	tcaacttcta	cccggctctt	gcgggtgccc	aggccctggc	23940
cacctaccac	atcttttttca	agaaccaaaa	gatccccgtc	tcctgccgcg	ccaaccgcac	24000
ccgcgccgac	gcccttttca	acctgggtcc	cggcgcccgc	ctacctgata	tcgcctcctt	24060
ggaagagggt	cccaagatct	tcgagggtct	gggcagcgac	gagactcggg	ccgcgaacgc	24120
tctgcaagga	gaaggaggag	agcatgagca	ccacagcgcc	ctggtcgagt	tggaaggcga	24180
caacgcgcgg	ctggcggtgc	tcaaacgcac	ggtcgagctg	acccatttcg	cgtacccggc	24240
tctgaacctg	ccccccaaag	tcatgagcgc	ggtcatggac	caggtgctca	tcaagcgcg	24300
gtcgcccatc	tccgaggacg	agggcatgca	agactccgag	gagggcaagc	ccgtgggtcag	24360
cgacgagcag	ctggcccggg	ggctgggtcc	taatgctagt	ccccagagtt	tggaagagcg	24420

gcgcaaaactc atgatggccg tggctcctggt gaccgtggag ctggagtgcc tgcgccgctt 24480
 cttcgccgac gcggagaccc tgcgcaaggt cgaggagaac ctgcactacc tcttcaggca 24540
 cgggttcgtg cgccaggcct gcaagatctc caacgtggag ctgaccaacc tggctctcgta 24600
 catgggcatc ttgcacgaga accgcctggg gcagaacgtg ctgcacacca ccctgcgcgg 24660
 ggaggcccg cgcgactaca tccgcgactg cgtctacctc tacctctgcc acacctggca 24720
 gacgggcatg ggcgtgtggc agcagtgtct ggaggagcag aacctgaaag agctctgcaa 24780
 gctcctgcag aagaactcaa gggctctgtg accgggttcg acgagcgcac caccgcctcg 24840
 gacctggccg acctcathtt ccccgagcgc ctgaggctga cgctgcgcaa cggcctgccc 24900
 gactttatga gccaaagcat gttgcaaaac tttcgctctt tcatcctcga acgctccgga 24960
 atcctgcccg ccacctgctc cgcgctgccc tcggacttcg tgccgctgac cttccgcgag 25020
 tgccccccgc cgctgtggag ccactgctac ctgctgcgcc tggccaacta cctggcctac 25080
 cactcggacg tgatcgagga cgtcagcggc gagggcctgc tcgagtgcc a ctgccgctgc 25140
 aacctctgca cgccgcaccg ctccctggcc tgcaaccccc agctgytgag cgagaccag 25200
 atcatcggca ccttcgagtt gcaagggccc agcgaaggcg agggttcagc cgccaagggg 25260
 ggtctgaaac tcaccccggg gctgtggacc tcggcctact tgcgcaagtt cgtgcccag 25320
 gactaccatc ccttcgagat caggttctac gaggaccaat cccatccgcc caaggccgag 25380
 ctgtcggcct gcgtcatcac ccagggggcg atcctggccc aattgcaagc catccagaaa 25440
 tccccccaag aattcttgct gaaaaagggc cgcggggtct acctcgacct ccagaccggt 25500
 gaggagctca accccggctt cccccaggat gccccgagga aacaagaagc tgaaagtgga 25560
 gctgccgccc gtggaggatt tggaggaaga ctgggagaac agcagtcagg cagaggagga 25620
 ggagatggag gaagactggg acagcactca ggcagaggag gacagcctgc aagacagtct 25680
 ggaggaagac gaggaggagg cagaggagga ggtggaagaa gcagccgccg ccagaccgtc 25740
 gtcctcggcg ggggagaaa caagcagcac ggataccatc tccgctccgg gtcgggggtc 25800
 cgctcgacca cacagtagat gggacgagac cggacgattc ccgaaccca ccaccagac 25860
 cggtaaagaag gagcggcagg gatacaagtc ctggcggggg caaaaaacg ccatcgtctc 25920
 ctgcttgca gctgcgggg gcaacatctc cttaccccg cgctacctgc tcttccaccg 25980
 cgggggtgaac tttccccga acatcttgca ttactaccgt cacctccaca gcccctacta 26040
 cttccaagaa gaggcagcag cagcagaaaa agaccagcag aaaaccagca gctagaaaat 26100
 ccacagcggc ggcagcaggt ggactgagga tcgcggcgaa cgagccggcg caaaccggg 26160
 agctgaggaa ccggatcttt cccaccctct atgccatctt ccagcagagt cgggggcagg 26220
 agcaggaact gaaagtcaag aaccgttctc tgcgctcgct caccgcagt tgtctgtatc 26280

acaagagcga agaccaactt cagcgactc tcgaggacgc cgaggctctc ttcaacaagt 26340
 actgcgcgct cactcttaaa gagtagcccg cgcccgcca gtcgcagaaa aaggcgggaa 26400
 ttacgtcacc tgtgcccttc gccctagccg cctccacca tcatcatgag caaagagatt 26460
 cccacgcctt acatgtggag ctaccagccc cagatgggccc tggccgcggtg tgccgcccag 26520
 gactactcca cccgcatgaa ttggctcagc gccggggccc cgatgatctc acgggtgaat 26580
 gacatccgcg cccaccgaaa ccagatactc ctagaacagt cagcgctcac cgccacgccc 26640
 cgcaatcacc tcaatccgcg taattggccc gccgccttggtgtgtaccagga aattccccag 26700
 cccacgaccg tactacttcc gcgagacgcc caggccgaag tccagctgac taactcaggt 26760
 gtccagctgg cgggcggcgc caccctgtgt cgtcaccgcc ccgctcaggg tataaagcgg 26820
 ctggtgatcc ggggcagaag cacacagctc aacgacgaag tggtagctc ttcgctgggt 26880
 ctgcgacctg acggagtctt ccaactcgcc ggatcgggga gatcttcctt cagcctcgt 26940
 caggccgtcc tgactttgga gagttcgtcc tcgcagcccc gctcgggtgg catcggcact 27000
 ctccagttcg tggaggagtt cactccctcg gtctacttca accccttctc cggtccccc 27060
 ggccactacc cggacgagtt catcccgaac ttcgacgcca tcagcgagtc ggtggacggc 27120
 tacgattgaa tgtcccatgg tggcgcagct gacctagctc ggcttcgaca cctggaccac 27180
 tgccgcgct tccgctgctt cgctcgggat ctcgccgagt ttgcctactt tgagctgccc 27240
 gaggagcacc ctcagggccc ggccacgga gtgcggatcg tcgtcgaagg gggcctcgac 27300
 tcccactgc ttcggatctt cagccagcgt ccgatcctgg tcgagcgca gcaaggacag 27360
 acccttctga ctctgtactg catctgcaac caccggccc tgcatgaaag tctttgttgt 27420
 ctgctgtgta ctgagtataa taaaagctga gacagcgact actccggact tccgtgtgtt 27480
 cctgaatcca tcaaccagtc tttgttcttc accgggaacg agaccgagct ccagctccag 27540
 tgtaagcccc acaagaagta cctcacctgg ctgttccagg gctccccgat cgccgttgtc 27600
 aaccactgcg acaacgacgg agtcctgstg agcggccctg ccaaccwtac tttttccacc 27660
 cgcagaagca agctccagct sttccaaccc ttcctccccg ggacctatca gtgcgtctcg 27720
 ggaccctgcc atcacacctt ccacctgatc ccgaatacca cagcgctcgt ccccgmtact 27780
 aacaaccaa ctaacctcca ccaacgccac cgtcgcgacc tttctgaatc taatactacc 27840
 acccacaccg gaggtgagct ccgaggtcaa ccaacctctg ggatttacta cgccccctgg 27900
 gaggtgggtg ggtaataaac gctaggccta gttgcgggtg ggcttttggt tctctgctac 27960
 ctatacctcc cttgctgttc gtacttagtg gtgctgtgtt gctggtttaa gaaatgggga 28020
 agatcacctc agtgagctgc ggtgcgctgg tggcgggtgt gctttcgatt gtgggactgg 28080
 gcggtgcggc tgtantgaaa gagaaggccg atccctgctt gcatttcaat cccaacaaat 28140

gccagctgag ttttcagccc gatggcaatc ggtgcgcggt actgatcaag tgcggatggg 28200
 aatgcgagaa cgtgagaatc gagtacaata acaagactcg gaacaatact ctcgcgtccg 28260
 tgtggcagcc cggggacccc gagtgggtaca ccgtctctgt ccccggtgct gacggctccc 28320
 cgcgcacccgt gaataatact ttcattttttg cgcacatgtg cgacacgggtc atgtggatga 28380
 gcaagcagta cgatatgtgg cccccacga aggagaacat cgtggtcttc tccatcgctt 28440
 acagcctgtg cacggcgcta atcaccgcta tcgtgtgcct gagcattcac atgctcatcg 28500
 ctattcgccc cagaaataat gccgaaaaag aaaaacagcc ataacgtttt ttttcacacc 28560
 tttttcagac catggcctct gttaaatttt tgcttttatt tgccagtctc attgccgtca 28620
 ttcattggaat gagtaatgag aaaattacta tttacactgg cactaatcac acattgaaag 28680
 gtccagaaaa agccacagaa gtttcatggt attgttattt taatgaatca gatgtatcta 28740
 ctgaactctg tggaacaat aacaaaaaaa atgagagcat tactctcatc aagtttcaat 28800
 gtggatctga cttaacccta attaacatca ctagagacta tgtaggtatg tattatggaa 28860
 ctacagcagg catttcggac atggaatttt atcaagtttc tgtgtctgaa cccaccacgc 28920
 ctagaatgac cacaaccaca aaaactacac ctgttaccac tatgcagctc actaccaata 28980
 acatttttgc catgcgtaa atggtcaaca atagcactca acccaccaca cccagtgagg 29040
 aaattcccaa atccatgatt ggcattattg ttgctgtagt ggtgtgcatg ttgatcatcg 29100
 ccttgtgcat ggtgtactat gccttctgct acagaaagca cagactgaac gacaagctgg 29160
 aacacttact aagtgttgaa ttttaatttt ttagaaccat gaagatccta ggccttttaa 29220
 ttttttctat cattacctct gctctatgca attctgacaa tgaggacgtt actgtcgttg 29280
 tcggatcaaa ttatacactg aaaggtccag cgaagggtat gctttcgtgg tattgctatt 29340
 ttggatctga cactacagaa actgaattat gcnatcttaa gaatggcaaa attcaaaatt 29400
 cttaaaatta acaattatat atgcaatggg actgatctga tactcctcaa tatcacgaaa 29460
 tcatatgstg gcagttacac ctgccctgga gatgatgctg acagtatgat tttttacaaa 29520
 gtaactgttg ttgatcccat actccacctc caccaccac aattactcac accacacaca 29580
 cagatcaaac cgagcagag gaggcagcaa agttagcctt gcagggtcaa gacagttcat 29640
 ttgttggcat taccctaca catgatcagc ggtgtccggg gctgctagtc agcggcattg 29700
 tcggtgtgct ttcgggatta gcagtcataa tcatctgcat gttcattttt gcttctgct 29760
 atagaaggct ttaccgacaa aaatcagacc cactgctgaa cctctatggt taattttttc 29820
 cagagtcatg aaggcagtta gcgctctagt tttttgttct wtgattggca ttgttttttg 29880
 caatcctatt cctaaagtta gctttattaa agatgtgaat gttactgagg ggggcaatgt 29940
 gacactggta ggtgtagagg gtgctgaaaa caccacctgg acaaaatacc acctcaatgg 30000

gtggaaagat atttgcaatt ggagtgtatt agtttataca tgtgagggag ttaatcttac 30060
 cattgtcaat gccacctcag ctcaaaatgg tagaattcaa ggacaaagtg tcagtgtatc 30120
 taatgggtat ttacccaac atacttttat ctatgacgtt aaagtcatac cactgcwtac 30180
 gcttagccca cttagcatta ccacacagac aaccacatt acacagacaa ccacatacag 30240
 tacattaaat cagcbtacca ccactacagc agcagagggt gccagctcgt ctgggggtccg 30300
 agtggcattt ttgatgtggg ccccatmtag cagtcccact gctagtacca atgagcagac 30360
 tactgaattt ttgtccactg tcgagagcca caccacagct acctccagt ccttctctag 30420
 caccgccaat ctctcctcgc tttcctntac accaatcagt cccgytaata ctcttagccc 30480
 cgtcctcttc ccactcccct gaagcaaaca gacggcggca tgcaatggca gatcaccctg 30540
 ctcatgtga tcgggttggc catcctggcc gtgttgctct actacatctt ctgccgccgc 30600
 attcccaacg cgcaccgcaa gccggtatac aagcccatca ttgtcgggca gccggagccg 30660
 cttcaggtgg aaggggtct aaggaatctt ctcttctctt ttacagtatg gtgattgaac 30720
 tatgattcct agacaattct tgatcactat tcttatctgc ctctccaag tctgtgccac 30780
 cctcgtctg gtggccaacg ccagtccaga ctgtattggg cccttcgcct cctacgtgct 30840
 ctttgccctc accacctgca tctgctgctg tagcatagtc tgcctgctta tcaccttctt 30900
 ccagttcatt gactggatct ttgtgcgcat cgcmtacctg cgccaccacc cccagtaccg 30960
 cgaccagcga gtggcgcggc tgctcaggct cctctgataa gcatgcgggc tgtgntactt 31020
 ctgcgccttc tgctgttagt gctccccctg cccgtcgacc cccggtcccc caccagtc 31080
 cccgaggagg tccgcaaagt caaatccaa gaacctgga aattcctcaa atgctaccgc 31140
 caaaaatcag acatgcatcc cagctggatc atgatcattg ggatcgtgaa cattctggcc 31200
 tgcacctca tctcctttgt gatttaccct tgctttgact ttggttgga ctcgccagag 31260
 gcgctctatc tccgcctga acctgacaca ccaccacagc aacctcaggc acacgcacta 31320
 ccaccactac agcctaggcc acaatacatg cccatattag actatgaggc cgagccacag 31380
 cgacctatgc tccccgctat tagttacttc aatctaaccg gcggagatga ctgacctact 31440
 ggccaacaac aacgtcaacg accttctcct ggacatggac ggccgcgcct cggagcagcg 31500
 actcgcccaa cttcgcatc gccagcagca ggagagagcc gtcaaggagc tgcaggatgc 31560
 ggtggccatc caccagtgc agagaggcat cttctgcctg gtgaaacagg ccaagatctc 31620
 ctacgaggtc actccaaacg accatcgctt ctctacgag ctctgcagc agcgccagaa 31680
 gttcacctgc ctggtcggag tcaaccccat cgtcatcacc cagcagtctg gcgataccaa 31740
 ggggtgcac cactgctcct gcgactcccc cgactgcgtc cacactctga tcaagaccct 31800
 stgcggcctc cgcgacctcc tccccatgaa ctaatcacc ccttatccag tgaaataaag 31860

atcatattga tgatgatttt acagaaataa aaaataatca tttgatttga aataaagata 31920
 caatcatatt gatgatttga gtttaacaaa aaaataaaga atcacttact tgaaatctga 31980
 taccaggtct ctgtccatgt tttctgcaa caccacttca ctcccctctt cccagctctg 32040
 gtactgcagg ccccggcggg ctgcaaactt cctccacacg ctgaagggga tgtcaaattc 32100
 ctctgtccc tcaatcttca ttttatcttc tatcagatgt ccaaaaagcg cgtccgggtg 32160
 gatgatgact tcgaccccggt ctacccttac gatgcagaca acgcaccgac cgtgcccttc 32220
 atcaaccccc ccttcgtctc ttcagatgga ttccaagaga agcccctggg ggtgttggtc 32280
 ctgcgactgg ccgaccccggt caccaccaag aacggggaaa tcaccctcaa gctgggagag 32340
 ggggtggacc tcgattcctc gggaaaactc atctccaaca cggccaccaa ggccgcccgc 32400
 cctctcagtt tttccaacaa caccatttcc cttaacatgg atcaccctt ttactactaa 32460
 gatggaaaat tatccttaca agtttctcca ccattaaata tactgagaac aagcattcta 32520
 aacacactag ctttaggttt tggatcaggt ttaggactcc gtggctctgc cttggcagta 32580
 cagttagtct ctccacttac atttgatact gatggaaaca taaagcttac cttagacaga 32640
 ggtttgcatg ttacaacagg agatgcaatt gaaagcaaca taagctgggc taaaggttta 32700
 aaatttgaag atggagccat agcaaccaac attggaaatg ggtagagtt tggaagcagt 32760
 agtacagaaa caggtgttga tgatgcttac ccaatccaag ttaaacttgg atctggcctt 32820
 agctttgaca gtacaggagc cataatggct ggtaacaaag aagacgataa actcactttg 32880
 tggacaacac ctgatccatc accaaactgt caaatactcg cagaaaatga tgcaaaacta 32940
 acactttgct tgactaaatg tggtagtcaa atactggcca ctgtgtcagt cttagtgtga 33000
 ggaagtggaa acctaaaccc cattactggc accgtaagca gtgctcaggt gtttctacgt 33060
 tttgatgcaa acggtgttct tttacagaa cattctacac taaaaaata ctgggggtat 33120
 aggcagggag atagcataga tggcactcca tataccaatg ctgtaggatt catgcccaat 33180
 ttaaaagctt atccaaagtc acaaagttct actactaaaa ataatatagt agggcaagta 33240
 tacatgaatg gagatgtttc aaaacctatg cttctcacta taaccctcaa tggtagtgat 33300
 gacagcaaca gtacatatc aatgtcattt tcatacacct ggactaatgg aagctatggt 33360
 ggagcaacat ttggggctaa ctcttatacc ttctcataca tcgcccaga atgaacactg 33420
 tatcccaccc tgcattgcaa cccttcccac cccactctgt ggaacaaact ctgaaacaca 33480
 aaataaaata aagttcaagt gttttattga ttcaacagtt ttacaggatt cgagcagtta 33540
 ttttctctc accctcccag gacatggaat acaccacct ctccccccgc acagccttga 33600
 acatctgaat gccattggtg atggacatgc ttttggctc caggttccac acagtttcag 33660
 agcgagccag tctcgggtcg gtcagggaga tgaaaccctc cgggcactcc cgcattctga 33720

cctcacagct caacagctga ggattgtcct cgggtggtcgg gatcacggtt atctggaaga 33780
agcagaagag cggcgggtggg aatcatagtc cgcgaacggg atcggccggg ggtgtcgcat 33840
caggccccgc agcagtcgct gccgccgccg ctccgtcaag ctgctgctca ggggggtccgg 33900
gtccagggac tccctcagca tgatgccac ggccctcagc atcagtcgtc tgggtgcggcg 33960
ggcgcagcag cgcattgcgga tctcgtctcag gtcgtctcag tacgtgcaac acagaaccac 34020
cagggttggtc aacagtccat agttcaacac gctccagccg aaactcatcg cgggaaggat 34080
gctaccacacg tggccgtcgt accagatcct caggtaaatac aagtgggtgcc ccctccagaa 34140
cacgctgccc acgtacatga tctccttggg catgtggcgg ttcaccacct cccggtacca 34200
catcacccctc tgggtgaaca tgcagccccg gatgatcctg cggaaccaca gggccagcac 34260
cgccccgccc gccatgcagc gaagagaccc cgggtcccgg caatggcaat ggaggacca 34320
ccgctcgtac ccgtggatca tctgggagct gaacaagtct atgttggcac agcacaggca 34380
tatgctcatg catctcttca gcaactctca ctcctcgggg gtcaaaacca tatcccaggg 34440
cacggggaac tcttgcagga cagcgaaccc cgcagaacag ggcaatcctc gcacagaact 34500
tacattgtgc atggacaggg tatcgcaatc aggcagcacc ggggtgatcct ccaccagaga 34560
agcgcgggtc tcggtctcct cacagcgtgg taagggggcc ggccgatacg ggtgatggcg 34620
ggacgcggct gatcgtgttc gcgaccgtgt catgatgcag ttgctttcgg acattttcgt 34680
acttgctgta gcagaacctg gtccggggcg tgcacaccga tcgccggcg cgggtctcggc 34740
gcttggaacg ctcggtgttg aaattgtaaa acagccactc tctcagaccg tgcagcagat 34800
ctagggcctc aggagtgatg aagatcccat catgcctgat ggctctgatc acatcgacca 34860
ccgtggaatg ggccagaccc agccagatga tgcaattttg ttgggtttcg gtgacggcgg 34920
gggaggggaag aacaggaaga accatgatta acttttaatac caaacggtct cggagtactt 34980
caaaatgaag atcgcggaga tggcacctct cgcgcccgct gtgttggtgg aaaataacag 35040
ccaggtaaaa ggtgatacgg ttctcgagat gttccacggt ggcttccagc aaagcctcca 35100
cgcgcacatc cagaaacaag acaatagcga aagcgggagg gttctctaata tcctcaatca 35160
tcatgttaca ctctgcacc atccccagat aattttcatt tttccagcct tgaatgattc 35220
gaactagttc gtgaggtaaa tccaagccag ccatgataaa gagctcgcgc agagcgccct 35280
ccaccggcat tcttaagcac accctcataa ttccaagata ttctgctcct gggttcacctg 35340
cagcagattg acaagcggaa tatcaaaatc tctgccgcga tccctgagct cctccctcag 35400
caataactgt aagtactctt tcatatcctc tccgaaatct ttagccatag gaccaccagg 35460
aataagatta gggcaagcca cagtacagat aaaccgaagt cctccccagt gagcattgcc 35520
aaatgcaaga ctgctataag catgctggct agaccgggtg atatcttcca gataactgga 35580

```

cagaaaatcg cccaggcaat ttttaagaaa atcaacaaaa gaaaaatcct ccagggtggac 35640
gttttagagcc tcgggaacaa cgatgaagta aatgcaagcg gtgcgttcca gcatgggtag 35700
ttagctgatc ttagaaaaaa acaaaaatga acattaaacc atgctagcct ggcgaaacagg 35760
tgggtaaaatc gttctctcca gcaccaggca ggccacgggg tctccggcgc gaccctcgta 35820
aaaattgtcg ctatgattga aaaccatcac agagagacgt tcccgggtggc cggcgtgaat 35880
gattcgacaa gatgaatata cccccggaac attggcgtcc gcgagtgaaa aaaagcgccc 35940
gaggaagcaa taaggcacta caatgctcag tctcaagtcc agcaaagcga tgccatgcgg 36000
atgaagcaca aaattctcag gtgcgtacaa aatgtaatta ctcccctcct gcacaggcag 36060
caaagccccc gatccctcca ggtacacata caaagcctca gcgtccatag cttaccgagc 36120
agcagcacac aacaggcgca agagtcagag aaaggctgag ctctaacctg tccacccgct 36180
ctctgctcaa tatatagccc agatctacac tgacgtaaag gccaaagtct aaaaataccc 36240
gccaaataat cacacacgcc cagcacacgc ccagaaaccg gtgacacact caaaaaata 36300
cgcgcacttc ctcaaagcc caaaactgcc gtcatttccg ggttcccacg ctacgtcatc 36360
aaaacacgac tttcaaattc cgtcgaccgt taaaacgtc acccgccccg cccctaacgg 36420
tcgcccgtct ctgagccaat cagcgccccg catccccaaa ttcaaacacc tcatttgcac 36480
attaacgcgc aaaaaagtt tgaggtatat tattgatgat g 36521

```

```

<210> 34
<211> 314
<212> PRT
<213> Human adenovirus type 4

```

```

<400> 34

```

```

Asn Thr Cys Gln Trp Lys Asp Ser Asp Ser Lys Met His Thr Phe Gly
1           5           10           15
Ala Ala Ala Met Pro Gly Val Thr Gly Lys Lys Ile Glu Ala Asp Gly
20           25           30
Leu Pro Ile Arg Ile Asp Ser Thr Ser Gly Thr Asp Thr Val Ile Tyr
35           40           45
Ala Asp Lys Thr Phe Gln Pro Glu Pro Gln Val Gly Asn Asp Ser Trp
50           55           60
Val Asp Thr Asn Gly Ala Glu Glu Lys Tyr Gly Gly Arg Ala Leu Lys
65           70           75           80
Asp Thr Thr Lys Met Asn Pro Cys Tyr Gly Ser Phe Ala Lys Pro Thr
85           90           95
Asn Lys Glu Gly Gly Gln Ala Asn Leu Lys Asp Ser Glu Pro Ala Ala
100          105          110

```


Thr Thr Pro Asn Tyr Asp Ile Asp Leu Ala Phe Phe Asp Ser Lys Thr
 115 120 125
 Ile Val Ala Asn Tyr Asp Pro Asp Ile Val Met Tyr Thr Glu Asn Val
 130 135 140
 Asp Leu Gln Thr Pro Asp Thr His Ile Val Tyr Lys Pro Gly Thr Glu
 145 150 155 160
 Asp Thr Ser Ser Glu Ser Asn Leu Gly Gln Gln Ala Met Pro Asn Arg
 165 170 175
 Pro Asn Tyr Ile Gly Phe Arg Asp Asn Phe Ile Gly Leu Met Tyr Tyr
 180 185 190
 Asn Ser Thr Gly Asn Met Gly Val Leu Ala Gly Gln Ala Ser Gln Leu
 195 200 205
 Asn Ala Val Val Asp Leu Gln Asp Arg Asn Thr Glu Leu Ser Tyr Gln
 210 215 220
 Leu Leu Leu Asp Ser Leu Gly Asp Arg Thr Arg Tyr Phe Ser Met Trp
 225 230 235 240
 Asn Gln Ala Val Asp Ser Tyr Asp Pro Asp Val Arg Ile Ile Glu Asn
 245 250 255
 His Gly Val Glu Asp Glu Leu Pro Asn Tyr Cys Phe Pro Leu Asn Gly
 260 265 270
 Val Gly Leu Thr Asp Thr Tyr Gln Gly Val Lys Val Lys Thr Asp Ala
 275 280 285
 Gly Ser Glu Lys Trp Asp Lys Asp Asp Thr Thr Val Ser Asn Ala Asn
 290 295 300
 Glu Ile His Val Gly Asn Pro Phe Ala Met
 305 310

<210> 35
 <211> 318
 <212> PRT
 <213> Human adenovirus type 16

<400> 35

Asn Thr Cys Gln Trp Lys Asp Ser Asp Ser Lys Met His Thr Phe Gly
 1 5 10 15
 Val Ala Ala Met Pro Gly Val Thr Gly Lys Lys Ile Glu Ala Asp Gly
 20 25 30
 Leu Pro Ile Gly Ile Asp Ser Thr Ser Gly Thr Asp Thr Val Ile Tyr
 35 40 45
 Ala Asp Lys Thr Phe Gln Pro Glu Pro Gln Val Gly Asn Ala Ser Trp
 50 55 60
 Val Asp Ala Asn Gly Thr Glu Glu Lys Tyr Gly Gly Arg Ala Leu Lys
 65 70 75 80

Asp Thr Thr Lys Met Lys Pro Cys Tyr Gly Ser Phe Ala Lys Pro Thr
 85 90 95
 Asn Lys Glu Gly Gly Gln Ala Asn Leu Lys Asp Ser Glu Thr Ala Ala
 100 105 110
 Thr Thr Pro Asn Tyr Asp Ile Asp Leu Ala Phe Phe Asp Asn Lys Asn
 115 120 125
 Ile Ala Ala Asn Tyr Asp Pro Asp Ile Val Met Tyr Thr Glu Asn Val
 130 135 140
 Asp Leu Gln Thr Pro Asp Thr His Ile Val Tyr Lys Pro Gly Thr Glu
 145 150 155 160
 Asp Thr Ser Ser Glu Ser Asn Leu Gly Gln Gln Ala Met Pro Asn Arg
 165 170 175
 Pro Asn Tyr Ile Gly Phe Arg Asp Asn Phe Ile Gly Leu Met Tyr Tyr
 180 185 190
 Asn Ser Thr Gly Asn Met Gly Val Leu Ala Gly Gln Ala Ser Gln Leu
 195 200 205
 Asn Ala Val Val Asp Leu Gln Asp Arg Asn Thr Glu Leu Ser Tyr Gln
 210 215 220
 Leu Leu Leu Asp Ser Leu Gly Asp Arg Thr Arg Tyr Phe Ser Met Trp
 225 230 235 240
 Asn Gln Ala Val Asp Ser Tyr Asp Pro Asp Val Arg Ile Ile Glu Asn
 245 250 255
 His Gly Val Glu Asp Glu Leu Pro Asn Tyr Cys Phe Pro Leu Asn Gly
 260 265 270
 Val Gly Phe Thr Asp Thr Tyr Gln Gly Val Lys Val Lys Thr Asp Ala
 275 280 285
 Val Ala Gly Thr Ser Gly Thr Gln Trp Asp Lys Asp Asp Thr Thr Val
 290 295 300
 Ser Thr Ala Asn Glu Ile His Gly Gly Asn Pro Phe Ala Met
 305 310 315

<210> 36

<211> 323

<212> PRT

<213> Human adenovirus type 3

<400> 36

Asn Thr Ser Gln Trp Ile Val Thr Thr Asn Gly Asp Asn Ala Val Thr
 1 5 10 15
 Thr Thr Thr Asn Thr Phe Gly Ile Ala Ser Met Lys Gly Gly Asn Ile
 20 25 30
 Thr Lys Glu Gly Leu Gln Ile Gly Lys Asp Ile Thr Thr Thr Glu Gly
 35 40 45

Glu Glu Lys Pro Ile Tyr Ala Asp Lys Thr Tyr Gln Pro Glu Pro Gln
 50 55 60
 Val Gly Glu Glu Ser Trp Thr Asp Thr Asp Gly Thr Asn Glu Lys Phe
 65 70 75 80
 Gly Gly Arg Ala Leu Lys Pro Ala Thr Asn Met Lys Pro Cys Tyr Gly
 85 90 95
 Ser Phe Ala Arg Pro Thr Asn Ile Lys Gly Gly Gln Ala Lys Asn Arg
 100 105 110
 Lys Val Lys Pro Thr Thr Glu Gly Gly Val Glu Thr Glu Glu Pro Asp
 115 120 125
 Ile Asp Met Glu Phe Phe Asp Gly Arg Asp Ala Val Ala Gly Ala Leu
 130 135 140
 Ala Pro Glu Ile Val Leu Tyr Thr Glu Asn Val Asn Leu Glu Thr Pro
 145 150 155 160
 Asp Ser His Val Val Tyr Lys Pro Glu Thr Ser Asn Asn Ser His Ala
 165 170 175
 Asn Leu Gly Gln Gln Ala Met Pro Asn Arg Pro Asn Tyr Ile Gly Phe
 180 185 190
 Arg Asp Asn Phe Val Gly Leu Met Tyr Tyr Asn Ser Thr Gly Asn Met
 195 200 205
 Gly Val Leu Ala Gly Gln Ala Ser Gln Leu Asn Ala Val Val Asp Leu
 210 215 220
 Gln Asp Arg Asn Thr Glu Leu Ser Tyr Gln Leu Leu Leu Asp Ser Leu
 225 230 235 240
 Gly Asp Arg Thr Arg Tyr Phe Ser Met Trp Asn Gln Ala Val Asp Ser
 245 250 255
 Tyr Asp Pro Asp Val Arg Ile Ile Glu Asn His Gly Ile Glu Asp Glu
 260 265 270
 Leu Pro Asn Tyr Cys Phe Pro Leu Asn Gly Ile Gly Pro Gly His Thr
 275 280 285
 Tyr Gln Gly Ile Lys Lys Val Lys Thr Asp Asp Thr Asn Gly Trp Glu
 290 295 300
 Lys Asp Ala Asn Val Ala Pro Ala Asn Glu Ile Thr Ile Gly Asn Asn
 305 310 315 320
 Leu Ala Met

<210> 37
 <211> 315
 <212> PRT
 <213> Human adenovirus type 7

<400> 37

```

Asn Thr Ser Gln Trp Ile Val Thr Ala Gly Glu Glu Arg Ala Val Thr
1          5          10          15

Thr Thr Thr Asn Thr Phe Gly Ile Ala Ser Met Lys Gly Asp Asn Ile
20          25          30

Thr Lys Glu Gly Leu Glu Ile Gly Lys Asp Ile Thr Ala Asp Asn Lys
35          40          45

Pro Ile Tyr Ala Asp Lys Thr Tyr Gln Pro Glu Pro Gln Val Gly Glu
50          55          60

Glu Ser Trp Thr Asp Thr Asp Gly Thr Asn Glu Lys Phe Gly Gly Arg
65          70          75          80

Ala Leu Lys Pro Ala Thr Lys Met Lys Pro Cys Tyr Gly Ser Phe Ala
85          90          95

Arg Pro Thr Asn Ile Lys Gly Gly Gln Ala Lys Asn Arg Lys Val Lys
100         105         110

Pro Thr Glu Gly Asp Val Glu Thr Glu Glu Pro Asp Ile Asp Met Glu
115         120         125

Phe Phe Asp Gly Arg Glu Ala Ala Asp Ala Phe Ser Pro Glu Ile Val
130         135         140

Leu Tyr Thr Glu Asn Val Asn Leu Glu Thr Pro Asp Ser His Val Val
145         150         155         160

Tyr Lys Pro Gly Thr Ser Asp Asp Asn Ser His Ala Asn Leu Gly Gln
165         170         175

Gln Ala Met Pro Asn Arg Pro Asn Tyr Ile Gly Phe Arg Asp Asn Phe
180         185         190

Val Gly Leu Met Tyr Tyr Asn Ser Thr Gly Asn Met Gly Val Leu Ala
195         200         205

Gly Gln Ala Ser Gln Leu Asn Ala Val Val Asp Leu Gln Asp Arg Asn
210         215         220

Thr Glu Leu Ser Tyr Gln Leu Leu Leu Asp Ser Leu Gly Asp Arg Thr
225         230         235         240

Arg Tyr Phe Ser Met Trp Asn Gln Ala Val Asp Ser Tyr Asp Pro Asp
245         250         255

Val Arg Ile Ile Glu Asn His Gly Ile Glu Asp Glu Leu Pro Asn Tyr
260         265         270

Cys Phe Pro Leu Asp Gly Ile Gly Pro Ala Lys Thr Tyr Gln Gly Ile
275         280         285

Lys Ser Lys Asp Asn Gly Trp Glu Lys Asp Asp Asn Val Ser Lys Ser
290         295         300

Asn Glu Ile Ala Ile Gly Asn Asn Gln Ala Met
305         310         315

```

<210> 38
 <211> 345
 <212> PRT
 <213> Human adenovirus type 2

<400> 38

```

Asn Ser Cys Glu Trp Glu Gln Thr Glu Asp Ser Gly Arg Ala Val Ala
1      5      10      15
Glu Asp Glu Glu Glu Glu Asp Glu Asp Glu Glu Glu Glu Glu Glu
20      25      30
Gln Asn Ala Arg Asp Gln Ala Thr Lys Lys Thr His Val Tyr Ala Gln
35      40      45
Ala Pro Leu Ser Gly Glu Thr Leu Thr Lys Ser Gly Leu Gln Ile Gly
50      55      60
Ser Lys Asn Ala Glu Thr Gln Ala Lys Pro Val Tyr Ala Asp Pro Ser
65      70      75      80
Tyr Gln Pro Glu Pro Gln Ile Gly Glu Ser Gln Trp Asn Glu Ala Asp
85      90      95
Ala Asn Ala Ala Gly Gly Arg Val Leu Lys Lys Thr Thr Pro Met Lys
100     105     110
Pro Tyr Gly Ser Tyr Ala Arg Pro Thr Asn Pro Phe Gly Gly Gln Ser
115     120     125
Val Leu Val Pro Asp Glu Lys Gly Val Pro Leu Pro Lys Val Asp Leu
130     135     140
Gln Phe Phe Ser Asn Thr Thr Ser Leu Asn Asp Arg Gln Gly Asn Ala
145     150     155     160
Thr Lys Pro Lys Val Val Leu Tyr Ser Glu Asp Val Asn Met Glu Thr
165     170     175
Pro Asp Thr His Leu Ser Tyr Lys Pro Gly Lys Gly Asp Glu Asn Ser
180     185     190
Lys Ala Met Leu Gly Gln Gln Ser Met Pro Asn Arg Pro Asn Tyr Ile
195     200     205
Ala Phe Arg Asp Asn Phe Ile Gly Leu Met Tyr Tyr Asn Ser Thr Gly
210     215     220
Asn Met Gly Val Leu Ala Gly Gln Ala Ser Gln Leu Asn Ala Val Val
225     230     235     240
Asp Leu Gln Asp Arg Asn Thr Glu Leu Ser Tyr Gln Leu Leu Leu Asp
245     250     255
Ser Ile Gly Asp Arg Thr Arg Tyr Phe Ser Met Trp Asn Gln Ala Val
260     265     270
Asp Ser Tyr Asp Pro Asp Val Arg Ile Ile Glu Asn His Gly Thr Glu
275     280     285

```

Asp Glu Leu Pro Asn Tyr Cys Phe Pro Leu Gly Gly Ile Gly Val Thr
 290 295 300
 Asp Thr Tyr Gln Ala Ile Lys Ala Asn Gly Asn Gly Ser Gly Asp Asn
 305 310 315 320
 Gly Asp Thr Thr Trp Thr Lys Asp Glu Thr Phe Ala Thr Arg Asn Glu
 325 330 335
 Ile Gly Val Gly Asn Asn Phe Ala Met
 340 345

<210> 39
 <211> 183
 <212> PRT
 <213> human adenovirus protein

<400> 39

Thr Leu Trp Thr Thr Pro Asp Pro Ser Pro Asn Cys Arg Ile His Ser
 1 5 10 15
 Asp Asn Asp Cys Lys Phe Thr Leu Val Leu Thr Lys Cys Gly Ser Gln
 20 25 30
 Val Leu Ala Thr Val Ala Ala Leu Ala Val Ser Gly Asp Leu Ser Ser
 35 40 45
 Met Thr Gly Thr Val Ala Ser Val Ser Ile Phe Leu Arg Phe Asp Gln
 50 55 60
 Asn Gly Val Leu Met Glu Asn Ser Ser Leu Lys Lys His Tyr Trp Asn
 65 70 75 80
 Phe Arg Asn Gly Asn Ser Thr Asn Ala Asn Pro Tyr Thr Asn Ala Val
 85 90 95
 Gly Phe Met Pro Asn Leu Leu Ala Tyr Pro Lys Thr Gln Ser Gln Thr
 100 105 110
 Ala Lys Asn Asn Ile Val Ser Gln Val Tyr Leu His Gly Asp Lys Thr
 115 120 125
 Lys Pro Met Ile Leu Thr Ile Thr Leu Asn Gly Thr Ser Glu Ser Thr
 130 135 140
 Glu Thr Ser Glu Val Ser Thr Tyr Ser Met Ser Phe Thr Trp Ser Trp
 145 150 155 160
 Glu Ser Gly Lys Tyr Thr Thr Glu Thr Phe Ala Thr Asn Ser Tyr Thr
 165 170 175
 Phe Ser Tyr Ile Ala Gln Glu
 180

<210> 40
 <211> 182
 <212> PRT
 <213> human adenovirus protein

<400> 40

```

Thr Leu Trp Thr Thr Pro Ala Pro Ser Pro Asn Cys Arg Leu Asn Ala
1      5      10      15

Glu Lys Asp Ala Lys Leu Thr Leu Val Leu Thr Lys Cys Gly Ser Gln
      20      25      30

Ile Leu Ala Thr Val Ser Val Leu Ala Val Lys Gly Ser Leu Ala Pro
      35      40      45

Ile Ser Gly Thr Val Gln Ser Ala His Leu Ile Ile Arg Phe Asp Glu
      50      55      60

Asn Gly Val Leu Leu Asn Asn Ser Phe Leu Asp Pro Glu Tyr Trp Asn
      65      70      75      80

Phe Arg Asn Gly Asp Leu Thr Glu Gly Thr Ala Tyr Thr Asn Ala Val
      85      90      95

Gly Phe Met Pro Asn Leu Ser Ala Tyr Pro Lys Ser His Gly Lys Thr
      100      105      110

Ala Lys Ser Asn Ile Val Ser Gln Val Tyr Leu Asn Gly Asp Lys Thr
      115      120      125

Lys Pro Val Thr Leu Thr Ile Thr Leu Asn Gly Thr Gln Glu Thr Gly
      130      135      140

Asp Thr Thr Pro Ser Ala Tyr Ser Met Ser Phe Ser Trp Asp Trp Ser
      145      150      155      160

Gly His Asn Tyr Ile Asn Glu Ile Phe Ala Thr Ser Ser Tyr Thr Phe
      165      170      175

Ser Tyr Ile Ala Gln Glu
      180

```

<210> 41

<211> 338

<212> PRT

<213> human adenovirus protein

<400> 41

```

Ala Pro Lys Gly Ala Pro Asn Pro Cys Glu Trp Asp Glu Ala Ala Thr
1      5      10      15

Ala Leu Glu Ile Asn Leu Glu Glu Glu Asp Asp Asp Asn Glu Asp Glu
      20      25      30

Val Asp Glu Gln Ala Glu Gln Gln Lys Thr His Val Phe Gly Gln Ala
      35      40      45

Pro Tyr Ser Gly Ile Asn Ile Thr Lys Glu Gly Ile Gln Ile Gly Val
      50      55      60

Glu Gly Gln Thr Pro Lys Tyr Ala Asp Lys Thr Phe Gln Pro Glu Pro
      65      70      75      80

```

Gln Ile Gly Glu Ser Gln⁷ Trp Tyr Glu Thr Glu Ile Asn His Ala Ala
 85 90 95
 Gly Arg Val Leu Lys Lys Thr Thr Pro Met Lys Pro Cys Tyr Gly Ser
 100 105 110
 Tyr Ala Lys Pro Thr Asn Glu Asn Gly Gly Gln Gly Ile Leu Val Lys
 115 120 125
 Gln Gln Asn Gly Lys Leu Glu Ser Gln Val Glu Met Gln Phe Phe Ser
 130 135 140
 Thr Thr Glu Ala Thr Ala Gly Asn Gly Asp Asn Leu Thr Pro Lys Val
 145 150 155 160
 Val Leu Tyr Ser Glu Asp Val Asp Ile Glu Thr Pro Asp Thr His Ile
 165 170 175
 Ser Tyr Met Pro Thr Ile Lys Glu Gly Asn Ser Arg Glu Leu Met Gly
 180 185 190
 Gln Gln Ser Met Pro Asn Arg Pro Asn Tyr Ile Ala Phe Arg Asp Asn
 195 200 205
 Phe Ile Gly Leu Met Tyr Tyr Asn Ser Thr Gly Asn Met Gly Val Leu
 210 215 220
 Ala Gly Gln Ala Ser Gln Leu Asn Ala Val Val Asp Leu Gln Asp Arg
 225 230 235 240
 Asn Thr Glu Leu Ser Tyr Gln Leu Leu Leu Asp Ser Ile Gly Asp Arg
 245 250 255
 Thr Arg Tyr Phe Ser Met Trp Asn Gln Ala Val Asp Ser Tyr Asp Pro
 260 265 270
 Asp Val Arg Ile Ile Glu Asn His Gly Thr Glu Asp Glu Leu Pro Asn
 275 280 285
 Tyr Cys Phe Pro Leu Gly Gly Val Ile Asn Thr Glu Thr Leu Thr Lys
 290 295 300
 Val Lys Pro Lys Thr Gly Gln Glu Asn Gly Trp Glu Lys Asp Ala Thr
 305 310 315 320
 Glu Phe Ser Asp Lys Asn Glu Ile Arg Val Gly Asn Asn Phe Ala Met
 325 330 335
 Glu Ile